

Tennessee Together Student Survey Results Compendium

2018-2019

Detailed Tables

TN  **TOGETHER**

TN Department of
**Mental Health &
Substance Abuse Services**


EVALUATION • MANAGEMENT • TRAINING

This project is funded under a contract with the State of Tennessee, Department of Mental Health and Substance Abuse Services. The survey and analysis were conducted by Evaluation, Management, and Training (EMT) Associates, Inc.

This page was intentionally left blank to
optimize double-sided printing.

Contents

Introduction	10
Table A. Unweighted and Weighted Sample by County	11
Demographic Characteristics of the Respondent Sample.....	12
Table D.1. Age of Sample	12
Table D.2. Grade Level of Sample.....	12
Table D.3. Gender of Sample	12
Table D.4. Race and Ethnicity of Sample	12
Table D.5. Living Situation of Sample.....	12
Table D.6. School Grades of Sample	12
Statewide Results Summary.....	13
Table TN.1. Lifetime Prescription Drug Misuse	13
Table TN.2. Lifetime Alcohol Use.....	13
Table TN.3. Lifetime Binge Drinking	14
Table TN.4. Lifetime Cigarette Use.....	14
Table TN.5. Lifetime Electronic Cigarette Use.....	15
Table TN.6. Lifetime Smokeless Tobacco Use.....	15
Table TN.7. Lifetime Marijuana Use.....	16
Table TN.8. Lifetime Methamphetamines Use.....	16
Table TN.9. Lifetime Other Illegal Drug Use	17
Table TN.10. Age of First Prescription Drug Misuse.....	18
Table TN.11. Age of First Alcohol Use.....	19
Table TN.12. Age of First Cigarette Use	20
Table TN.13. Age of First Electronic Cigarette Use	21
Table TN.14. Age of First Marijuana Use.....	22
Table TN.15. Past Month Prescription Drug Misuse.....	23
Table TN.16. Past Month Alcohol Use	23
Table TN.17. Past Month Binge Drinking.....	24
Table TN.18. Past Month Cigarette Use	24
Table TN.19. Past Month Electronic Cigarette Use	25
Table TN.20. Past Month Smokeless Tobacco Use	25
Table TN.21. Past Month Marijuana Use	26
Table TN.22. Past Month Cocaine Use.....	26
Table TN.23. Past Month Inhalants Use	27
Table TN.24. Past Month Hallucinogens Use	27
Table TN.25. Past Month Heroin Use	28
Table TN.26. Past Month Steroids Use.....	28
Table TN.27. Past Month Ecstasy Use	29
Table TN.28. Past Month Methamphetamines Use.....	29
Table TN.29. Availability of Prescription Drugs.....	30
Table TN.30. Availability of Alcohol.....	30
Table TN.31. Availability of Tobacco.....	31
Table TN.32. Availability of Marijuana.....	31
Table TN.33. Availability of Prescription Drugs.....	32
Table TN.34. Availability of Alcohol.....	32
Table TN.35. Availability of Prescription Drugs – Sources	33
Table TN.36. Availability of Alcohol – Sources	34
Table TN.37. Asked to Show Proof of Age to Buy Alcohol	35
Table TN.38. Ability to Purchase Alcohol.....	35
Table TN.39. Ever Ridden with a Prescription Drugs Impaired Driver.....	36
Table TN.40. Frequency Riding with a Prescription Drugs Impaired Driver	36
Table TN.41. Ever Ridden with an Alcohol or Drug Impaired Driver	37
Table TN.42. Frequency Riding with an Alcohol or Drug Impaired Driver	37
Table TN.43. Parents Feel Misusing Prescription Drugs is Wrong.....	38
Table TN.44. Parents Feel Drinking Alcohol is Wrong.....	38
Table TN.45. Parents Feel Regularly Drinking Alcohol is Wrong	39

Table TN.46.	Parents Feel Smoking Tobacco is Wrong.....	39
Table TN.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong.....	40
Table TN.48.	Parents Feel Using Electronic Cigarettes is Wrong.....	40
Table TN.49.	Parents Feel Smoking Marijuana is Wrong.....	41
Table TN.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong.....	41
Table TN.51.	Friends Feel Misusing Prescription Drugs is Wrong.....	42
Table TN.52.	Friends Feel Drinking Alcohol is Wrong.....	42
Table TN.53.	Friends Feel Smoking Tobacco is Wrong.....	43
Table TN.54.	Friends Feel Using Electronic Cigarettes is Wrong.....	43
Table TN.55.	Friends Feel Smoking Marijuana is Wrong.....	44
Table TN.56.	Friends Misused Prescription Drugs.....	44
Table TN.57.	Friends Tried Alcohol.....	45
Table TN.58.	Friends Binge Drinking.....	45
Table TN.59.	Friends Smoked Cigarettes.....	46
Table TN.60.	Friends Used Electronic Cigarettes.....	46
Table TN.61.	Friends Used Marijuana.....	47
Table TN.62.	Perceived Risk of Misusing Prescription Drugs.....	47
Table TN.63.	Perceived Risk of Regularly Drinking Alcohol.....	48
Table TN.64.	Perceived Risk of Binge Drinking.....	48
Table TN.65.	Perceived Risk of Smoking Cigarettes.....	49
Table TN.66.	Perceived Risk of Using Electronic Cigarettes.....	49
Table TN.67.	Perceived Risk of Trying Marijuana.....	50
Table TN.68.	Perceived Risk of Regularly Smoking Marijuana.....	50
Table TN.69.	Wrong to Misuse Prescription Drugs.....	51
Table TN.70.	Wrong to Regularly Drink Alcohol.....	51
Table TN.71.	Wrong to Smoke Tobacco.....	52
Table TN.72.	Wrong to Use Electronic Cigarettes.....	52
Table TN.73.	Wrong to Smoke Marijuana.....	53
Table TN.74.	Parents Talked about Prescription Drug Misuse.....	53
Table TN.75.	Parents Talked about Alcohol and Other Drugs (ATOD).....	54
Table TN.76.	Exposure to Prescription Drug Misuse Media Messaging.....	54

Behavioral Health Planning Region 1 Results Summary..... 55

Table BH1.1.	Lifetime Prescription Drug Misuse.....	55
Table BH1.2.	Lifetime Alcohol Use.....	55
Table BH1.3.	Lifetime Binge Drinking.....	56
Table BH1.4.	Lifetime Cigarette Use.....	56
Table BH1.5.	Lifetime Electronic Cigarette Use.....	57
Table BH1.6.	Lifetime Smokeless Tobacco Use.....	57
Table BH1.7.	Lifetime Marijuana Use.....	58
Table BH1.8.	Lifetime Methamphetamines Use.....	58
Table BH1.9.	Lifetime Other Illegal Drug Use.....	59
Table BH1.10.	Age of First Prescription Drug Misuse.....	60
Table BH1.11.	Age of First Alcohol Use.....	61
Table BH1.12.	Age of First Cigarette Use.....	62
Table BH1.13.	Age of First Electronic Cigarette Use.....	63
Table BH1.14.	Age of First Marijuana Use.....	64
Table BH1.15.	Past Month Prescription Drug Misuse.....	65
Table BH1.16.	Past Month Alcohol Use.....	65
Table BH1.17.	Past Month Binge Drinking.....	66
Table BH1.18.	Past Month Cigarette Use.....	66
Table BH1.19.	Past Month Electronic Cigarette Use.....	67
Table BH1.20.	Past Month Smokeless Tobacco Use.....	67
Table BH1.21.	Past Month Marijuana Use.....	68
Table BH1.22.	Past Month Cocaine Use.....	68
Table BH1.23.	Past Month Inhalants Use.....	69
Table BH1.24.	Past Month Hallucinogens Use.....	69
Table BH1.25.	Past Month Heroin Use.....	70
Table BH1.26.	Past Month Steroids Use.....	70
Table BH1.27.	Past Month Ecstasy Use.....	71

Table BH1.28.	Past Month Methamphetamines Use	71
Table BH1.29.	Availability of Prescription Drugs	72
Table BH1.30.	Availability of Alcohol	72
Table BH1.31.	Availability of Tobacco	73
Table BH1.32.	Availability of Marijuana	73
Table BH1.33.	Availability of Prescription Drugs	74
Table BH1.34.	Availability of Alcohol	74
Table BH1.35.	Availability of Prescription Drugs – Sources	75
Table BH1.36.	Availability of Alcohol – Sources	76
Table BH1.37.	Asked to Show Proof of Age to Buy Alcohol	77
Table BH1.38.	Ability to Purchase Alcohol	77
Table BH1.39.	Ever Ridden with a Prescription Drugs Impaired Driver	78
Table BH1.40.	Frequency Riding with a Prescription Drugs Impaired Driver	78
Table BH1.41.	Ever Ridden with an Alcohol or Drug Impaired Driver	79
Table BH1.42.	Frequency Riding with an Alcohol or Drug Impaired Driver	79
Table BH1.43.	Parents Feel Misusing Prescription Drugs is Wrong	80
Table BH1.44.	Parents Feel Drinking Alcohol is Wrong	80
Table BH1.45.	Parents Feel Regularly Drinking Alcohol is Wrong	81
Table BH1.46.	Parents Feel Smoking Tobacco is Wrong	81
Table BH1.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong	82
Table BH1.48.	Parents Feel Using Electronic Cigarettes is Wrong	82
Table BH1.49.	Parents Feel Smoking Marijuana is Wrong	83
Table BH1.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong	83
Table BH1.51.	Friends Feel Misusing Prescription Drugs is Wrong	84
Table BH1.52.	Friends Feel Drinking Alcohol is Wrong	84
Table BH1.53.	Friends Feel Smoking Tobacco is Wrong	85
Table BH1.54.	Friends Feel Using Electronic Cigarettes is Wrong	85
Table BH1.55.	Friends Feel Smoking Marijuana is Wrong	86
Table BH1.56.	Friends Misused Prescription Drugs	86
Table BH1.57.	Friends Tried Alcohol	87
Table BH1.58.	Friends Binge Drinking	87
Table BH1.59.	Friends Smoked Cigarettes	88
Table BH1.60.	Friends Used Electronic Cigarettes	88
Table BH1.61.	Friends Used Marijuana	89
Table BH1.62.	Perceived Risk of Misusing Prescription Drugs	89
Table BH1.63.	Perceived Risk of Regularly Drinking Alcohol	90
Table BH1.64.	Perceived Risk of Binge Drinking	90
Table BH1.65.	Perceived Risk of Smoking Cigarettes	91
Table BH1.66.	Perceived Risk of Using Electronic Cigarettes	91
Table BH1.67.	Perceived Risk of Trying Marijuana	92
Table BH1.68.	Perceived Risk of Regularly Smoking Marijuana	92
Table BH1.69.	Wrong to Misuse Prescription Drugs	93
Table BH1.70.	Wrong to Regularly Drink Alcohol	93
Table BH1.71.	Wrong to Smoke Tobacco	94
Table BH1.72.	Wrong to Use Electronic Cigarettes	94
Table BH1.73.	Wrong to Smoke Marijuana	95
Table BH1.74.	Parents Talked about Prescription Drug Misuse	95
Table BH1.75.	Parents Talked about Alcohol and Other Drugs (ATOD)	96
Table BH1.76.	Exposure to Prescription Drug Misuse Media Messaging	96

Behavioral Health Planning Region 2 Results Summary **97**

Table BH2.1.	Lifetime Prescription Drug Misuse	97
Table BH2.2.	Lifetime Alcohol Use	97
Table BH2.3.	Lifetime Binge Drinking	98
Table BH2.4.	Lifetime Cigarette Use	98
Table BH2.5.	Lifetime Electronic Cigarette Use	99
Table BH2.6.	Lifetime Smokeless Tobacco Use	99
Table BH2.7.	Lifetime Marijuana Use	100
Table BH2.8.	Lifetime Methamphetamines Use	100
Table BH2.9.	Lifetime Other Illegal Drug Use	101

Table BH2.10.	Age of First Prescription Drug Misuse	102
Table BH2.11.	Age of First Alcohol Use	103
Table BH2.12.	Age of First Cigarette Use	104
Table BH2.13.	Age of First Electronic Cigarette Use	105
Table BH2.14.	Age of First Marijuana Use	106
Table BH2.15.	Past Month Prescription Drug Misuse	107
Table BH2.16.	Past Month Alcohol Use	107
Table BH2.17.	Past Month Binge Drinking	108
Table BH2.18.	Past Month Cigarette Use	108
Table BH2.19.	Past Month Electronic Cigarette Use	109
Table BH2.20.	Past Month Smokeless Tobacco Use	109
Table BH2.21.	Past Month Marijuana Use	110
Table BH2.22.	Past Month Cocaine Use	110
Table BH2.23.	Past Month Inhalants Use	111
Table BH2.24.	Past Month Hallucinogens Use	111
Table BH2.25.	Past Month Heroin Use	112
Table BH2.26.	Past Month Steroids Use	112
Table BH2.27.	Past Month Ecstasy Use	113
Table BH2.28.	Past Month Methamphetamines Use	113
Table BH2.29.	Availability of Prescription Drugs	114
Table BH2.30.	Availability of Alcohol	114
Table BH2.31.	Availability of Tobacco	115
Table BH2.32.	Availability of Marijuana	115
Table BH2.33.	Availability of Prescription Drugs	116
Table BH2.34.	Availability of Alcohol	116
Table BH2.35.	Availability of Prescription Drugs – Sources	117
Table BH2.36.	Availability of Alcohol – Sources	118
Table BH2.37.	Asked to Show Proof of Age to Buy Alcohol	119
Table BH2.38.	Ability to Purchase Alcohol	119
Table BH2.39.	Ever Ridden with a Prescription Drugs Impaired Driver	120
Table BH2.40.	Frequency Riding with a Prescription Drugs Impaired Driver	120
Table BH2.41.	Ever Ridden with an Alcohol or Drug Impaired Driver	121
Table BH2.42.	Frequency Riding with an Alcohol or Drug Impaired Driver	121
Table BH2.43.	Parents Feel Misusing Prescription Drugs is Wrong	122
Table BH2.44.	Parents Feel Drinking Alcohol is Wrong	122
Table BH2.45.	Parents Feel Regularly Drinking Alcohol is Wrong	123
Table BH2.46.	Parents Feel Smoking Tobacco is Wrong	123
Table BH2.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong	124
Table BH2.48.	Parents Feel Using Electronic Cigarettes is Wrong	124
Table BH2.49.	Parents Feel Smoking Marijuana is Wrong	125
Table BH2.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong	125
Table BH2.51.	Friends Feel Misusing Prescription Drugs is Wrong	126
Table BH2.52.	Friends Feel Drinking Alcohol is Wrong	126
Table BH2.53.	Friends Feel Smoking Tobacco is Wrong	127
Table BH2.54.	Friends Feel Using Electronic Cigarettes is Wrong	127
Table BH2.55.	Friends Feel Smoking Marijuana is Wrong	128
Table BH2.56.	Friends Misused Prescription Drugs	128
Table BH2.57.	Friends Tried Alcohol	129
Table BH2.58.	Friends Binge Drinking	129
Table BH2.59.	Friends Smoked Cigarettes	130
Table BH2.60.	Friends Used Electronic Cigarettes	130
Table BH2.61.	Friends Used Marijuana	131
Table BH2.62.	Perceived Risk of Misusing Prescription Drugs	131
Table BH2.63.	Perceived Risk of Regularly Drinking Alcohol	132
Table BH2.64.	Perceived Risk of Binge Drinking	132
Table BH2.65.	Perceived Risk of Smoking Cigarettes	133
Table BH2.66.	Perceived Risk of Using Electronic Cigarettes	133
Table BH2.67.	Perceived Risk of Trying Marijuana	134
Table BH2.68.	Perceived Risk of Regularly Smoking Marijuana	134
Table BH2.69.	Wrong to Misuse Prescription Drugs	135

Table BH2.70.	Wrong to Regularly Drink Alcohol	135
Table BH2.71.	Wrong to Smoke Tobacco	136
Table BH2.72.	Wrong to Use Electronic Cigarettes	136
Table BH2.73.	Wrong to Smoke Marijuana	137
Table BH2.74.	Parents Talked about Prescription Drug Misuse	137
Table BH2.75.	Parents Talked about Alcohol and Other Drugs (ATOD)	138
Table BH2.76.	Exposure to Prescription Drug Misuse Media Messaging.....	138

Behavioral Health Planning Region 3 Results Summary..... 139

Table BH3.1.	Lifetime Prescription Drug Misuse	139
Table BH3.2.	Lifetime Alcohol Use.....	139
Table BH3.3.	Lifetime Binge Drinking	140
Table BH3.4.	Lifetime Cigarette Use	140
Table BH3.5.	Lifetime Electronic Cigarette Use.....	141
Table BH3.6.	Lifetime Smokeless Tobacco Use.....	141
Table BH3.7.	Lifetime Marijuana Use.....	142
Table BH3.8.	Lifetime Methamphetamines Use.....	142
Table BH3.9.	Lifetime Other Illegal Drug Use	143
Table BH3.10.	Age of First Prescription Drug Misuse.....	144
Table BH3.11.	Age of First Alcohol Use	145
Table BH3.12.	Age of First Cigarette Use	146
Table BH3.13.	Age of First Electronic Cigarette Use	147
Table BH3.14.	Age of First Marijuana Use	148
Table BH3.15.	Past Month Prescription Drug Misuse	149
Table BH3.16.	Past Month Alcohol Use	149
Table BH3.17.	Past Month Binge Drinking.....	150
Table BH3.18.	Past Month Cigarette Use	150
Table BH3.19.	Past Month Electronic Cigarette Use	151
Table BH3.20.	Past Month Smokeless Tobacco Use	151
Table BH3.21.	Past Month Marijuana Use	152
Table BH3.22.	Past Month Cocaine Use.....	152
Table BH3.23.	Past Month Inhalants Use	153
Table BH3.24.	Past Month Hallucinogens Use	153
Table BH3.25.	Past Month Heroin Use	154
Table BH3.26.	Past Month Steroids Use.....	154
Table BH3.27.	Past Month Ecstasy Use	155
Table BH3.28.	Past Month Methamphetamines Use	155
Table BH3.29.	Availability of Prescription Drugs.....	156
Table BH3.30.	Availability of Alcohol.....	156
Table BH3.31.	Availability of Tobacco.....	157
Table BH3.32.	Availability of Marijuana.....	157
Table BH3.33.	Availability of Prescription Drugs.....	158
Table BH3.34.	Availability of Alcohol.....	158
Table BH3.35.	Availability of Prescription Drugs – Sources	159
Table BH3.36.	Availability of Alcohol – Sources	160
Table BH3.37.	Asked to Show Proof of Age to Buy Alcohol	161
Table BH3.38.	Ability to Purchase Alcohol.....	161
Table BH3.39.	Ever Ridden with a Prescription Drugs Impaired Driver.....	162
Table BH3.40.	Frequency Riding with a Prescription Drugs Impaired Driver	162
Table BH3.41.	Ever Ridden with an Alcohol or Drug Impaired Driver.....	163
Table BH3.42.	Frequency Riding with an Alcohol or Drug Impaired Driver	163
Table BH3.43.	Parents Feel Misusing Prescription Drugs is Wrong.....	164
Table BH3.44.	Parents Feel Drinking Alcohol is Wrong.....	164
Table BH3.45.	Parents Feel Regularly Drinking Alcohol is Wrong	165
Table BH3.46.	Parents Feel Smoking Tobacco is Wrong.....	165
Table BH3.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong	166
Table BH3.48.	Parents Feel Using Electronic Cigarettes is Wrong	166
Table BH3.49.	Parents Feel Smoking Marijuana is Wrong.....	167
Table BH3.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong	167
Table BH3.51.	Friends Feel Misusing Prescription Drugs is Wrong	168

Table BH3.52.	Friends Feel Drinking Alcohol is Wrong	168
Table BH3.53.	Friends Feel Smoking Tobacco is Wrong	169
Table BH3.54.	Friends Feel Using Electronic Cigarettes is Wrong.....	169
Table BH3.55.	Friends Feel Smoking Marijuana is Wrong	170
Table BH3.56.	Friends Misused Prescription Drugs	170
Table BH3.57.	Friends Tried Alcohol.....	171
Table BH3.58.	Friends Binge Drinking	171
Table BH3.59.	Friends Smoked Cigarettes	172
Table BH3.60.	Friends Used Electronic Cigarettes	172
Table BH3.61.	Friends Used Marijuana	173
Table BH3.62.	Perceived Risk of Misusing Prescription Drugs	173
Table BH3.63.	Perceived Risk of Regularly Drinking Alcohol	174
Table BH3.64.	Perceived Risk of Binge Drinking	174
Table BH3.65.	Perceived Risk of Smoking Cigarettes	175
Table BH3.66.	Perceived Risk of Using Electronic Cigarettes	175
Table BH3.67.	Perceived Risk of Trying Marijuana.....	176
Table BH3.68.	Perceived Risk of Regularly Smoking Marijuana	176
Table BH3.69.	Wrong to Misuse Prescription Drugs	177
Table BH3.70.	Wrong to Regularly Drink Alcohol	177
Table BH3.71.	Wrong to Smoke Tobacco	178
Table BH3.72.	Wrong to Use Electronic Cigarettes	178
Table BH3.73.	Wrong to Smoke Marijuana	179
Table BH3.74.	Parents Talked about Prescription Drug Misuse	179
Table BH3.75.	Parents Talked about Alcohol and Other Drugs (ATOD)	180
Table BH3.76.	Exposure to Prescription Drug Misuse Media Messaging.....	180

Behavioral Health Planning Region 5 Results Summary..... 181

Table BH5.1.	Lifetime Prescription Drug Misuse	181
Table BH5.2.	Lifetime Alcohol Use.....	181
Table BH5.3.	Lifetime Binge Drinking	182
Table BH5.4.	Lifetime Cigarette Use.....	182
Table BH5.5.	Lifetime Electronic Cigarette Use	183
Table BH5.6.	Lifetime Smokeless Tobacco Use	183
Table BH5.7.	Lifetime Marijuana Use.....	184
Table BH5.8.	Lifetime Methamphetamines Use.....	184
Table BH5.9.	Lifetime Other Illegal Drug Use	185
Table BH5.10.	Age of First Prescription Drug Misuse	186
Table BH5.11.	Age of First Alcohol Use	187
Table BH5.12.	Age of First Cigarette Use	188
Table BH5.13.	Age of First Electronic Cigarette Use	189
Table BH5.14.	Age of First Marijuana Use	190
Table BH5.15.	Past Month Prescription Drug Misuse	191
Table BH5.16.	Past Month Alcohol Use	191
Table BH5.17.	Past Month Binge Drinking	192
Table BH5.18.	Past Month Cigarette Use	192
Table BH5.19.	Past Month Electronic Cigarette Use	193
Table BH5.20.	Past Month Smokeless Tobacco Use	193
Table BH5.21.	Past Month Marijuana Use	194
Table BH5.22.	Past Month Cocaine Use.....	194
Table BH5.23.	Past Month Inhalants Use	195
Table BH5.24.	Past Month Hallucinogens Use	195
Table BH5.25.	Past Month Heroin Use	196
Table BH5.26.	Past Month Steroids Use.....	196
Table BH5.27.	Past Month Ecstasy Use	197
Table BH5.28.	Past Month Methamphetamines Use	197
Table BH5.29.	Availability of Prescription Drugs	198
Table BH5.30.	Availability of Alcohol.....	198
Table BH5.31.	Availability of Tobacco	199
Table BH5.32.	Availability of Marijuana.....	199
Table BH5.33.	Availability of Prescription Drugs	200

Table BH5.34.	Availability of Alcohol.....	200
Table BH5.35.	Availability of Prescription Drugs – Sources	201
Table BH5.36.	Availability of Alcohol – Sources	202
Table BH5.37.	Asked to Show Proof of Age to Buy Alcohol	203
Table BH5.38.	Ability to Purchase Alcohol.....	203
Table BH5.39.	Ever Ridden with a Prescription Drugs Impaired Driver.....	204
Table BH5.40.	Frequency Riding with a Prescription Drugs Impaired Driver	204
Table BH5.41.	Ever Ridden with an Alcohol or Drug Impaired Driver.....	205
Table BH5.42.	Frequency Riding with an Alcohol or Drug Impaired Driver	205
Table BH5.43.	Parents Feel Misusing Prescription Drugs is Wrong.....	206
Table BH5.44.	Parents Feel Drinking Alcohol is Wrong.....	206
Table BH5.45.	Parents Feel Regularly Drinking Alcohol is Wrong	207
Table BH5.46.	Parents Feel Smoking Tobacco is Wrong.....	207
Table BH5.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong	208
Table BH5.48.	Parents Feel Using Electronic Cigarettes is Wrong	208
Table BH5.49.	Parents Feel Smoking Marijuana is Wrong.....	209
Table BH5.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong	209
Table BH5.51.	Friends Feel Misusing Prescription Drugs is Wrong	210
Table BH5.52.	Friends Feel Drinking Alcohol is Wrong	210
Table BH5.53.	Friends Feel Smoking Tobacco is Wrong	211
Table BH5.54.	Friends Feel Using Electronic Cigarettes is Wrong.....	211
Table BH5.55.	Friends Feel Smoking Marijuana is Wrong	212
Table BH5.56.	Friends Misused Prescription Drugs	212
Table BH5.57.	Friends Tried Alcohol.....	213
Table BH5.58.	Friends Binge Drinking	213
Table BH5.59.	Friends Smoked Cigarettes	214
Table BH5.60.	Friends Used Electronic Cigarettes.....	214
Table BH5.61.	Friends Used Marijuana	215
Table BH5.62.	Perceived Risk of Misusing Prescription Drugs	215
Table BH5.63.	Perceived Risk of Regularly Drinking Alcohol	216
Table BH5.64.	Perceived Risk of Binge Drinking	216
Table BH5.65.	Perceived Risk of Smoking Cigarettes	217
Table BH5.66.	Perceived Risk of Using Electronic Cigarettes	217
Table BH5.67.	Perceived Risk of Trying Marijuana.....	218
Table BH5.68.	Perceived Risk of Regularly Smoking Marijuana	218
Table BH5.69.	Wrong to Misuse Prescription Drugs.....	219
Table BH5.70.	Wrong to Regularly Drink Alcohol	219
Table BH5.71.	Wrong to Smoke Tobacco	220
Table BH5.72.	Wrong to Use Electronic Cigarettes	220
Table BH5.73.	Wrong to Smoke Marijuana.....	221
Table BH5.74.	Parents Talked about Prescription Drug Misuse.....	221
Table BH5.75.	Parents Talked about Alcohol and Other Drugs (ATOD)	222
Table BH5.76.	Exposure to Prescription Drug Misuse Media Messaging.....	222

Behavioral Health Planning Region 6 Results Summary..... 223

Table BH6.1.	Lifetime Prescription Drug Misuse	223
Table BH6.2.	Lifetime Alcohol Use.....	223
Table BH6.3.	Lifetime Binge Drinking	224
Table BH6.4.	Lifetime Cigarette Use.....	224
Table BH6.5.	Lifetime Electronic Cigarette Use.....	225
Table BH6.6.	Lifetime Smokeless Tobacco Use.....	225
Table BH6.7.	Lifetime Marijuana Use.....	226
Table BH6.8.	Lifetime Methamphetamines Use.....	226
Table BH6.9.	Lifetime Other Illegal Drug Use	227
Table BH6.10.	Age of First Prescription Drug Misuse.....	228
Table BH6.11.	Age of First Alcohol Use	229
Table BH6.12.	Age of First Cigarette Use	230
Table BH6.13.	Age of First Electronic Cigarette Use	231
Table BH6.14.	Age of First Marijuana Use	232
Table BH6.15.	Past Month Prescription Drug Misuse	233

Table BH6.16.	Past Month Alcohol Use	233
Table BH6.17.	Past Month Binge Drinking	234
Table BH6.18.	Past Month Cigarette Use	234
Table BH6.19.	Past Month Electronic Cigarette Use	235
Table BH6.20.	Past Month Smokeless Tobacco Use	235
Table BH6.21.	Past Month Marijuana Use	236
Table BH6.22.	Past Month Cocaine Use.....	236
Table BH6.23.	Past Month Inhalants Use	237
Table BH6.24.	Past Month Hallucinogens Use	237
Table BH6.25.	Past Month Heroin Use	238
Table BH6.26.	Past Month Steroids Use.....	238
Table BH6.27.	Past Month Ecstasy Use	239
Table BH6.28.	Past Month Methamphetamines Use	239
Table BH6.29.	Availability of Prescription Drugs.....	240
Table BH6.30.	Availability of Alcohol.....	240
Table BH6.31.	Availability of Tobacco.....	241
Table BH6.32.	Availability of Marijuana.....	241
Table BH6.33.	Availability of Prescription Drugs.....	242
Table BH6.34.	Availability of Alcohol.....	242
Table BH6.35.	Availability of Prescription Drugs – Sources	243
Table BH6.36.	Availability of Alcohol – Sources	244
Table BH6.37.	Asked to Show Proof of Age to Buy Alcohol	245
Table BH6.38.	Ability to Purchase Alcohol.....	245
Table BH6.39.	Ever Ridden with a Prescription Drugs Impaired Driver.....	246
Table BH6.40.	Frequency Riding with a Prescription Drugs Impaired Driver	246
Table BH6.41.	Ever Ridden with an Alcohol or Drug Impaired Driver.....	247
Table BH6.42.	Frequency Riding with an Alcohol or Drug Impaired Driver	247
Table BH6.43.	Parents Feel Misusing Prescription Drugs is Wrong.....	248
Table BH6.44.	Parents Feel Drinking Alcohol is Wrong.....	248
Table BH6.45.	Parents Feel Regularly Drinking Alcohol is Wrong	249
Table BH6.46.	Parents Feel Smoking Tobacco is Wrong.....	249
Table BH6.47.	Parents Feel Smoking One or More Packs of Cigarettes is Wrong	250
Table BH6.48.	Parents Feel Using Electronic Cigarettes is Wrong	250
Table BH6.49.	Parents Feel Smoking Marijuana is Wrong.....	251
Table BH6.50.	Parents Feel Smoking Marijuana Once or Twice a Week is Wrong	251
Table BH6.51.	Friends Feel Misusing Prescription Drugs is Wrong	252
Table BH6.52.	Friends Feel Drinking Alcohol is Wrong	252
Table BH6.53.	Friends Feel Smoking Tobacco is Wrong	253
Table BH6.54.	Friends Feel Using Electronic Cigarettes is Wrong.....	253
Table BH6.55.	Friends Feel Smoking Marijuana is Wrong	254
Table BH6.56.	Friends Misused Prescription Drugs	254
Table BH6.57.	Friends Tried Alcohol.....	255
Table BH6.58.	Friends Binge Drinking	255
Table BH6.59.	Friends Smoked Cigarettes.....	256
Table BH6.60.	Friends Used Electronic Cigarettes.....	256
Table BH6.61.	Friends Used Marijuana	257
Table BH6.62.	Perceived Risk of Misusing Prescription Drugs	257
Table BH6.63.	Perceived Risk of Regularly Drinking Alcohol	258
Table BH6.64.	Perceived Risk of Binge Drinking	258
Table BH6.65.	Perceived Risk of Smoking Cigarettes	259
Table BH6.66.	Perceived Risk of Using Electronic Cigarettes	259
Table BH6.67.	Perceived Risk of Trying Marijuana.....	260
Table BH6.68.	Perceived Risk of Regularly Smoking Marijuana	260
Table BH6.69.	Wrong to Misuse Prescription Drugs.....	261
Table BH6.70.	Wrong to Regularly Drink Alcohol	261
Table BH6.71.	Wrong to Smoke Tobacco.....	262
Table BH6.72.	Wrong to Use Electronic Cigarettes	262
Table BH6.73.	Wrong to Smoke Marijuana.....	263
Table BH6.74.	Parents Talked about Prescription Drug Misuse.....	263
Table BH6.75.	Parents Talked about Alcohol and Other Drugs (ATOD)	264

Introduction

The *Tennessee Together Student Survey Results Compendium 2018-2019: Detailed Tables* is a compilation of tables presenting weighted statewide statistics from the Tennessee Together Student Survey administered during the 2018-2019 school year. The 2018-2019 Tennessee Together Student Survey was administered to 8th-, 10th-, and 12th-grade students enrolled in Tennessee public schools. The final sample included 28 counties with 21,766 survey respondents. The detailed tables in this report present weighted state and Behavioral Health Planning Region summaries of survey responses profiling student tobacco, alcohol, marijuana, prescription drug, and other drug use. State-level data tables are named with the prefix “TN” and Behavioral Health Planning Region tables are named with the prefix “BH” and the region number. Survey results are also presented by grade level, as well as by race/ethnicity and gender. The findings from this report can be used by the state and local school districts to identify emergent needs related to alcohol and drug use prevention; to monitor changes in alcohol, tobacco, and drug use prevalence and attitudes over time; and to assess the impacts of community-wide efforts to prevent and reduce alcohol and drug misuse among youth within local schools and communities.

The detailed tables in this document include weighted prevalence estimates (%) and 95 percent confidence intervals. Post-stratification weighting of survey responses by grade level was used to adjust for school and student nonresponse. The overall weights were scaled so that the weighted count of students equaled the total state enrolled population, and the weighted proportions of students in each grade matched the state population proportions. Enrollment data from the Tennessee Department of Education 2017-2018 Membership File were used to estimate student populations. To account for the complex sampling designs, statistical analyses were conducted on the weighted data using STATA statistical software. Table A documents the size of the unweighted survey sample for each participating county.

This document was developed to accompany the *2018-2019 TN Together Student Survey Results* narrative report which provides a more descriptive introduction about the survey administration and methodology, as well as a comprehensive analysis of survey results.

Table A. Unweighted and Weighted Sample by County

	Unweighted Sample (n)	Weighted Sample (n)	Proportion of Weighted N
Blount County	3,028	25,174	11.8%
Carter County	338	4,029	1.9%
Clay County	219	1,271	0.6%
DeKalb County	495	3,927	1.8%
Dickson County	970	11,223	5.3%
Dyer County	160	1,768	0.8%
Gibson County	683	5,334	2.5%
Greene County	385	7,851	3.7%
Hamilton County	1,449	17,220	8.1%
Henry County	732	6,301	3.0%
Johnson County	321	2,992	1.4%
Lawrence County	1,032	9,353	4.4%
Macon County	556	5,270	2.5%
Monroe County	1,062	8,872	4.2%
Overton County	244	3,000	1.4%
Putnam County	1,502	13,241	6.2%
Roane County	990	8,236	3.9%
Scott County	756	5,938	2.8%
Sevier County	489	12,372	5.8%
Smith County	387	4,189	2.0%
Stewart County	353	3,100	1.5%
Sullivan County	714	10,100	4.7%
Tipton County	1,558	15,604	7.3%
Unicoi County	359	2,388	1.1%
Union County	175	1,225	0.6%
Weakley County	390	5,815	2.7%
White County	381	3,594	1.7%
Williamson County	2,038	1,4399	6.7%
Total (N)	21,766	213,786	100.0%

Demographic Characteristics of the Respondent Sample

Table D.1. Age of Sample

	(n=21,753)	%
10 years old	(n=21)	0.1%
11 years old	(n=5)	0.0%
12 years old	(n=39)	0.2%
13 years old	(n=5,489)	25.2%
14 years old	(n=3,733)	17.2%
15 years old	(n=4,098)	18.8%
16 years old	(n=2,890)	13.3%
17 years old	(n=3,360)	15.4%
18+ years old	(n=2,118)	9.7%

Table D.2. Grade Level of Sample

	(n=21,766)	%
8th Grade	(n=9,562)	43.9%
10th Grade	(n=6,895)	31.7%
12th Grade	(n=5,309)	24.4%

Table D.3. Gender of Sample

	(n=21,520)	%
Male	(n=10,634)	50.3%
Female	(n=10,508)	49.7%
Other	(n=378)	1.8%

Note: The demographic data presented represents the unweighted respondent sample.

Table D.4. Race and Ethnicity of Sample

	(n=21,648)	%
Am. Indian/Alaska Native	(n=95)	0.4%
Asian/Pacific Islander	(n=344)	1.6%
Black/African American	(n=1,432)	6.6%
Hispanic	(n=974)	4.5%
White	(n=16,705)	77.2%
Other	(n=278)	1.3%
Multi-Race	(n=1,820)	8.4%

Table D.5. Living Situation of Sample

	(n=21,450)	%
Both parents	(n=11,859)	55.6%
Father and stepmother	(n=941)	4.3%
Mother and stepfather	(n=2,699)	12.3%
Father only	(n=767)	3.5%
Mother only	(n=2,984)	13.9%
Other	(n=989)	4.6%
Grandparents	(n=1,031)	4.8%
Foster family	(n=180)	0.8%

Table D.6. School Grades of Sample

	(n=21,412)	%
Mostly A's	(n=8,948)	41.8%
Mostly B's	(n=7,985)	37.3%
Mostly C's	(n=3,591)	16.8%
Mostly D's	(n=612)	2.9%
Mostly F's	(n=276)	1.3%

Statewide Results Summary

Table TN.1. Lifetime Prescription Drug Misuse

		On how many occasions (if any) have you ever <u>used prescription drugs not prescribed to you?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	91.1	4.6	1.7	0.9	0.7	0.2	0.9	8.9
(n=21,312)	CI	90.6–91.5	4.2–4.9	1.5–1.9	0.7–1.0	0.5–0.8	0.2–0.3	0.8–1.1	8.5–9.4
Male	%	92.2	3.4	1.4	0.8	0.6	0.3	1.3	7.8
(n=10,381)	CI	91.6–92.8	3.0–3.8	1.2–1.7	0.6–1.0	0.5–0.9	0.2–0.4	1.0–1.6	7.2–8.4
Female	%	90.5	5.5	1.9	0.8	0.7	0.2	0.4	9.5
(n=10,370)	CI	89.9–91.2	5.0–6.0	1.6–2.2	0.7–1.1	0.5–0.9	0.1–0.3	0.3–0.6	8.8–10.1
White	%	91.4	4.5	1.6	0.8	0.6	0.3	0.8	8.6
(n=16,416)	CI	90.9–91.9	4.1–4.9	1.4–1.9	0.6–0.9	0.5–0.8	0.2–0.3	0.6–1.0	8.1–9.1
Black	%	89.9	5.0	1.7	0.6	0.5	0.4	1.9	10.1
(n=1,392)	CI	88.1–91.5	3.9–6.5	1.1–2.5	0.3–1.2	0.2–1.0	0.2–0.9	1.3–2.9	8.5–11.9
Hispanic	%	93.8	3.1	1.2	0.9	0.5	0.0	0.5	6.2
(n=952)	CI	91.8–95.3	2.1–4.6	0.7–2.3	0.4–1.8	0.2–1.4	–	0.2–1.2	4.7–8.2
8th	%	94.2	3.5	1.0	0.5	0.3	0.1	0.4	5.8
(n=9,414)	CI	93.7–94.7	3.1–3.9	0.8–1.3	0.4–0.7	0.2–0.5	0.0–0.2	0.3–0.5	5.3–6.3
10th	%	89.7	5.2	2.0	1.0	0.6	0.3	1.2	10.3
(n=6,739)	CI	88.8–90.4	4.6–5.9	1.6–2.4	0.7–1.3	0.4–0.9	0.2–0.5	1.0–1.5	9.6–11.2
12th	%	89.1	5.0	2.1	1.1	1.1	0.4	1.2	10.9
(n=5,159)	CI	88.1–90.0	4.4–5.8	1.7–2.5	0.8–1.4	0.8–1.5	0.2–0.6	0.9–1.7	10.0–11.9

Note: The confidence interval (CI) is 95 percent.

Table TN.2. Lifetime Alcohol Use

		On how many occasions (if any) have you ever <u>had one or more drinks of an alcoholic beverage?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	60.1	17.8	7.5	4.2	3.7	2.4	4.4	39.9
(n=21,431)	CI	59.4–60.9	17.2–18.4	7.1–7.9	3.9–4.5	3.4–4.0	2.1–2.6	4.0–4.7	39.1–40.6
Male	%	61.5	16.0	6.8	4.1	3.7	2.2	5.6	38.5
(n=10,456)	CI	60.4–62.5	15.2–16.8	6.3–7.4	3.7–4.6	3.3–4.2	1.9–2.6	5.1–6.2	37.5–39.6
Female	%	59.3	19.4	8.1	4.2	3.7	2.5	2.8	40.7
(n=10,405)	CI	58.2–60.4	18.5–20.3	7.5–8.8	3.8–4.7	3.3–4.2	2.1–2.9	2.4–3.2	39.6–41.8
White	%	59.8	17.5	7.5	4.3	3.8	2.5	4.6	40.2
(n=16,493)	CI	59.0–60.7	16.8–18.2	7.0–7.9	4.0–4.7	3.4–4.1	2.2–2.8	4.2–5.1	39.3–41.0
Black	%	65.4	17.9	6.8	3.1	3.6	1.1	2.1	34.6
(n=1,407)	CI	62.6–68.2	15.7–20.3	5.6–8.4	2.2–4.2	2.5–5.0	0.6–2.0	1.4–3.1	31.8–37.4
Hispanic	%	65.7	20.0	5.1	2.4	2.5	1.9	2.5	34.3
(n=956)	CI	62.2–69.0	17.3–23.1	3.7–6.9	1.6–3.6	1.5–4.0	1.1–3.2	1.5–4.0	31.0–37.8
8th	%	74.3	16.5	4.5	2.0	1.2	0.7	0.8	25.7
(n=9,453)	CI	73.4–75.3	15.7–17.3	4.1–5.0	1.7–2.3	1.0–1.5	0.5–0.9	0.6–1.0	24.7–26.6
10 th	%	56.1	19.3	8.9	5.1	4.4	2.5	3.8	43.9
(n=6,781)	CI	54.8–57.4	18.2–20.3	8.1–9.7	4.5–5.7	3.9–5.0	2.1–2.9	3.3–4.3	42.6–45.2
12 th	%	48.5	17.6	9.4	5.6	5.7	4.1	9.1	51.5
(n=5,197)	CI	47.0–50.1	16.4–18.8	8.5–10.3	5.0–6.3	5.1–6.4	3.5–4.9	8.2–10.0	49.9–53.0

Note: The confidence interval (CI) is 95 percent.

Table TN.3. Lifetime Binge Drinking

		On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	81.4	6.7	3.7	2.4	2.1	1.4	2.3	18.6
(n=21,411)	CI	80.7–82.0	6.3–7.1	3.4–4.0	2.2–2.7	1.9–2.4	1.2–1.6	2.0–2.5	18.0–19.3
Male	%	80.5	6.1	3.6	2.7	2.3	1.8	3.2	19.5
(n=10,434)	CI	79.5–81.4	5.6–6.6	3.2–4.0	2.3–3.1	2.0–2.6	1.4–2.1	2.7–3.6	18.6–20.5
Female	%	82.6	7.3	3.8	2.2	1.9	1.0	1.2	17.4
(n=10,411)	CI	81.7–83.5	6.7–7.9	3.4–4.3	1.9–2.6	1.6–2.3	0.8–1.3	0.9–1.4	16.5–18.3
White	%	80.8	6.7	3.8	2.6	2.3	1.5	2.4	19.2
(n=16,486)	CI	80.0–81.5	6.3–7.2	3.5–4.1	2.3–2.9	2.0–2.6	1.3–1.8	2.2–2.8	18.5–20.0
Black	%	86.2	5.6	3.5	1.9	0.9	0.8	1.0	13.8
(n=1,402)	CI	83.9–88.3	4.4–7.2	2.3–5.3	1.2–2.8	0.5–1.7	0.5–1.5	0.6–1.7	11.7–16.1
Hispanic	%	86.7	5.5	2.4	2.0	1.2	1.0	1.2	13.3
(n=953)	CI	83.9–89.1	4.0–7.6	1.5–3.9	1.2–3.2	0.6–2.6	0.5–2.0	0.6–2.1	10.9–16.1
8th	%	93.7	3.5	1.2	0.7	0.4	0.2	0.3	6.3
(n=9,441)	CI	93.2–94.2	3.1–3.9	1.0–1.4	0.5–0.9	0.3–0.6	0.1–0.3	0.2–0.5	5.8–6.8
10th	%	80.1	7.9	4.1	3.0	1.9	1.3	1.8	19.9
(n=6,779)	CI	79.0–81.1	7.2–8.7	3.6–4.6	2.6–3.5	1.6–2.3	1.0–1.6	1.5–2.1	18.9–21.0
12th	%	68.8	9.0	6.2	3.8	4.3	2.9	5.0	31.2
(n=5,191)	CI	67.4–70.2	8.2–10.0	5.5–6.9	3.3–4.4	3.7–5.0	2.4–3.5	4.3–5.8	29.8–32.6

Note: The confidence interval (CI) is 95 percent.

Table TN.4. Lifetime Cigarette Use

		On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	80.7	8.7	2.8	1.7	1.4	1.0	3.7	19.3
(n=21,424)	CI	80.1–81.4	8.2–9.1	2.5–3.1	1.5–2.0	1.3–1.6	0.8–1.2	3.4–4.0	18.6–19.9
Male	%	80.5	8.2	2.7	1.6	1.5	1.1	4.3	19.5
(n=10,454)	CI	79.6–81.4	7.7–8.8	2.4–3.1	1.4–1.9	1.3–1.8	0.8–1.4	3.8–4.9	18.6–20.4
Female	%	81.6	8.8	2.8	1.8	1.3	0.8	2.9	18.4
(n=10,401)	CI	80.7–82.4	8.2–9.5	2.4–3.2	1.5–2.1	1.1–1.6	0.6–1.0	2.5–3.3	17.6–19.3
White	%	79.7	8.9	3.0	1.8	1.5	1.1	4.1	20.3
(n=16,495)	CI	79.0–80.4	8.4–9.4	2.7–3.3	1.6–2.1	1.3–1.8	0.9–1.3	3.7–4.5	19.6–21.0
Black	%	88.0	5.9	1.8	1.2	1.1	0.4	1.4	12.0
(n=1,402)	CI	85.9–89.8	4.7–7.4	1.1–3.0	0.6–2.3	0.6–2.0	0.2–1.0	0.9–2.3	10.2–14.1
Hispanic	%	87.0	6.9	1.9	1.5	0.6	0.6	1.5	13.0
(n=958)	CI	84.5–89.2	5.3–8.9	1.0–3.3	0.9–2.5	0.3–1.2	0.2–1.7	0.9–2.7	10.8–15.5
8th	%	88.5	6.8	1.8	0.8	0.7	0.5	0.9	11.5
(n=9,457)	CI	87.8–89.1	6.3–7.4	1.5–2.1	0.6–1.0	0.5–0.9	0.4–0.7	0.7–1.1	10.9–12.2
10th	%	79.3	9.5	2.9	1.8	1.6	1.0	3.9	20.7
(n=6,783)	CI	78.2–80.4	8.8–10.4	2.5–3.4	1.5–2.2	1.3–2.0	0.7–1.2	3.4–4.4	19.6–21.8
12th	%	73.6	9.7	3.8	2.7	2.1	1.5	6.6	26.4
(n=5,184)	CI	72.2–74.9	8.9–10.6	3.3–4.4	2.2–3.2	1.7–2.6	1.2–2.0	5.8–7.5	25.1–27.8

Note: The confidence interval (CI) is 95 percent.

Table TN.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	70.2	9.5	4.2	2.4	2.4	2.0	9.3	29.8
(n=21,368)	CI	69.5–70.9	9.1–10.0	3.9–4.5	2.2–2.7	2.2–2.7	1.8–2.2	8.8–9.8	29.1–30.5
Male	%	69.0	9.3	4.2	2.0	2.2	2.0	11.3	31.0
(n=10,425)	CI	67.9–70.0	8.7–9.9	3.8–4.7	1.8–2.4	1.9–2.6	1.7–2.4	10.5–12.0	30.0–32.1
Female	%	71.9	9.7	4.2	2.7	2.5	1.9	7.0	28.1
(n=10,377)	CI	70.9–72.9	9.1–10.4	3.8–4.7	2.3–3.1	2.2–2.9	1.6–2.2	6.5–7.7	27.1–29.1
White	%	68.5	9.8	4.4	2.4	2.6	2.1	10.2	31.5
(n=16,456)	CI	67.7–69.3	9.3–10.3	4.1–4.8	2.2–2.7	2.3–2.9	1.9–2.4	9.6–10.7	30.7–32.3
Black	%	82.4	6.6	2.3	1.4	1.4	1.1	4.8	17.6
(n=1,396)	CI	80.0–84.6	5.2–8.4	1.5–3.3	0.9–2.2	0.9–2.2	0.5–2.4	3.7–6.2	15.4–20.0
Hispanic	%	81.5	6.6	3.1	2.2	1.4	1.8	3.4	18.5
(n=954)	CI	78.6–84.1	5.1–8.7	2.0–4.8	1.3–3.8	0.8–2.3	1.1–2.9	2.4–4.7	15.9–21.4
8th	%	81.6	8.1	3.0	1.8	1.5	1.2	2.9	18.4
(n=9,437)	CI	80.8–82.4	7.6–8.7	2.6–3.3	1.5–2.1	1.2–1.7	1.0–1.5	2.6–3.3	17.6–19.2
10th	%	66.0	10.7	4.9	2.8	2.9	2.1	10.6	34.0
(n=6,749)	CI	64.7–67.3	9.9–11.5	4.4–5.6	2.4–3.3	2.5–3.4	1.7–2.5	9.8–11.4	32.7–35.3
12th	%	62.1	9.8	4.8	2.7	2.9	2.7	15.1	37.9
(n=5,182)	CI	60.5–63.6	8.9–10.8	4.1–5.5	2.2–3.2	2.4–3.5	2.3–3.3	14.0–16.3	36.4–39.5

Note: The confidence interval (CI) is 95 percent.

Table TN.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	88.2	4.3	1.5	1.0	0.9	0.6	3.5	11.8
(n=21,362)	CI	87.7–88.8	4.0–4.6	1.3–1.7	0.9–1.2	0.7–1.0	0.5–0.7	3.2–3.9	11.2–12.3
Male	%	82.8	5.4	2.1	1.4	1.4	0.9	6.0	17.2
(n=10,437)	CI	81.9–83.6	4.9–6.0	1.8–2.5	1.2–1.7	1.2–1.7	0.7–1.1	5.4–6.6	16.4–18.1
Female	%	94.1	3.1	0.7	0.5	0.3	0.3	0.9	5.9
(n=10,358)	CI	93.5–94.6	2.7–3.6	0.6–0.9	0.4–0.8	0.2–0.4	0.2–0.5	0.7–1.2	5.4–6.5
White	%	87.1	4.6	1.6	1.1	0.9	0.7	4.1	12.9
(n=16,442)	CI	86.5–87.7	4.2–5.0	1.4–1.8	0.9–1.3	0.7–1.0	0.5–0.8	3.8–4.5	12.3–13.5
Black	%	93.9	2.5	0.9	0.4	0.8	0.3	1.2	6.1
(n=1,400)	CI	92.2–95.3	1.7–3.6	0.4–2.0	0.2–0.9	0.3–1.9	0.1–1.4	0.7–2.0	4.7–7.8
Hispanic	%	94.4	2.8	0.6	0.9	0.3	0.2	0.7	5.6
(n=953)	CI	92.6–95.7	1.9–4.1	0.3–1.4	0.4–1.9	0.1–0.9	0.1–1.0	0.3–1.6	4.3–7.4
8th	%	93.7	3.0	0.9	0.6	0.4	0.4	1.0	6.3
(n=9,437)	CI	93.2–94.2	2.7–3.4	0.7–1.1	0.5–0.8	0.3–0.5	0.3–0.6	0.8–1.2	5.8–6.8
10th	%	87.2	4.5	1.7	1.0	1.1	0.7	3.7	12.8
(n=6,748)	CI	86.3–88.1	4.0–5.1	1.4–2.1	0.8–1.4	0.9–1.5	0.5–0.9	3.2–4.2	11.9–13.7
12th	%	83.2	5.4	1.9	1.4	1.1	0.8	6.3	16.8
(n=5,177)	CI	81.9–84.4	4.7–6.2	1.5–2.3	1.1–1.8	0.8–1.4	0.5–1.1	5.5–7.1	15.6–18.1

Note: The confidence interval (CI) is 95 percent.

Table TN.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.8	6.1	2.9	1.8	2.0	1.5	5.8	20.2
(n=21,380)	CI	79.2–80.5	5.8–6.5	2.7–3.2	1.6–2.0	1.8–2.3	1.3–1.7	5.4–6.1	19.5–20.8
Male	%	80.0	5.5	2.6	1.7	2.2	1.4	6.7	20.0
(n=10,435)	CI	79.1–80.9	5.0–6.0	2.3–2.9	1.4–2.0	1.8–2.6	1.1–1.7	6.1–7.3	19.1–20.9
Female	%	80.2	6.8	3.2	1.9	1.9	1.6	4.6	19.8
(n=10,377)	CI	79.3–81.1	6.2–7.4	2.8–3.6	1.6–2.2	1.6–2.2	1.3–1.9	4.1–5.0	18.9–20.7
White	%	80.8	5.8	2.7	1.7	1.9	1.5	5.5	19.2
(n=16,468)	CI	80.1–81.5	5.4–6.3	2.5–3.0	1.5–2.0	1.7–2.2	1.2–1.7	5.1–5.9	18.5–19.9
Black	%	72.7	8.2	3.5	2.8	2.9	1.9	8.0	27.3
(n=1,397)	CI	69.9–75.3	6.5–10.2	2.6–4.7	2.0–4.0	2.0–4.0	1.2–3.1	6.5–9.8	24.7–30.1
Hispanic	%	84.8	5.8	2.5	1.1	1.8	1.1	2.7	15.2
(n=953)	CI	82.0–87.3	4.3–7.9	1.7–3.8	0.6–2.2	1.0–3.2	0.6–2.3	1.8–4.1	12.7–18.0
8th	%	91.5	3.7	1.4	0.7	0.8	0.7	1.2	8.5
(n=9,430)	CI	90.9–92.1	3.3–4.1	1.2–1.7	0.5–0.9	0.6–1.0	0.5–0.9	1.0–1.4	7.9–9.1
10th	%	77.1	7.1	3.3	2.2	2.2	1.9	6.1	22.9
(n=6,772)	CI	76.0–78.2	6.4–7.8	2.9–3.9	1.8–2.6	1.9–2.7	1.5–2.3	5.5–6.8	21.8–24.0
12th	%	69.6	7.8	4.2	2.5	3.3	2.0	10.6	30.4
(n=5,178)	CI	68.2–71.0	7.0–8.7	3.7–4.8	2.1–3.0	2.8–3.9	1.6–2.5	9.6–11.5	29.0–31.8

Note: The confidence interval (CI) is 95 percent.

Table TN.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	98.7	0.6	0.2	0.2	0.1	0.0	0.3	1.3
(n=21,395)	CI	98.5–98.8	0.5–0.7	0.1–0.2	0.1–0.3	0.1–0.2	0.0–0.1	0.2–0.3	1.2–1.5
Male	%	98.6	0.6	0.2	0.2	0.1	0.0	0.2	1.4
(n=10,441)	CI	98.3–98.9	0.5–0.9	0.1–0.4	0.1–0.3	0.0–0.2	0.0–0.1	0.1–0.3	1.1–1.7
Female	%	99.1	0.5	0.1	0.1	0.1	0.0	0.1	0.9
(n=10,386)	CI	98.8–99.3	0.3–0.7	0.0–0.2	0.1–0.2	0.1–0.3	0.0–0.1	0.1–0.2	0.7–1.2
White	%	98.9	0.6	0.1	0.1	0.1	0.0	0.1	1.1
(n=16,477)	CI	98.7–99.1	0.4–0.7	0.1–0.2	0.1–0.2	0.1–0.2	0.0–0.1	0.1–0.2	0.9–1.3
Black	%	97.7	1.0	0.4	0.3	0.0	0.1	0.6	2.3
(n=1,395)	CI	96.6–98.4	0.5–1.8	0.1–1.1	0.1–0.8	–	0.0–0.9	0.3–1.2	1.6–3.4
Hispanic	%	99.3	0.1	0.2	0.0	0.1	0.0	0.4	0.7
(n=955)	CI	98.4–99.7	0.0–0.5	0.0–0.8	–	0.0–0.5	–	0.1–1.2	0.3–1.6
8th	%	99.4	0.3	0.1	0.1	0.1	0.0	0.1	0.6
(n=9,435)	CI	99.2–99.5	0.2–0.4	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.1–0.2	0.5–0.8
10th	%	98.4	0.7	0.2	0.2	0.1	0.1	0.3	1.6
(n=6,782)	CI	98.0–98.7	0.5–1.0	0.1–0.4	0.1–0.3	0.1–0.4	0.0–0.2	0.2–0.4	1.3–2.0
12th	%	98.2	0.8	0.2	0.3	0.2	0.0	0.4	1.8
(n=5,178)	CI	97.7–98.5	0.5–1.2	0.1–0.3	0.2–0.6	0.1–0.3	0.0–0.1	0.3–0.6	1.5–2.3

Note: The confidence interval (CI) is 95 percent.

Table TN.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	97.0	1.3	0.5	0.3	0.3	0.1	0.4	3.0
(n=21,411)	CI	96.7–97.3	1.2–1.5	0.4–0.6	0.2–0.4	0.2–0.4	0.1–0.2	0.3–0.5	2.7–3.3
Male	%	96.7	1.4	0.6	0.4	0.4	0.1	0.4	3.3
(n=10,441)	CI	96.2–97.1	1.2–1.7	0.4–0.8	0.3–0.6	0.3–0.7	0.0–0.4	0.3–0.6	2.9–3.8
Female	%	97.8	1.1	0.4	0.2	0.1	0.1	0.3	2.2
(n=10,402)	CI	97.4–98.1	0.9–1.4	0.3–0.5	0.1–0.3	0.1–0.3	0.1–0.2	0.2–0.5	1.9–2.6
White	%	97.2	1.3	0.5	0.3	0.3	0.1	0.3	2.8
(n=16,481)	CI	96.9–97.5	1.1–1.5	0.4–0.6	0.2–0.4	0.2–0.4	0.1–0.3	0.2–0.5	2.5–3.1
Black	%	97.5	0.8	0.4	0.3	0.1	0.0	0.9	2.5
(n=1,397)	CI	96.3–98.2	0.4–1.6	0.2–1.1	0.1–0.8	0.0–0.6	–	0.5–1.6	1.8–3.7
Hispanic	%	97.3	1.5	0.3	0.6	0.1	0.0	0.2	2.7
(n=959)	CI	95.8–98.3	0.8–2.9	0.1–0.9	0.2–1.6	0.0–0.5	–	0.0–0.9	1.7–4.2
8th	%	98.7	0.7	0.1	0.1	0.1	0.0	0.2	1.3
(n=9,443)	CI	98.5–98.9	0.6–0.9	0.1–0.2	0.1–0.2	0.0–0.2	0.0–0.1	0.1–0.3	1.1–1.5
10th	%	96.8	1.5	0.6	0.3	0.2	0.1	0.5	3.2
(n=6,784)	CI	96.3–97.2	1.2–1.8	0.4–0.8	0.2–0.4	0.1–0.4	0.1–0.2	0.4–0.8	2.8–3.7
12th	%	95.4	1.8	0.8	0.6	0.6	0.2	0.6	4.6
(n=5,184)	CI	94.6–96.0	1.4–2.2	0.6–1.1	0.4–0.9	0.4–1.0	0.1–0.6	0.4–0.9	4.0–5.4

Note: The confidence interval (CI) is 95 percent.

Table TN.10. Age of First Prescription Drug Misuse

		How old were you when you first <u>used prescription drugs not prescribed to you?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	92.3	1.2	0.5	0.8	1.2	1.1	1.4	0.9	0.7	13.47
(n=21,473)	CI	91.9–92.7	1.1–1.4	0.4–0.6	0.7–0.9	1.0–1.3	1.0–1.3	1.2–1.6	0.7–1.0	0.5–0.8	13.3–13.6
Male	%	93.7	1.1	0.4	0.6	0.8	0.7	1.2	0.8	0.7	13.54
(n=10,479)	CI	93.1–94.2	0.9–1.4	0.3–0.6	0.4–0.7	0.6–1.1	0.6–1.0	1.0–1.5	0.6–1.0	0.5–0.9	13.3–13.7
Female	%	91.4	1.1	0.6	1.0	1.4	1.4	1.5	0.9	0.7	13.50
(n=10,423)	CI	90.8–92.0	0.9–1.4	0.5–0.7	0.8–1.2	1.2–1.7	1.2–1.7	1.3–1.8	0.7–1.2	0.5–0.9	13.3–13.7
White	%	92.6	1.1	0.5	0.7	1.1	1.1	1.4	0.9	0.6	13.53
(n=16,535)	CI	92.1–93.0	1.0–1.3	0.4–0.6	0.5–0.8	1.0–1.4	0.9–1.3	1.2–1.6	0.7–1.1	0.5–0.8	13.4–13.7
Black	%	91.9	1.4	0.5	1.4	1.0	1.0	1.1	0.8	1.0	13.40
(n=1,402)	CI	90.2–93.3	0.9–2.2	0.2–1.0	0.9–2.2	0.6–1.7	0.5–1.8	0.6–2.0	0.4–1.5	0.6–1.8	12.9–13.9
Hispanic	%	94.7	0.7	0.2	0.4	0.6	1.3	1.1	0.1	0.9	13.89
(n=958)	CI	92.9–96.1	0.3–1.5	0.0–0.6	0.1–1.3	0.2–1.5	0.7–2.4	0.6–2.3	0.0–0.6	0.4–2.0	13.2–14.6
8th	%	95.2	1.3	0.7	0.9	1.4	0.5	0.0	0.0	0.0	11.81
(n=9,474)	CI	94.7–95.7	1.1–1.6	0.5–0.9	0.8–1.2	1.1–1.6	0.3–0.7	0.0–0.1	–	0.0–0.1	11.7–11.9
10th	%	91.0	1.4	0.5	1.1	1.2	1.8	2.4	0.5	0.1	13.20
(n=6,803)	CI	90.2–91.7	1.2–1.8	0.3–0.8	0.8–1.4	0.9–1.5	1.5–2.2	2.0–2.8	0.3–0.7	0.0–0.2	13.0–13.4
12th	%	90.4	0.8	0.3	0.3	0.9	1.1	1.8	2.3	2.1	14.69
(n=5,196)	CI	89.4–91.2	0.6–1.2	0.2–0.5	0.2–0.5	0.6–1.3	0.9–1.5	1.4–2.2	1.9–2.8	1.7–2.5	14.5–14.9

Note: The confidence interval (CI) is 95 percent.

Table TN.11. Age of First Alcohol Use

How old were you when you first had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	63.7	4.6	2.1	3.3	5.3	5.8	6.6	5.1	3.4	13.74
(n=21,420)	CI	62.9–64.4	4.3–4.9	1.9–2.3	3.1–3.6	5.0–5.7	5.4–6.2	6.2–7.0	4.8–5.5	3.1–3.8	13.7–13.8
Male	%	65.1	5.2	2.3	3.4	4.7	5.2	6.2	4.3	3.5	13.61
(n=10,454)	CI	64.0–66.1	4.7–5.7	2.0–2.6	3.1–3.9	4.3–5.2	4.7–5.7	5.6–6.7	3.9–4.9	3.1–4.0	13.5–13.7
Female	%	62.7	3.7	1.9	3.2	5.9	6.2	7.1	6.0	3.4	13.92
(n=10,398)	CI	61.6–63.8	3.3–4.1	1.6–2.2	2.8–3.5	5.4–6.4	5.6–6.8	6.6–7.7	5.4–6.6	3.0–3.9	13.8–14.0
White	%	63.2	4.2	2.0	3.4	5.3	6.1	6.9	5.3	3.6	13.81
(n=16,500)	CI	62.3–64.0	3.9–4.6	1.8–2.3	3.1–3.8	4.9–5.7	5.7–6.6	6.4–7.4	4.9–5.8	3.2–3.9	13.7–13.9
Black	%	70.6	4.3	2.4	2.4	3.9	3.2	5.2	4.3	3.6	13.71
(n=1,401)	CI	67.8–73.2	3.2–5.7	1.7–3.4	1.8–3.4	2.9–5.1	2.3–4.5	4.1–6.7	3.2–5.6	2.5–5.3	13.4–14.0
Hispanic	%	68.5	3.9	2.6	3.2	4.9	4.3	5.6	3.6	3.4	13.64
(n=955)	CI	65.1–71.7	2.8–5.3	1.6–4.0	2.2–4.8	3.7–6.5	3.1–6.1	4.1–7.5	2.4–5.3	2.2–5.3	13.4–13.9
8th	%	78.1	5.0	3.0	4.3	7.0	2.5	0.1	0.0	0.0	11.96
(n=9,453)	CI	77.2–79.0	4.6–5.5	2.6–3.4	3.9–4.8	6.5–7.5	2.2–2.8	0.0–0.2	–	–	11.9–12.0
10th	%	60.0	5.0	1.7	3.4	5.8	9.5	10.9	3.4	0.3	13.53
(n=6,783)	CI	58.7–61.3	4.4–5.6	1.5–2.1	2.9–3.9	5.2–6.5	8.7–10.3	10.2–11.8	3.0–3.9	0.2–0.5	13.5–13.6
12th	%	51.4	3.6	1.6	2.2	2.9	5.3	9.1	12.9	10.9	14.85
(n=5,184)	CI	49.8–52.9	3.1–4.2	1.3–2.0	1.8–2.6	2.5–3.5	4.6–6.1	8.3–10.1	11.8–14.0	10.0–12.0	14.8–14.9

Note: The confidence interval (CI) is 95 percent.

Table TN.12. Age of First Cigarette Use

		How old were you when you first <u>smoked a cigarette, even just a puff?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	81.3	3.6	1.6	2.2	2.5	2.5	2.6	2.0	1.7	13.23
(n=21,439)	CI	80.7–81.9	3.3–3.9	1.4–1.8	2.0–2.5	2.3–2.8	2.2–2.8	2.3–2.8	1.8–2.3	1.5–1.9	13.1–13.3
Male	%	81.2	4.0	1.7	2.3	2.5	2.1	2.4	2.0	1.9	13.14
(n=10,463)	CI	80.3–82.0	3.6–4.5	1.5–2.0	1.9–2.6	2.2–2.8	1.8–2.4	2.1–2.8	1.7–2.4	1.6–2.2	13.0–13.3
Female	%	82.0	2.9	1.4	2.1	2.5	2.8	2.7	2.0	1.5	13.37
(n=10,407)	CI	81.1–82.9	2.5–3.3	1.2–1.7	1.8–2.4	2.2–2.9	2.4–3.2	2.4–3.1	1.7–2.4	1.2–1.8	13.2–13.5
White	%	80.3	3.7	1.6	2.3	2.6	2.6	2.8	2.3	1.8	13.29
(n=16,506)	CI	79.5–81.0	3.4–4.0	1.4–1.8	2.1–2.6	2.4–2.9	2.4–3.0	2.5–3.1	2.0–2.6	1.5–2.0	13.2–13.4
Black	%	89.0	2.6	0.9	1.5	1.2	0.8	1.6	1.2	1.3	13.16
(n=1,400)	CI	86.9–90.7	1.7–3.8	0.5–1.6	0.9–2.3	0.7–1.9	0.4–1.5	1.0–2.6	0.6–2.3	0.7–2.4	12.7–13.7
Hispanic	%	87.3	2.0	1.0	1.8	2.0	1.8	1.4	0.8	1.9	13.35
(n=959)	CI	84.8–89.5	1.1–3.6	0.6–1.8	1.1–3.0	1.3–3.1	1.1–3.1	0.7–2.5	0.4–1.6	1.1–3.1	12.9–13.8
8th	%	89.1	3.3	1.9	2.3	2.5	0.9	0.0	0.0	0.0	11.64
(n=9,466)	CI	88.4–89.7	2.9–3.7	1.6–2.2	2.0–2.6	2.2–2.9	0.7–1.1	0.0–0.1	0.0–0.1	0.0–0.1	11.6–11.7
10th	%	79.8	4.5	1.6	2.3	3.1	3.7	3.9	0.8	0.1	12.77
(n=6,785)	CI	78.7–80.9	4.0–5.1	1.3–2.0	2.0–2.8	2.6–3.6	3.2–4.2	3.4–4.4	0.6–1.1	0.1–0.3	12.7–12.9
12th	%	74.2	2.9	1.2	2.0	1.9	2.9	3.9	5.7	5.3	14.39
(n=5,188)	CI	72.8–75.5	2.4–3.5	0.9–1.5	1.6–2.5	1.5–2.3	2.4–3.6	3.4–4.6	5.0–6.5	4.7–6.0	14.2–14.5

Note: The confidence interval (CI) is 95 percent.

Table TN.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	71.4	1.2	1.2	2.2	4.3	4.8	6.6	4.4	4.1	14.37
(n=21,339)	CI	70.7–72.1	1.0–1.3	1.0–1.3	2.0–2.4	4.0–4.6	4.4–5.1	6.2–7.0	4.1–4.8	3.7–4.4	14.3–14.4
Male	%	70.3	1.3	1.1	2.6	4.4	5.0	6.7	4.5	4.1	14.33
(n=10,402)	CI	69.3–71.3	1.1–1.5	1.0–1.4	2.3–2.9	4.0–4.8	4.5–5.5	6.2–7.3	4.0–5.0	3.6–4.6	14.3–14.4
Female	%	72.8	0.9	1.1	1.8	4.1	4.5	6.3	4.4	4.1	14.47
(n=10,373)	CI	71.8–73.8	0.7–1.2	0.9–1.4	1.5–2.1	3.7–4.5	4.0–5.0	5.8–6.9	3.9–4.9	3.7–4.6	14.4–14.5
White	%	69.6	1.2	1.2	2.3	4.3	5.2	7.0	4.9	4.4	14.42
(n=16,428)	CI	68.7–70.4	1.0–1.3	1.0–1.4	2.0–2.5	4.0–4.7	4.8–5.6	6.6–7.5	4.5–5.3	4.0–4.8	14.4–14.5
Black	%	83.9	0.9	0.6	1.5	2.6	1.5	3.7	3.2	2.2	14.36
(n=1,402)	CI	81.5–86.0	0.5–1.6	0.3–1.1	1.0–2.2	1.8–3.7	0.9–2.3	2.7–4.9	2.2–4.6	1.3–3.8	14.0–14.7
Hispanic	%	82.8	0.5	0.4	1.8	3.1	2.8	3.5	1.8	3.3	14.41
(n=953)	CI	79.9–85.3	0.2–1.2	0.2–1.0	1.2–2.8	2.2–4.3	1.9–4.2	2.3–5.3	1.1–2.9	2.1–5.2	14.1–14.7
8th	%	82.7	1.6	1.8	3.6	7.3	2.9	0.1	0.0	0.0	12.50
(n=9,430)	CI	81.9–83.5	1.4–1.9	1.5–2.1	3.2–4.0	6.8–7.8	2.6–3.3	0.1–0.2	–	0.0–0.1	12.4–12.6
10th	%	67.2	1.2	1.2	1.8	3.4	7.8	13.6	3.6	0.2	14.19
(n=6,741)	CI	65.9–68.4	0.9–1.5	1.0–1.5	1.5–2.2	2.9–3.9	7.1–8.5	12.7–14.6	3.2–4.2	0.1–0.4	14.1–14.3
12th	%	63.3	0.7	0.4	1.1	1.9	3.5	5.9	10.3	13.0	15.57
(n=5,168)	CI	61.7–64.8	0.5–1.0	0.2–0.6	0.8–1.4	1.6–2.4	2.9–4.2	5.2–6.7	9.4–11.2	12.0–14.1	15.5–15.6

Note: The confidence interval (CI) is 95 percent.

Table TN.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	80.0	1.2	1.0	1.9	2.9	3.4	4.4	2.9	2.2	14.12
(n=21,478)	CI	79.4–80.7	1.1–1.4	0.8–1.2	1.7–2.1	2.6–3.1	3.1–3.7	4.1–4.7	2.7–3.2	2.0–2.5	14.1–14.2
Male	%	80.4	1.5	1.3	2.0	2.7	2.9	4.3	2.7	2.1	13.96
(n=10,482)	CI	79.5–81.3	1.3–1.8	1.0–1.6	1.7–2.3	2.4–3.1	2.6–3.3	3.9–4.8	2.3–3.2	1.7–2.4	13.9–14.1
Female	%	80.2	0.8	0.6	1.6	2.9	3.8	4.4	3.2	2.4	14.35
(n=10,425)	CI	79.3–81.1	0.6–1.0	0.5–0.8	1.4–1.9	2.6–3.2	3.4–4.3	3.9–4.9	2.8–3.6	2.1–2.9	14.3–14.4
White	%	81.0	1.0	0.9	1.8	2.7	3.2	4.4	2.9	2.2	14.20
(n=16,538)	CI	80.2–81.7	0.8–1.1	0.7–1.1	1.6–2.0	2.5–3.0	2.9–3.5	4.0–4.8	2.6–3.2	1.9–2.5	14.1–14.3
Black	%	73.2	2.7	1.3	2.3	3.3	4.1	5.8	3.8	3.4	14.04
(n=1,403)	CI	70.5–75.8	1.8–3.9	0.7–2.4	1.6–3.2	2.5–4.4	3.0–5.6	4.5–7.4	2.8–5.1	2.3–5.1	13.8–14.3
Hispanic	%	85.2	0.6	0.7	1.3	2.5	3.2	3.0	2.1	1.4	14.11
(n=959)	CI	82.4–87.6	0.3–1.3	0.3–1.5	0.7–2.7	1.6–3.7	2.1–4.9	2.0–4.5	1.3–3.3	0.7–2.8	13.8–14.4
8th	%	91.7	1.0	0.9	2.0	3.1	1.2	0.0	0.0	0.0	12.33
(n=9,480)	CI	91.1–92.3	0.8–1.3	0.7–1.1	1.7–2.3	2.8–3.5	1.0–1.5	0.0–0.1	–	0.0–0.1	12.2–12.4
10th	%	77.4	1.5	1.2	2.0	3.1	5.4	7.5	1.8	0.1	13.75
(n=6,800)	CI	76.3–78.5	1.3–1.9	0.9–1.6	1.7–2.4	2.7–3.6	4.8–6.0	6.8–8.2	1.5–2.2	0.0–0.2	13.7–13.8
12th	%	69.8	1.2	0.8	1.5	2.3	3.8	5.9	7.5	7.2	15.00
(n=5,198)	CI	68.3–71.2	0.9–1.5	0.6–1.2	1.2–1.9	1.9–2.8	3.2–4.4	5.2–6.6	6.7–8.4	6.4–8.1	14.9–15.1

Note: The confidence interval (CI) is 95 percent.

Table TN.15. Past Month Prescription Drug Misuse

		During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	96.3	2.0	0.8	0.3	0.2	0.1	0.2	3.7
(n=21,519)	CI	96.0–96.6	1.8–2.2	0.7–1.0	0.2–0.3	0.2–0.3	0.1–0.2	0.2–0.3	3.4–4.0
Male	%	97.1	1.5	0.6	0.2	0.2	0.2	0.1	2.9
(n=10,506)	CI	96.7–97.5	1.2–1.8	0.4–0.8	0.2–0.4	0.2–0.3	0.1–0.4	0.1–0.2	2.5–3.3
Female	%	96.0	2.4	0.8	0.2	0.2	0.1	0.2	4.0
(n=10,430)	CI	95.6–96.4	2.1–2.8	0.7–1.1	0.2–0.4	0.2–0.3	0.0–0.2	0.1–0.3	3.6–4.4
White	%	96.7	1.9	0.6	0.2	0.2	0.1	0.2	3.3
(n=16,561)	CI	96.4–97.0	1.7–2.2	0.5–0.8	0.2–0.3	0.1–0.3	0.1–0.3	0.1–0.3	3.0–3.6
Black	%	95.0	2.3	1.3	0.4	0.4	0.2	0.4	5.0
(n=1,408)	CI	93.6–96.1	1.6–3.5	0.8–2.1	0.2–0.9	0.2–0.9	0.1–0.6	0.2–1.0	3.9–6.4
Hispanic	%	97.2	0.9	1.1	0.4	0.2	0.0	0.1	2.8
(n=961)	CI	95.8–98.2	0.5–1.8	0.5–2.3	0.2–1.3	0.0–0.7	–	0.0–1.0	1.8–4.2
8th	%	97.4	1.3	0.6	0.2	0.2	0.1	0.2	2.6
(n=9,486)	CI	97.1–97.7	1.1–1.6	0.4–0.8	0.1–0.3	0.1–0.3	0.1–0.2	0.1–0.3	2.3–2.9
10th	%	95.6	2.4	1.0	0.3	0.3	0.1	0.3	4.4
(n=6,817)	CI	95.0–96.1	2.0–2.9	0.8–1.3	0.2–0.5	0.2–0.5	0.1–0.2	0.2–0.4	3.9–5.0
12th	%	95.9	2.3	0.8	0.3	0.2	0.2	0.3	4.1
(n=5,216)	CI	95.3–96.5	1.8–2.8	0.6–1.2	0.2–0.5	0.1–0.4	0.1–0.6	0.1–0.5	3.5–4.7

Note: The confidence interval (CI) is 95 percent.

Table TN.16. Past Month Alcohol Use

		During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	83.2	9.2	3.9	1.3	1.4	0.5	0.5	16.8
(n=21,449)	CI	82.6–83.8	8.8–9.7	3.6–4.2	1.1–1.5	1.2–1.6	0.4–0.6	0.4–0.6	16.2–17.4
Male	%	84.1	8.0	3.8	1.6	1.6	0.5	0.4	15.9
(n=10,470)	CI	83.2–84.9	7.4–8.6	3.4–4.3	1.3–1.9	1.3–2.0	0.3–0.7	0.3–0.6	15.1–16.8
Female	%	82.8	10.4	4.0	1.0	1.1	0.4	0.4	17.2
(n=10,398)	CI	81.9–83.7	9.7–11.1	3.5–4.5	0.8–1.2	0.9–1.4	0.2–0.6	0.3–0.6	16.3–18.1
White	%	83.3	9.0	4.0	1.3	1.4	0.5	0.4	16.7
(n=16,509)	CI	82.6–84.0	8.5–9.6	3.7–4.4	1.1–1.6	1.2–1.6	0.4–0.7	0.3–0.6	16.0–17.4
Black	%	84.4	9.7	2.8	1.7	0.6	0.4	0.5	15.6
(n=1,402)	CI	81.9–86.5	8.0–11.7	2.0–3.9	1.0–3.0	0.3–1.1	0.2–1.1	0.2–1.0	13.5–18.1
Hispanic	%	85.6	9.1	2.9	0.7	1.0	0.4	0.4	14.4
(n=956)	CI	82.8–88.0	7.2–11.5	1.9–4.4	0.3–1.6	0.4–2.3	0.1–1.0	0.2–1.2	12.0–17.2
8th	%	91.5	5.6	1.4	0.4	0.6	0.3	0.3	8.5
(n=9,461)	CI	90.8–92.0	5.1–6.1	1.2–1.7	0.3–0.6	0.4–0.8	0.2–0.4	0.2–0.4	8.0–9.2
10th	%	81.9	10.3	4.1	1.4	1.3	0.5	0.5	18.1
(n=6,790)	CI	80.8–82.9	9.5–11.2	3.6–4.6	1.1–1.8	1.0–1.6	0.3–0.8	0.3–0.7	17.1–19.2
12th	%	75.4	12.1	6.4	2.2	2.5	0.7	0.7	24.6
(n=5,198)	CI	74.0–76.7	11.1–13.1	5.7–7.3	1.7–2.7	2.0–3.1	0.5–1.1	0.5–1.0	23.3–26.0

Note: The confidence interval (CI) is 95 percent.

Table TN.17. Past Month Binge Drinking

		During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	91.4	4.1	2.2	0.8	0.8	0.2	0.3	8.6
(n=21,477)	CI	90.9–91.9	3.8–4.5	2.0–2.5	0.7–1.0	0.7–1.0	0.2–0.4	0.2–0.4	8.1–9.1
Male	%	90.5	4.4	2.4	1.1	1.0	0.3	0.3	9.5
(n=10,483)	CI	89.8–91.2	4.0–4.9	2.0–2.8	0.9–1.4	0.8–1.3	0.2–0.5	0.2–0.4	8.8–10.2
Female	%	92.6	3.9	2.0	0.5	0.6	0.1	0.2	7.4
(n=10,411)	CI	92.0–93.3	3.5–4.3	1.7–2.5	0.4–0.7	0.4–0.9	0.1–0.3	0.1–0.3	6.7–8.0
White	%	91.3	4.3	2.2	0.9	0.9	0.2	0.2	8.7
(n=16,536)	CI	90.8–91.8	3.9–4.6	1.9–2.5	0.7–1.1	0.7–1.1	0.1–0.4	0.2–0.3	8.2–9.2
Black	%	92.0	3.2	2.5	0.7	0.9	0.2	0.5	8.0
(n=1,404)	CI	89.9–93.7	2.3–4.6	1.5–4.3	0.3–1.3	0.5–1.6	0.0–0.8	0.2–1.1	6.3–10.1
Hispanic	%	93.9	2.5	2.0	0.9	0.2	0.1	0.3	6.1
(n=959)	CI	91.8–95.5	1.7–3.8	1.1–3.6	0.4–2.2	0.0–0.8	0.0–0.6	0.1–1.1	4.5–8.2
8th	%	97.4	1.3	0.6	0.2	0.2	0.1	0.1	2.6
(n=9,475)	CI	97.0–97.7	1.1–1.6	0.5–0.8	0.1–0.3	0.1–0.3	0.1–0.2	0.1–0.2	2.3–3.0
10th	%	91.1	4.5	2.1	0.9	0.9	0.2	0.3	8.9
(n=6,800)	CI	90.3–91.9	3.9–5.0	1.8–2.5	0.7–1.2	0.7–1.3	0.1–0.4	0.2–0.5	8.1–9.7
12th	%	84.9	6.9	4.2	1.6	1.5	0.4	0.5	15.1
(n=5,202)	CI	83.7–86.0	6.2–7.7	3.5–5.0	1.2–2.0	1.1–2.0	0.2–0.8	0.4–0.8	14.0–16.3

Note: The confidence interval (CI) is 95 percent.

Table TN.18. Past Month Cigarette Use

		During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	91.5	3.4	1.6	0.6	1.0	0.5	1.4	8.5
(n=21,476)	CI	91.0–91.9	3.2–3.7	1.4–1.8	0.5–0.7	0.8–1.2	0.4–0.6	1.2–1.7	8.1–9.0
Male	%	91.4	3.2	1.8	0.6	1.0	0.6	1.5	8.6
(n=10,482)	CI	90.7–92.0	2.8–3.6	1.5–2.1	0.4–0.7	0.8–1.3	0.5–0.8	1.2–1.8	8.0–9.3
Female	%	92.0	3.5	1.3	0.6	0.9	0.4	1.3	8.0
(n=10,411)	CI	91.3–92.6	3.1–3.9	1.1–1.6	0.5–0.8	0.7–1.2	0.3–0.5	1.0–1.6	7.4–8.7
White	%	91.0	3.6	1.6	0.6	1.1	0.5	1.5	9.0
(n=16,524)	CI	90.5–91.5	3.3–3.9	1.4–1.9	0.5–0.8	0.9–1.3	0.4–0.7	1.3–1.8	8.5–9.5
Black	%	94.5	2.2	1.2	0.5	0.5	0.4	0.7	5.5
(n=1,408)	CI	92.9–95.8	1.4–3.3	0.7–2.0	0.2–1.1	0.2–1.7	0.1–0.9	0.4–1.5	4.2–7.1
Hispanic	%	94.3	2.7	0.6	0.7	0.7	0.1	0.8	5.7
(n=961)	CI	92.6–95.6	1.9–3.9	0.3–1.3	0.3–1.4	0.3–1.7	0.0–0.7	0.4–1.8	4.4–7.4
8th	%	94.7	2.7	0.9	0.4	0.6	0.2	0.5	5.3
(n=9,476)	CI	94.2–95.2	2.4–3.0	0.7–1.1	0.3–0.6	0.4–0.8	0.2–0.3	0.4–0.6	4.8–5.8
10th	%	91.2	3.3	1.5	0.7	0.9	0.7	1.6	8.8
(n=6,799)	CI	90.4–92.0	2.9–3.8	1.3–1.9	0.5–0.9	0.7–1.2	0.5–0.9	1.3–2.0	8.0–9.6
12th	%	88.0	4.4	2.3	0.8	1.5	0.7	2.3	12.0
(n=5,201)	CI	86.9–89.0	3.8–5.1	1.9–2.8	0.6–1.1	1.1–2.0	0.5–1.0	1.8–2.9	11.0–13.1

Note: The confidence interval (CI) is 95 percent.

Table TN.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	80.9	5.6	3.4	1.6	2.3	1.6	4.6	19.1
(n=21,427)	CI	80.3–81.5	5.2–5.9	3.1–3.7	1.4–1.8	2.1–2.5	1.5–1.9	4.3–5.0	18.5–19.7
Male	%	80.0	5.6	3.3	1.4	2.2	2.0	5.4	20.0
(n=10,448)	CI	79.1–80.9	5.1–6.1	2.9–3.7	1.2–1.7	1.9–2.6	1.7–2.4	4.9–6.0	19.1–20.9
Female	%	82.2	5.5	3.5	1.6	2.3	1.2	3.7	17.8
(n=10,402)	CI	81.3–83.0	5.0–6.0	3.1–3.9	1.4–1.9	2.0–2.7	1.0–1.5	3.2–4.2	17.0–18.7
White	%	79.8	5.8	3.4	1.6	2.5	1.9	5.0	20.2
(n=16,480)	CI	79.1–80.5	5.4–6.2	3.1–3.8	1.4–1.8	2.2–2.8	1.6–2.1	4.6–5.4	19.5–20.9
Black	%	88.7	3.6	1.7	1.0	1.3	0.3	3.5	11.3
(n=1,408)	CI	86.5–90.5	2.7–4.7	1.0–2.7	0.6–1.7	0.8–2.1	0.1–0.8	2.4–5.3	9.5–13.5
Hispanic	%	88.0	3.6	3.4	0.7	1.6	1.2	1.5	12.0
(n=959)	CI	85.5–90.1	2.6–5.0	2.2–5.1	0.3–1.6	1.0–2.6	0.5–2.8	0.9–2.4	9.9–14.5
8th	%	89.3	4.2	2.2	0.8	1.6	0.8	1.1	10.7
(n=9,457)	CI	88.6–90.0	3.8–4.7	1.9–2.6	0.6–1.0	1.3–1.9	0.6–1.0	0.9–1.3	10.0–11.4
10th	%	77.7	6.7	4.4	1.9	2.5	1.9	4.9	22.3
(n=6,781)	CI	76.5–78.8	6.1–7.4	3.9–5.0	1.6–2.3	2.2–3.0	1.5–2.3	4.3–5.5	21.2–23.5
12th	%	75.0	5.8	3.5	2.1	2.9	2.4	8.3	25.0
(n=5,189)	CI	73.6–76.3	5.2–6.6	3.0–4.2	1.7–2.5	2.4–3.5	1.9–2.9	7.5–9.3	23.7–26.4

Note: The confidence interval (CI) is 95 percent.

Table TN.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	94.5	1.6	0.7	0.4	0.6	0.4	1.8	5.5
(n=21,499)	CI	94.1–94.9	1.4–1.8	0.6–0.9	0.3–0.5	0.5–0.7	0.3–0.6	1.6–2.0	5.1–5.9
Male	%	91.3	2.1	1.2	0.5	1.0	0.8	3.1	8.7
(n=10,485)	CI	90.7–92.0	1.8–2.4	1.0–1.5	0.4–0.7	0.8–1.2	0.6–1.0	2.7–3.6	8.0–9.3
Female	%	98.0	0.9	0.2	0.2	0.1	0.1	0.3	2.0
(n=10,431)	CI	97.7–98.3	0.7–1.2	0.2–0.4	0.1–0.4	0.1–0.3	0.0–0.2	0.2–0.5	1.7–2.3
White	%	94.0	1.6	0.8	0.4	0.6	0.5	2.1	6.0
(n=16,546)	CI	93.5–94.4	1.4–1.9	0.6–1.0	0.3–0.5	0.5–0.8	0.4–0.7	1.8–2.3	5.6–6.5
Black	%	97.6	1.0	0.2	0.3	0.4	0.2	0.3	2.4
(n=1,408)	CI	96.4–98.4	0.5–1.7	0.1–0.7	0.1–0.9	0.1–1.7	0.0–0.8	0.1–0.7	1.6–3.6
Hispanic	%	97.4	1.6	0.1	0.1	0.1	0.0	0.6	2.6
(n=957)	CI	96.2–98.3	1.0–2.7	0.0–0.7	0.0–0.9	0.0–1.0	–	0.3–1.3	1.7–3.8
8th	%	97.0	1.1	0.5	0.3	0.2	0.2	0.6	3.0
(n=9,485)	CI	96.6–97.3	0.9–1.4	0.4–0.7	0.2–0.4	0.1–0.3	0.1–0.4	0.5–0.8	2.7–3.4
10th	%	93.8	1.9	0.8	0.4	0.8	0.5	1.8	6.2
(n=6,806)	CI	93.2–94.4	1.6–2.3	0.6–1.1	0.3–0.6	0.6–1.1	0.3–0.7	1.5–2.2	5.6–6.8
12th	%	92.4	1.7	0.8	0.5	0.8	0.6	3.1	7.6
(n=5,208)	CI	91.5–93.3	1.4–2.2	0.6–1.2	0.3–0.7	0.6–1.1	0.4–1.0	2.6–3.7	6.7–8.5

Note: The confidence interval (CI) is 95 percent.

Table TN.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	88.8	3.8	2.0	1.0	1.3	1.0	2.0	11.2
(n=21,451)	CI	88.2–89.3	3.5–4.1	1.8–2.3	0.9–1.2	1.1–1.5	0.9–1.2	1.8–2.3	10.7–11.8
Male	%	89.1	3.2	1.8	1.0	1.3	1.1	2.5	10.9
(n=10,470)	CI	88.3–89.7	2.8–3.6	1.6–2.2	0.8–1.3	1.1–1.6	0.9–1.4	2.2–2.9	10.3–11.7
Female	%	88.9	4.4	2.2	1.1	1.2	0.9	1.4	11.1
(n=10,404)	CI	88.1–89.6	4.0–4.9	1.8–2.5	0.9–1.3	1.0–1.5	0.7–1.1	1.2–1.8	10.4–11.9
White	%	89.7	3.5	1.8	0.9	1.3	1.0	1.8	10.3
(n=16,521)	CI	89.1–90.3	3.2–3.9	1.6–2.1	0.8–1.1	1.1–1.5	0.8–1.2	1.6–2.1	9.7–10.9
Black	%	82.2	5.3	3.0	1.6	1.9	1.5	4.4	17.8
(n=1,396)	CI	79.7–84.5	4.1–6.9	2.1–4.2	0.9–2.7	1.3–2.9	0.9–2.4	3.2–6.2	15.5–20.3
Hispanic	%	91.9	3.6	1.6	0.8	0.4	0.7	1.0	8.1
(n=958)	CI	89.8–93.6	2.5–5.1	1.0–2.8	0.3–1.6	0.1–1.0	0.3–1.6	0.5–2.0	6.4–10.2
8th	%	95.0	2.1	1.0	0.4	0.7	0.4	0.5	5.0
(n=9,475)	CI	94.5–95.5	1.8–2.4	0.8–1.2	0.3–0.5	0.5–0.9	0.3–0.5	0.3–0.6	4.5–5.5
10th	%	86.9	4.6	2.3	1.3	1.4	1.1	2.3	13.1
(n=6,781)	CI	86.0–87.8	4.1–5.2	2.0–2.8	1.0–1.7	1.1–1.8	0.9–1.4	1.9–2.7	12.2–14.0
12th	%	83.7	4.9	2.8	1.5	1.9	1.7	3.5	16.3
(n=5,195)	CI	82.5–84.8	4.3–5.6	2.4–3.4	1.2–1.8	1.5–2.3	1.3–2.2	3.0–4.2	15.2–17.5

Note: The confidence interval (CI) is 95 percent.

Table TN.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.3	0.1	0.1	0.1	0.0	0.1	0.8
(n=21,536)	CI	99.0–99.3	0.2–0.4	0.1–0.2	0.0–0.1	0.0–0.2	0.0–0.1	0.1–0.2	0.7–1.0
Male	%	99.1	0.3	0.2	0.1	0.1	0.1	0.1	0.9
(n=10,514)	CI	98.8–99.4	0.2–0.5	0.1–0.4	0.0–0.2	0.1–0.4	0.0–0.1	0.1–0.2	0.6–1.2
Female	%	99.6	0.3	0.1	0.0	0.0	0.0	0.0	0.4
(n=10,438)	CI	99.4–99.7	0.2–0.4	0.0–0.1	–	0.0–0.1	0.0–0.1	0.0–0.1	0.3–0.6
White	%	99.3	0.3	0.1	0.0	0.1	0.0	0.1	0.7
(n=16,571)	CI	99.1–99.5	0.2–0.5	0.1–0.2	0.0–0.1	0.0–0.2	–	0.0–0.1	0.5–0.9
Black	%	98.8	0.1	0.3	0.2	0.1	0.1	0.4	1.2
(n=1,412)	CI	98.0–99.3	0.0–0.5	0.1–0.9	0.1–0.5	0.0–0.4	0.0–0.7	0.2–1.1	0.7–2.0
Hispanic	%	99.3	0.1	0.3	0.1	0.1	0.0	0.1	0.7
(n=960)	CI	98.5–99.7	0.0–0.7	0.1–1.1	0.0–0.5	0.0–0.7	–	0.0–1.0	0.3–1.5
8th	%	99.7	0.1	0.1	0.1	0.0	0.0	0.0	0.3
(n=9,494)	CI	99.6–99.8	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.2–0.4
10th	%	99.2	0.3	0.1	0.1	0.1	0.0	0.1	0.8
(n=6,821)	CI	98.9–99.4	0.2–0.5	0.1–0.4	0.0–0.2	0.0–0.2	0.0–0.1	0.1–0.3	0.6–1.1
12th	%	98.6	0.6	0.2	0.1	0.2	0.0	0.3	1.4
(n=5,221)	CI	98.1–98.9	0.3–0.9	0.1–0.4	0.0–0.3	0.1–0.6	0.0–0.2	0.1–0.5	1.1–1.9

Note: The confidence interval (CI) is 95 percent.

Table TN.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.3	0.3	0.1	0.1	0.0	0.1	0.2	0.7
(n=21,534)	CI	99.2–99.4	0.2–0.4	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.1–0.2	0.6–0.8
Male	%	99.3	0.3	0.2	0.1	0.0	0.0	0.1	0.7
(n=10,513)	CI	99.1–99.5	0.2–0.5	0.1–0.3	0.0–0.2	0.0–0.1	0.0–0.1	0.1–0.2	0.5–0.9
Female	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=10,437)	CI	99.6–99.8	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.2–0.4
White	%	99.5	0.2	0.1	0.1	0.0	0.0	0.1	0.5
(n=16,570)	CI	99.4–99.6	0.2–0.4	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.1–0.1	0.4–0.6
Black	%	98.7	0.1	0.2	0.2	0.1	0.3	0.3	1.3
(n=1,412)	CI	97.9–99.2	0.0–0.5	0.1–0.7	0.1–0.7	0.0–0.5	0.1–0.9	0.1–0.9	0.8–2.1
Hispanic	%	99.3	0.3	0.3	0.0	0.0	0.0	0.1	0.7
(n=960)	CI	98.5–99.6	0.1–0.9	0.1–0.8	–	–	–	0.0–1.0	0.4–1.5
8th	%	99.6	0.1	0.1	0.1	0.0	0.1	0.1	0.4
(n=9,493)	CI	99.4–99.7	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.2	0.3–0.6
10th	%	99.2	0.3	0.2	0.1	0.0	0.0	0.2	0.8
(n=6,823)	CI	98.9–99.4	0.2–0.6	0.1–0.3	0.0–0.2	0.0–0.1	0.0–0.1	0.1–0.3	0.6–1.1
12th	%	99.1	0.4	0.1	0.1	0.1	0.1	0.2	0.9
(n=5,218)	CI	98.7–99.3	0.2–0.6	0.1–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.1–0.4	0.7–1.3

Note: The confidence interval (CI) is 95 percent.

Table TN.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.9	0.5	0.2	0.1	0.1	0.1	0.1	1.1
(n=21,530)	CI	98.7–99.1	0.4–0.6	0.1–0.3	0.1–0.3	0.1–0.1	0.0–0.2	0.1–0.2	0.9–1.3
Male	%	98.6	0.6	0.3	0.2	0.1	0.1	0.1	1.4
(n=10,514)	CI	98.2–98.8	0.5–0.8	0.2–0.4	0.1–0.5	0.1–0.2	0.0–0.4	0.1–0.2	1.2–1.8
Female	%	99.6	0.2	0.1	0.0	0.0	0.0	0.0	0.4
(n=10,435)	CI	99.5–99.7	0.1–0.3	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.3–0.5
White	%	99.1	0.4	0.2	0.1	0.1	0.1	0.1	0.9
(n=16,568)	CI	98.9–99.3	0.3–0.5	0.1–0.3	0.1–0.3	0.0–0.1	0.0–0.2	0.0–0.1	0.7–1.1
Black	%	98.8	0.2	0.2	0.3	0.1	0.1	0.3	1.2
(n=1,412)	CI	98.0–99.3	0.1–0.5	0.1–0.7	0.1–0.8	0.0–0.9	0.0–0.7	0.1–0.9	0.7–2.0
Hispanic	%	98.9	0.5	0.2	0.1	0.1	0.0	0.1	1.1
(n=960)	CI	97.9–99.4	0.2–1.3	0.1–1.1	0.0–0.5	0.0–0.7	–	0.0–1.0	0.6–2.1
8th	%	99.7	0.1	0.1	0.1	0.0	0.0	0.1	0.3
(n=9,492)	CI	99.6–99.8	0.1–0.2	0.0–0.1	0.0–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.2–0.4
10th	%	99.0	0.4	0.1	0.1	0.2	0.0	0.1	1.0
(n=6,822)	CI	98.7–99.2	0.3–0.6	0.1–0.2	0.0–0.3	0.1–0.3	0.0–0.1	0.1–0.3	0.8–1.3
12th	%	97.9	0.9	0.4	0.2	0.1	0.2	0.2	2.1
(n=5,216)	CI	97.4–98.4	0.7–1.2	0.3–0.7	0.1–0.6	0.0–0.2	0.1–0.6	0.1–0.4	1.6–2.6

Note: The confidence interval (CI) is 95 percent.

Table TN.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.5	0.1	0.1	0.0	0.1	0.0	0.1	0.5
(n=21,533)	CI	99.3–99.6	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.2	0.0–0.1	0.1–0.2	0.4–0.7
Male	%	99.5	0.2	0.1	0.0	0.1	0.0	0.1	0.5
(n=10,512)	CI	99.3–99.7	0.1–0.3	0.0–0.2	0.0–0.1	0.0–0.4	0.0–0.1	0.1–0.2	0.3–0.7
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=10,438)	CI	99.6–99.9	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.3	0.0–0.1	0.1–0.4
White	%	99.6	0.1	0.1	0.0	0.1	0.0	0.1	0.4
(n=16,572)	CI	99.5–99.7	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.2	0.0–0.2	0.0–0.1	0.3–0.5
Black	%	99.0	0.2	0.0	0.1	0.2	0.1	0.4	1.0
(n=1,411)	CI	98.2–99.4	0.0–0.6	–	0.0–0.5	0.1–0.7	0.0–0.9	0.2–0.9	0.6–1.8
Hispanic	%	99.4	0.1	0.2	0.0	0.2	0.0	0.1	0.6
(n=959)	CI	98.7–99.7	0.0–0.7	0.0–0.7	–	0.0–0.7	–	0.0–1.0	0.3–1.3
8th	%	99.8	0.1	0.0	0.0	0.0	0.0	0.1	0.2
(n=9,493)	CI	99.6–99.8	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.2–0.4
10th	%	99.5	0.2	0.1	0.0	0.0	0.1	0.1	0.5
(n=6,823)	CI	99.2–99.6	0.1–0.4	0.0–0.2	0.0–0.1	0.0–0.1	0.0–0.4	0.1–0.2	0.4–0.8
12th	%	99.2	0.2	0.1	0.1	0.2	0.0	0.3	0.8
(n=5,217)	CI	98.8–99.4	0.1–0.3	0.1–0.3	0.0–0.2	0.0–0.6	0.0–0.2	0.1–0.4	0.6–1.2

Note: The confidence interval (CI) is 95 percent.

Table TN.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.5	0.1	0.1	0.1	0.0	0.0	0.1	0.5
(n=21,529)	CI	99.4–99.6	0.1–0.2	0.0–0.1	0.1–0.1	0.0–0.1	0.0–0.1	0.1–0.2	0.4–0.6
Male	%	99.6	0.1	0.1	0.1	0.0	0.0	0.1	0.4
(n=10,511)	CI	99.4–99.7	0.0–0.2	0.0–0.2	0.0–0.2	0.0–0.1	0.0–0.1	0.1–0.2	0.3–0.6
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=10,438)	CI	99.6–99.9	0.0–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.1–0.4
White	%	99.7	0.1	0.1	0.0	0.0	0.0	0.1	0.3
(n=16,569)	CI	99.6–99.8	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	–	0.0–0.1	0.2–0.4
Black	%	99.3	0.1	0.0	0.1	0.0	0.2	0.4	0.7
(n=1,411)	CI	98.5–99.6	0.0–0.6	–	0.0–0.5	–	0.0–0.7	0.1–1.0	0.4–1.5
Hispanic	%	99.4	0.1	0.0	0.2	0.0	0.1	0.2	0.6
(n=959)	CI	98.7–99.7	0.0–0.7	–	0.0–0.7	–	0.0–0.7	0.1–0.9	0.3–1.3
8th	%	99.7	0.1	0.0	0.0	0.0	0.0	0.0	0.3
(n=9,490)	CI	99.6–99.8	0.1–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.1	0.2–0.4
10th	%	99.5	0.2	0.1	0.1	0.1	0.0	0.1	0.5
(n=6,822)	CI	99.2–99.6	0.1–0.4	0.0–0.2	0.1–0.2	0.0–0.1	0.0–0.1	0.1–0.2	0.4–0.8
12th	%	99.3	0.1	0.1	0.1	0.0	0.0	0.3	0.7
(n=5,217)	CI	99.0–99.5	0.1–0.3	0.0–0.3	0.0–0.3	0.0–0.1	0.0–0.2	0.2–0.5	0.5–1.0

Note: The confidence interval (CI) is 95 percent.

Table TN.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you use ecstasy (MDMA)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.2	0.3	0.1	0.1	0.0	0.0	0.1	0.8
(n=21,530)	CI	99.1-99.4	0.3-0.4	0.1-0.2	0.1-0.2	0.0-0.1	0.0-0.2	0.1-0.2	0.6-0.9
Male	%	99.0	0.4	0.1	0.1	0.1	0.1	0.1	1.0
(n=10,513)	CI	98.7-99.3	0.3-0.6	0.1-0.3	0.0-0.4	0.0-0.1	0.0-0.4	0.1-0.2	0.7-1.3
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=10,438)	CI	99.7-99.9	0.1-0.2	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.1-0.3
White	%	99.4	0.3	0.1	0.1	0.0	0.0	0.1	0.6
(n=16,569)	CI	99.2-99.6	0.2-0.4	0.0-0.2	0.0-0.2	0.0-0.1	0.0-0.3	0.0-0.1	0.4-0.8
Black	%	99.0	0.1	0.1	0.2	0.1	0.0	0.5	1.0
(n=1,410)	CI	98.2-99.4	0.0-0.6	0.0-0.6	0.1-0.5	0.0-0.4	-	0.2-1.2	0.6-1.8
Hispanic	%	99.2	0.5	0.2	0.0	0.0	0.0	0.1	0.8
(n=960)	CI	98.2-99.6	0.2-1.4	0.0-0.6	-	-	-	0.0-1.0	0.4-1.8
8th	%	99.7	0.1	0.0	0.0	0.0	0.0	0.0	0.3
(n=9,491)	CI	99.6-99.8	0.0-0.2	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.2-0.4
10th	%	99.4	0.3	0.1	0.1	0.0	0.0	0.1	0.6
(n=6,822)	CI	99.1-99.6	0.2-0.5	0.0-0.3	0.0-0.2	0.0-0.1	0.0-0.1	0.0-0.2	0.4-0.9
12th	%	98.5	0.6	0.2	0.2	0.0	0.1	0.3	1.5
(n=5,217)	CI	98.0-98.9	0.5-0.9	0.1-0.3	0.1-0.6	0.0-0.2	0.0-0.8	0.2-0.5	1.1-2.0

Note: The confidence interval (CI) is 95 percent.

Table TN.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you use methamphetamines (crank, etc.)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.5	0.1	0.1	0.0	0.1	0.0	0.2	0.5
(n=21,531)	CI	99.3-99.5	0.1-0.2	0.1-0.2	0.0-0.1	0.0-0.1	-	0.1-0.2	0.5-0.7
Male	%	99.4	0.2	0.1	0.1	0.1	0.0	0.1	0.6
(n=10,514)	CI	99.2-99.6	0.1-0.3	0.1-0.2	0.0-0.1	0.0-0.2	0.0-0.1	0.1-0.2	0.4-0.8
Female	%	99.8	0.0	0.0	0.0	0.0	0.0	0.1	0.2
(n=10,437)	CI	99.7-99.9	0.0-0.1	0.0-0.1	-	0.0-0.1	0.0-0.1	0.0-0.1	0.1-0.3
White	%	99.6	0.1	0.1	0.0	0.0	0.0	0.1	0.4
(n=16,569)	CI	99.5-99.7	0.1-0.2	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.3-0.5
Black	%	99.0	0.1	0.1	0.1	0.1	0.0	0.6	1.0
(n=1,410)	CI	98.2-99.5	0.0-0.9	0.0-0.6	0.0-0.5	0.0-0.7	-	0.2-1.3	0.5-1.8
Hispanic	%	99.4	0.2	0.1	0.1	0.1	0.0	0.1	0.6
(n=962)	CI	98.7-99.7	0.0-0.7	0.0-0.7	0.0-0.5	0.0-0.6	-	0.0-1.0	0.3-1.3
8th	%	99.8	0.0	0.0	0.0	0.1	0.0	0.1	0.2
(n=9,486)	CI	99.6-99.8	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.1	0.2-0.4
10th	%	99.4	0.2	0.1	0.1	0.0	0.0	0.2	0.6
(n=6,820)	CI	99.2-99.5	0.1-0.3	0.1-0.3	0.0-0.2	0.0-0.1	0.0-0.1	0.1-0.3	0.5-0.8
12th	%	99.2	0.2	0.1	0.0	0.1	0.0	0.3	0.8
(n=5,225)	CI	98.9-99.4	0.1-0.4	0.1-0.3	0.0-0.1	0.0-0.2	0.0-0.1	0.2-0.5	0.6-1.1

Note: The confidence interval (CI) is 95 percent.

Table TN.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	24.1	23.9	16.9	35.1	52.0
(n=8,200)	CI	23.0–25.1	22.9–25.0	16.0–17.8	33.9–36.3	50.8–53.2
Male	%	23.0	21.5	17.4	38.2	55.5
(n=3,744)	CI	21.5–24.6	20.0–23.0	16.0–18.8	36.4–40.0	53.7–57.4
Female	%	23.3	26.8	16.8	33.2	49.9
(n=4,140)	CI	21.9–24.8	25.2–28.4	15.5–18.1	31.6–34.8	48.2–51.7
White	%	22.5	24.2	17.6	35.6	53.3
(n=6,460)	CI	21.4–23.7	23.0–25.4	16.6–18.7	34.3–37.0	51.9–54.6
Black	%	31.5	20.9	14.9	32.6	47.6
(n=426)	CI	27.0–36.5	17.0–25.4	11.5–19.2	28.1–37.5	42.5–52.7
Hispanic	%	23.9	18.7	20.4	37.1	57.5
(n=279)	CI	18.6–30.1	14.1–24.3	15.7–26.0	30.8–43.8	50.8–63.9
8th	%	19.9	21.4	13.6	45.1	58.7
(n=3,490)	CI	18.5–21.3	20.0–22.9	12.5–14.9	43.3–46.9	57.0–60.5
10th	%	26.0	25.3	18.0	30.8	48.8
(n=2,629)	CI	24.1–27.9	23.4–27.2	16.4–19.6	28.8–32.9	46.6–51.0
12th	%	26.5	25.2	19.1	29.2	48.3
(n=2,081)	CI	24.4–28.6	23.2–27.4	17.3–21.1	26.9–31.5	45.9–50.7

Note: The confidence interval (CI) is 95 percent.

Table TN.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	25.7	36.0	17.9	20.4	38.3
(n=11,768)	CI	24.8–26.6	35.1–37.0	17.1–18.7	19.5–21.2	37.3–39.3
Male	%	25.4	34.9	18.1	21.6	39.7
(n=5,494)	CI	24.1–26.8	33.4–36.3	17.0–19.4	20.4–22.9	38.3–41.2
Female	%	24.8	37.8	18.1	19.3	37.4
(n=5,870)	CI	23.6–26.1	36.4–39.3	16.9–19.2	18.2–20.5	36.0–38.8
White	%	24.5	37.1	18.1	20.3	38.4
(n=9,319)	CI	23.5–25.5	36.0–38.2	17.2–19.0	19.4–21.3	37.3–39.6
Black	%	31.7	30.1	13.4	24.8	38.2
(n=589)	CI	27.7–36.0	25.9–34.6	10.9–16.5	21.3–28.7	34.1–42.5
Hispanic	%	24.9	34.2	19.9	21.0	40.9
(n=428)	CI	20.5–29.8	29.3–39.5	15.7–24.8	16.9–25.8	35.6–46.4
8th	%	19.6	30.4	19.7	30.3	50.0
(n=4,475)	CI	18.4–20.9	29.0–31.9	18.5–21.0	28.8–31.7	48.4–51.5
10th	%	24.6	38.9	18.9	17.6	36.5
(n=3,934)	CI	23.1–26.2	37.2–40.6	17.5–20.3	16.3–19.0	34.8–38.2
12th	%	31.9	37.8	15.4	14.8	30.3
(n=3,359)	CI	30.1–33.7	36.0–39.7	14.0–17.0	13.5–16.3	28.5–32.1

Note: The confidence interval (CI) is 95 percent.

Table TN.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	39.9	27.8	12.4	19.9	32.3
(n=11,413)	CI	38.9–41.0	26.8–28.7	11.7–13.1	19.1–20.8	31.3–33.3
Male	%	42.0	26.3	12.3	19.4	31.7
(n=5,551)	CI	40.6–43.5	24.9–27.6	11.4–13.3	18.3–20.6	30.4–33.1
Female	%	36.8	29.9	12.6	20.7	33.3
(n=5,482)	CI	35.3–38.3	28.5–31.3	11.6–13.7	19.5–21.9	31.9–34.8
White	%	40.4	28.4	12.2	19.0	31.3
(n=9,197)	CI	39.2–41.5	27.3–29.5	11.5–13.0	18.2–20.0	30.2–32.3
Black	%	38.3	23.6	11.0	27.1	38.1
(n=519)	CI	33.7–43.0	19.9–27.8	8.0–15.1	23.2–31.3	33.6–42.9
Hispanic	%	33.6	27.0	13.1	26.2	39.4
(n=367)	CI	28.2–39.4	22.3–32.4	9.8–17.5	21.3–31.8	33.8–45.3
8th	%	23.9	26.3	16.3	33.4	49.7
(n=4,193)	CI	22.6–25.3	25.0–27.8	15.2–17.6	31.9–35.0	48.1–51.4
10th	%	37.2	32.2	13.3	17.3	30.6
(n=3,849)	CI	35.5–38.9	30.6–33.9	12.1–14.6	15.9–18.6	28.9–32.2
12th	%	55.3	24.5	8.3	11.9	20.2
(n=3,371)	CI	53.4–57.3	22.8–26.2	7.2–9.5	10.6–13.2	18.6–21.8

Note: The confidence interval (CI) is 95 percent.

Table TN.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	28.7	25.2	14.2	32.0	46.1
(n=9,332)	CI	27.7–29.8	24.2–26.2	13.4–15.0	30.9–33.0	45.0–47.3
Male	%	29.2	23.8	15.0	32.0	47.0
(n=4,418)	CI	27.7–30.8	22.4–25.3	13.8–16.2	30.5–33.5	45.3–48.6
Female	%	27.2	26.5	13.6	32.7	46.3
(n=4,580)	CI	25.7–28.7	25.1–28.1	12.6–14.8	31.2–34.2	44.7–47.9
White	%	27.0	24.9	14.8	33.4	48.1
(n=7,246)	CI	25.8–28.2	23.8–26.1	13.9–15.7	32.2–34.6	46.8–49.5
Black	%	44.4	24.5	8.7	22.4	31.0
(n=570)	CI	40.0–49.0	20.7–28.9	6.5–11.4	19.1–26.0	27.2–35.1
Hispanic	%	23.8	29.1	15.6	31.5	47.0
(n=349)	CI	19.2–29.1	23.8–35.2	11.8–20.3	26.2–37.3	41.1–53.1
8th	%	12.9	14.2	14.5	58.5	72.9
(n=3,252)	CI	11.7–14.2	13.0–15.5	13.2–15.8	56.6–60.3	71.3–74.5
10th	%	31.5	27.7	15.1	25.7	40.8
(n=3,278)	CI	29.7–33.3	26.0–29.4	13.8–16.5	24.1–27.5	39.0–42.8
12th	%	37.6	30.8	13.0	18.6	31.6
(n=2,802)	CI	35.6–39.7	28.8–32.8	11.7–14.4	17.1–20.3	29.7–33.6

Note: The confidence interval (CI) is 95 percent.

Table TN.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	1.9	98.1
(n=21,490)	CI	1.7–2.2	97.8–98.3
Male	%	1.7	98.3
(n=10,505)	CI	1.4–2.1	97.9–98.6
Female	%	1.8	98.2
(n=10,408)	CI	1.5–2.2	97.8–98.5
White	%	1.7	98.3
(n=16,537)	CI	1.4–1.9	98.1–98.6
Black	%	2.5	97.5
(n=1,401)	CI	1.7–3.5	96.5–98.3
Hispanic	%	1.7	98.3
(n=963)	CI	1.1–2.9	97.1–98.9
8th	%	1.1	98.9
(n=9,483)	CI	0.9–1.4	98.6–99.1
10th	%	2.3	97.7
(n=6,798)	CI	1.9–2.7	97.3–98.1
12th	%	2.5	97.5
(n=5,209)	CI	2.0–3.0	97.0–98.0

Note: The confidence interval (CI) is 95 percent.

Table TN.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	14.5	85.5
(n=21,374)	CI	13.9–15.0	85.0–86.1
Male	%	13.6	86.4
(n=10,423)	CI	12.8–14.4	85.6–87.2
Female	%	15.1	84.9
(n=10,374)	CI	14.2–15.9	84.1–85.8
White	%	14.7	85.3
(n=16,455)	CI	14.1–15.4	84.6–85.9
Black	%	12.9	87.1
(n=1,401)	CI	10.9–15.2	84.8–89.1
Hispanic	%	11.9	88.1
(n=952)	CI	9.7–14.5	85.5–90.3
8th	%	6.4	93.6
(n=9,436)	CI	5.9–6.9	93.1–94.1
10th	%	15.9	84.1
(n=6,780)	CI	14.9–16.9	83.1–85.1
12th	%	22.1	77.9
(n=5,158)	CI	20.8–23.4	76.6–79.2

Note: The confidence interval (CI) is 95 percent.

Table TN.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=383)	Male (n=162)	Female (n=180)	White (n=252)	Black (n=35)	Hispanic (n=16)	8th (n=107)	10th (n=157)	12th (n=119)
Got a prescription from one doctor	%	15.7	18.1	12.9	14.9	13.0	15.1	15.9	13.8	17.6
	CI	11.3–21.3	10.9–28.7	7.8–20.6	9.5–22.6	5.1–29.5	4.2–42.3	10.0–24.4	8.2–22.1	9.7–29.8
Got a prescription from more than one doctor	%	6.3	5.7	4.5	5.2	7.7	9.9	15.6	3.7	4.1
	CI	4.3–9.0	3.2–9.7	2.4–8.0	3.3–8.3	1.6–29.5	2.0–37.0	9.9–23.8	1.7–7.7	1.7–9.5
Bought on the Internet	%	10.0	13.2	2.1	6.4	17.7	13.7	7.1	7.8	13.8
	CI	7.2–13.7	8.4–20.1	0.7–5.9	3.8–10.4	6.6–39.6	2.7–47.2	3.5–13.8	4.4–13.4	8.4–21.8
Bought from a friend, relative, or someone you know	%	35.0	37.3	35.0	37.3	32.7	27.2	36.5	35.2	34.0
	CI	29.7–40.7	28.5–46.9	27.8–43.0	30.4–44.7	18.4–51.0	9.8–56.1	27.6–46.4	27.5–43.7	24.4–45.0
Got from a friend, relative, or someone you know for free	%	50.1	47.2	52.9	52.3	38.9	51.5	54.1	50.2	48.0
	CI	44.3–55.9	37.9–56.6	44.5–61.1	44.8–59.6	22.9–57.7	25.6–76.7	44.2–63.7	41.3–59.0	37.6–58.6
Took from a friend, relative, or someone you know without asking	%	22.9	18.3	25.7	24.4	12.2	14.7	28.2	23.1	19.8
	CI	18.0–28.6	11.2–28.5	18.7–34.2	18.0–32.2	4.3–30.3	4.0–41.5	20.5–37.5	15.7–32.7	11.9–31.1
Took from someone's house or workplace without asking	%	8.9	5.8	8.1	5.8	4.9	14.2	13.4	9.8	5.6
	CI	6.5–12.1	3.1–10.7	4.9–13.1	3.5–9.4	1.1–18.9	3.9–40.3	8.1–21.4	5.8–15.9	2.9–10.4
Got it at a party	%	27.7	31.3	23.1	24.7	39.7	18.1	24.7	28.7	28.2
	CI	22.7–33.4	22.8–41.2	17.1–30.5	18.7–31.9	23.5–58.5	4.8–49.1	17.1–34.1	21.6–37.1	19.2–39.5
Got it some other way	%	31.6	39.9	21.6	28.5	28.4	47.5	28.0	27.3	38.2
	CI	26.1–37.7	30.4–50.2	15.8–28.8	21.6–36.5	14.6–47.9	22.5–73.8	20.0–37.6	20.4–35.5	27.6–49.9

Note: The confidence interval (CI) is 95 percent.

Table TN.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=2,791)	Male (n=1,274)	Female (n=1,402)	White (n=2,177)	Black (n=160)	Hispanic (n=104)	8th (n=602)	10th (n=1,072)	12th (n=1,117)
I bought it in a store	%	4.5	5.6	2.7	3.7	7.5	9.3	2.4	3.3	6.3
	CI	3.7–5.5	4.4–7.1	1.9–4.0	2.9–4.7	4.1–13.3	4.2–19.5	1.4–3.9	2.4–4.5	4.9–8.0
I bought it in a gas station	%	5.8	6.9	3.9	5.4	6.7	4.3	3.1	3.6	8.4
	CI	4.8–6.9	5.4–8.8	2.8–5.3	4.3–6.7	3.6–12.1	1.7–10.4	2.0–4.8	2.7–4.9	6.6–10.6
I bought it in a restaurant	%	1.9	1.6	1.4	1.4	2.8	3.8	1.1	1.2	2.6
	CI	1.2–2.8	0.8–3.2	0.7–2.8	0.8–2.6	1.0–7.4	1.4–10.2	0.5–2.4	0.7–2.0	1.5–4.6
I bought it in a bar or club	%	2.5	2.2	2.3	2.0	4.1	7.6	1.3	1.6	3.7
	CI	1.9–3.3	1.4–3.6	1.5–3.4	1.4–2.9	1.8–9.1	3.0–18.0	0.7–2.6	1.0–2.4	2.6–5.4
I bought it at a public event	%	2.7	2.8	1.6	1.8	5.0	6.6	3.0	2.2	3.0
	CI	2.2–3.4	2.0–3.9	1.1–2.4	1.3–2.5	2.5–9.9	3.2–13.2	1.9–4.8	1.5–3.2	2.2–4.2
I gave someone money to buy it for me	%	28.4	30.3	26.8	30.2	18.1	27.7	14.1	26.6	34.6
	CI	26.5–30.4	27.4–33.3	24.2–29.6	28.0–32.5	12.4–25.5	19.1–38.3	11.5–17.2	23.7–29.7	31.4–37.9
I got it from a sibling (under 21)	%	5.5	4.9	5.5	4.9	10.3	7.2	9.3	5.7	4.1
	CI	4.7–6.5	3.8–6.4	4.3–6.9	4.0–6.0	6.3–16.4	3.5–14.5	7.2–11.9	4.3–7.5	3.0–5.4
I got it from a friend (under 21)	%	23.0	19.7	25.5	23.2	23.5	20.7	24.9	25.2	20.6
	CI	21.3–24.8	17.4–22.2	23.0–28.1	21.3–25.2	16.7–31.9	13.7–30.0	21.6–28.6	22.3–28.3	18.2–23.2
I got it at a party	%	43.5	46.2	41.2	41.9	56.3	54.2	27.0	40.9	51.0
	CI	41.3–45.6	43.0–49.4	38.3–44.2	39.5–44.3	47.3–64.8	43.3–64.8	23.5–30.7	37.6–44.3	47.6–54.4
I took it from a store or family member	%	12.0	12.5	10.5	11.0	11.7	16.3	19.1	12.3	9.4
	CI	10.7–13.4	10.6–14.7	8.9–12.4	9.6–12.6	7.5–17.8	10.3–24.7	16.0–22.5	10.3–14.6	7.6–11.6
I took it home without permission	%	17.8	15.4	19.1	17.4	12.4	22.6	26.6	23.7	10.1
	CI	16.3–19.5	13.3–17.8	16.7–21.6	15.7–19.3	7.9–18.9	14.6–33.3	23.2–30.4	20.8–26.8	8.2–12.3
I got it from my parents with permission	%	21.5	18.1	24.0	22.5	13.8	12.8	28.8	23.1	17.8
	CI	19.9–23.3	15.9–20.6	21.6–26.6	20.6–24.5	9.1–20.4	7.4–21.4	25.1–32.8	20.3–26.2	15.5–20.4
I got it from an older sibling (over 21)	%	12.0	10.3	13.2	11.4	17.5	12.7	11.2	12.4	11.8
	CI	10.7–13.4	8.5–12.4	11.3–15.4	10.0–13.0	11.8–25.2	6.6–22.8	8.7–14.3	10.3–14.9	9.9–14.1
I got it from an older friend (over 21)	%	25.3	24.5	26.1	26.1	19.9	22.1	14.3	23.2	30.7
	CI	23.4–27.3	21.7–27.5	23.4–28.9	23.9–28.4	13.7–27.8	14.7–31.8	11.7–17.4	20.5–26.1	27.5–34.0
I got it from some other adult	%	16.6	15.1	17.5	16.4	16.2	15.1	16.4	19.2	14.5
	CI	15.1–18.2	13.0–17.5	15.3–19.8	14.7–18.2	11.2–23.0	8.4–25.7	13.5–19.8	16.6–22.1	12.5–16.8
I got it some other way	%	21.9	26.5	16.8	20.7	27.8	26.0	24.2	22.6	20.5
	CI	20.2–23.7	23.7–29.4	14.7–19.2	18.8–22.8	20.3–36.8	17.6–36.6	20.8–28.0	19.8–25.8	17.9–23.4

Note: The confidence interval (CI) is 95 percent.

Table TN.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?			
I bought or tried to buy alcohol		Yes, asked for proof of age	No, not asked for proof of age
Total	%	3.4	28.0
(n=580)	CI	3.1–3.7	24.1–32.3
Male	%	4.1	27.6
(n=328)	CI	3.6–4.6	22.5–33.4
Female	%	2.1	27.1
(n=184)	CI	1.8–2.5	20.1–35.3
White	%	2.8	29.5
(n=381)	CI	2.5–3.2	24.4–35.1
Black	%	4.8	23.6
(n=49)	CI	3.7–6.3	13.2–38.5
Hispanic	%	4.4	25.5
(n=31)	CI	3.1–6.3	12.4–45.2
8th	%	1.9	19.6
(n=155)	CI	1.6–2.2	13.9–26.8
10th	%	3.3	22.2
(n=196)	CI	2.8–3.8	16.3–29.5
12th	%	5.2	35.2
(n=229)	CI	4.5–6.0	28.8–42.2

Note: The confidence interval (CI) is 95 percent.

Table TN.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?				
I bought or tried to buy alcohol		No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	3.4	36.3	29.5
(n=580)	CI	3.1–3.7	32.0–40.7	25.6–33.7
Male	%	4.1	36.1	34.0
(n=328)	CI	3.6–4.6	30.4–42.3	28.5–40.0
Female	%	2.1	43.1	23.2
(n=184)	CI	1.8–2.5	35.4–51.2	17.2–30.5
White	%	2.8	33.2	30.2
(n=381)	CI	2.5–3.2	28.0–38.9	25.3–35.6
Black	%	4.8	52.9	24.8
(n=49)	CI	3.7–6.3	38.7–66.7	14.4–39.3
Hispanic	%	4.4	49.6	24.3
(n=31)	CI	3.1–6.3	30.7–68.6	11.6–44.0
8th	%	1.9	47.1	20.4
(n=155)	CI	1.6–2.2	39.4–55.0	14.8–27.5
10th	%	3.3	38.9	30.1
(n=196)	CI	2.8–3.8	31.4–47.0	23.9–37.1
12th	%	5.2	30.0	32.8
(n=229)	CI	4.5–6.0	23.8–36.9	26.5–39.9

Note: The confidence interval (CI) is 95 percent.

Table TN.39. Ever Ridden with a Prescription Drugs Impaired Driver

Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		No	Yes	Not sure
Total	%	81.7	8.2	10.2
(n=20,399)	CI	81.1–82.3	7.7–8.6	9.7–10.6
Male	%	83.5	6.7	9.8
(n=9,880)	CI	82.6–84.3	6.2–7.3	9.1–10.4
Female	%	80.7	9.0	10.3
(n=9,965)	CI	79.8–81.6	8.4–9.7	9.6–11.0
White	%	82.1	8.0	10.0
(n=15,787)	CI	81.4–82.7	7.5–8.5	9.5–10.5
Black	%	80.7	8.0	11.2
(n=1,272)	CI	78.2–83.0	6.6–9.8	9.5–13.3
Hispanic	%	87.8	5.0	7.2
(n=891)	CI	85.3–89.9	3.6–6.8	5.7–9.2
8th	%	80.7	7.1	12.2
(n=8,965)	CI	79.8–81.6	6.5–7.7	11.5–13.0
10th	%	80.6	8.9	10.6
(n=6,464)	CI	79.4–81.6	8.1–9.7	9.7–11.4
12th	%	84.1	8.6	7.3
(n=4,970)	CI	82.9–85.2	7.8–9.5	6.6–8.2

Note: The confidence interval (CI) is 95 percent.

Table TN.40. Frequency Riding with a Prescription Drugs Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	95.4	1.5	1.3	0.5	1.3	4.6
(n=21,019)	CI	95.1–95.7	1.3–1.7	1.2–1.5	0.4–0.6	1.2–1.5	4.3–4.9
Male	%	96.0	1.2	1.2	0.4	1.1	4.0
(n=10,223)	CI	95.6–96.5	1.0–1.5	0.9–1.5	0.3–0.7	0.9–1.4	3.5–4.4
Female	%	95.3	1.7	1.4	0.5	1.0	4.7
(n=10,218)	CI	94.8–95.7	1.5–2.0	1.2–1.7	0.4–0.7	0.8–1.3	4.3–5.2
White	%	95.9	1.4	1.2	0.4	1.1	4.1
(n=16,209)	CI	95.5–96.3	1.2–1.6	1.0–1.4	0.3–0.5	0.9–1.3	3.7–4.5
Black	%	93.6	1.6	1.9	0.6	2.2	6.4
(n=1,356)	CI	92.0–95.0	1.1–2.5	1.2–2.9	0.3–1.3	1.5–3.3	5.0–8.0
Hispanic	%	96.9	1.6	0.7	0.5	0.3	3.1
(n=929)	CI	95.4–97.9	0.9–2.8	0.3–1.6	0.2–1.2	0.1–0.8	2.1–4.6
8th	%	96.0	1.4	1.1	0.5	0.9	4.0
(n=9,252)	CI	95.6–96.4	1.2–1.7	0.9–1.3	0.4–0.7	0.8–1.2	3.6–4.4
10th	%	94.9	1.6	1.6	0.4	1.5	5.1
(n=6,652)	CI	94.3–95.5	1.3–2.0	1.3–2.0	0.3–0.6	1.2–1.8	4.5–5.7
12th	%	95.3	1.4	1.3	0.5	1.6	4.7
(n=5,115)	CI	94.6–95.9	1.0–1.8	1.0–1.6	0.3–0.8	1.3–2.0	4.1–5.4

Note: The confidence interval (CI) is 95 percent.

Table TN.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	69.9	20.6	9.5
(n=21,053)	CI	69.2–70.6	20.0–21.2	9.1–10.0
Male	%	71.9	18.3	9.9
(n=10,243)	CI	70.9–72.8	17.4–19.2	9.3–10.5
Female	%	68.7	22.4	8.9
(n=10,237)	CI	67.6–69.7	21.5–23.3	8.3–9.5
White	%	70.5	20.3	9.2
(n=16,226)	CI	69.7–71.3	19.6–21.0	8.7–9.7
Black	%	66.2	22.7	11.1
(n=1,362)	CI	63.4–69.0	20.3–25.3	9.3–13.1
Hispanic	%	75.7	15.0	9.4
(n=935)	CI	72.6–78.5	12.8–17.5	7.5–11.6
8th	%	71.8	16.3	11.8
(n=9,273)	CI	70.9–72.8	15.5–17.1	11.2–12.6
10th	%	68.7	21.8	9.5
(n=6,670)	CI	67.5–70.0	20.7–22.9	8.7–10.3
12th	%	69.0	24.1	6.9
(n=5,110)	CI	67.5–70.4	22.8–25.4	6.2–7.7

Note: The confidence interval (CI) is 95 percent.

Table TN.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	88.9	4.2	3.6	1.0	2.2	11.1
(n=21,057)	CI	88.4–89.4	3.9–4.6	3.3–3.9	0.8–1.2	2.0–2.4	10.6–11.6
Male	%	90.2	3.4	3.2	1.0	2.1	9.8
(n=10,250)	CI	89.5–90.9	3.0–3.9	2.8–3.6	0.8–1.3	1.9–2.5	9.1–10.5
Female	%	88.1	5.1	4.1	1.0	1.7	11.9
(n=10,238)	CI	87.4–88.8	4.6–5.6	3.7–4.5	0.8–1.2	1.5–2.0	11.2–12.6
White	%	89.5	4.0	3.6	1.0	1.8	10.5
(n=16,232)	CI	89.0–90.1	3.7–4.4	3.3–4.0	0.8–1.2	1.6–2.1	9.9–11.0
Black	%	85.6	5.8	3.9	1.2	3.6	14.4
(n=1,365)	CI	83.4–87.5	4.5–7.4	2.8–5.2	0.7–1.9	2.7–4.8	12.5–16.6
Hispanic	%	91.1	4.0	3.6	0.4	0.9	8.9
(n=933)	CI	89.1–92.8	2.9–5.5	2.6–5.0	0.2–1.0	0.5–1.7	7.2–10.9
8th	%	91.4	3.6	2.9	0.7	1.5	8.6
(n=9,278)	CI	90.7–91.9	3.2–4.0	2.5–3.3	0.5–0.9	1.2–1.7	8.1–9.3
10th	%	88.2	4.3	4.2	1.1	2.2	11.8
(n=6,667)	CI	87.3–89.0	3.8–4.9	3.7–4.8	0.8–1.4	1.9–2.6	11.0–12.7
12th	%	87.1	4.8	3.8	1.3	3.0	12.9
(n=5,112)	CI	86.0–88.1	4.2–5.6	3.3–4.5	0.9–1.7	2.5–3.5	11.9–14.0

Note: The confidence interval (CI) is 95 percent.

Table TN.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.8	2.0	5.1	91.1	96.2
(n=20,975)	CI	1.6–2.0	1.8–2.3	4.8–5.5	90.6–91.5	95.9–96.5
Male	%	1.8	1.4	5.0	91.8	96.8
(n=10,206)	CI	1.5–2.1	1.2–1.7	4.5–5.5	91.2–92.4	96.4–97.2
Female	%	1.2	2.5	5.2	91.1	96.3
(n=10,198)	CI	1.0–1.5	2.2–2.9	4.7–5.7	90.4–91.7	95.8–96.7
White	%	1.3	2.0	5.1	91.6	96.7
(n=16,160)	CI	1.1–1.5	1.7–2.2	4.8–5.5	91.1–92.1	96.4–97.1
Black	%	4.2	1.8	5.6	88.5	94.1
(n=1,351)	CI	3.2–5.4	1.2–2.7	4.4–7.0	86.5–90.2	92.6–95.3
Hispanic	%	1.7	1.8	4.4	92.0	96.5
(n=933)	CI	1.0–2.9	1.1–3.1	3.2–6.1	89.9–93.7	94.9–97.6
8th	%	1.3	1.7	4.6	92.4	96.9
(n=9,253)	CI	1.1–1.6	1.5–2.0	4.1–5.0	91.8–92.9	96.5–97.3
10th	%	1.8	2.2	5.2	90.8	96.0
(n=6,635)	CI	1.4–2.1	1.8–2.7	4.7–5.8	90.0–91.6	95.5–96.5
12th	%	2.2	2.1	5.7	89.9	95.6
(n=5,087)	CI	1.9–2.7	1.7–2.7	5.1–6.4	89.0–90.8	94.9–96.2

Note: The confidence interval (CI) is 95 percent.

Table TN.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.4	4.8	10.9	82.0	92.9
(n=21,050)	CI	2.2–2.6	4.4–5.1	10.4–11.4	81.4–82.6	92.5–93.3
Male	%	2.7	5.0	11.3	81.0	92.3
(n=10,239)	CI	2.3–3.1	4.5–5.5	10.6–12.0	80.1–81.9	91.7–92.9
Female	%	1.6	4.5	10.3	83.6	93.9
(n=10,236)	CI	1.4–1.9	4.0–4.9	9.7–11.1	82.7–84.4	93.4–94.4
White	%	2.0	4.8	11.3	81.8	93.1
(n=16,219)	CI	1.8–2.3	4.4–5.2	10.8–11.9	81.1–82.5	92.7–93.6
Black	%	4.1	2.8	8.3	84.8	93.1
(n=1,355)	CI	3.1–5.4	2.0–4.0	6.8–10.1	82.5–86.8	91.4–94.4
Hispanic	%	2.6	2.7	9.2	85.5	94.7
(n=937)	CI	1.7–4.0	1.8–4.0	7.3–11.5	82.8–87.8	93.0–96.0
8th	%	1.2	3.5	8.0	87.3	95.3
(n=9,284)	CI	1.0–1.5	3.1–3.9	7.5–8.6	86.6–88.0	94.8–95.8
10th	%	2.2	4.5	10.9	82.4	93.3
(n=6,662)	CI	1.9–2.7	3.9–5.1	10.1–11.8	81.4–83.4	92.6–93.9
12th	%	3.9	6.6	14.1	75.5	89.6
(n=5,104)	CI	3.3–4.5	5.8–7.4	13.0–15.2	74.1–76.8	88.6–90.5

Note: The confidence interval (CI) is 95 percent.

Table TN.45. Parents Feel Regularly Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.8	8.6	11.9	73.7	85.5
(n=21,026)	CI	5.5–6.2	8.2–9.1	11.4–12.4	73.0–74.3	85.0–86.1
Male	%	6.0	8.0	12.2	73.8	86.0
(n=10,233)	CI	5.5–6.6	7.4–8.6	11.5–12.9	72.8–74.8	85.2–86.8
Female	%	5.2	9.1	11.5	74.1	85.7
(n=10,222)	CI	4.7–5.7	8.5–9.8	10.9–12.3	73.1–75.1	84.8–86.5
White	%	5.6	8.8	12.3	73.3	85.6
(n=16,206)	CI	5.2–6.1	8.3–9.3	11.7–12.8	72.5–74.1	84.9–86.2
Black	%	5.7	6.7	9.1	78.6	87.7
(n=1,353)	CI	4.4–7.2	5.3–8.4	7.5–11.0	76.0–81.0	85.5–89.5
Hispanic	%	5.0	6.5	10.5	77.9	88.5
(n=932)	CI	3.7–6.8	5.0–8.5	8.6–12.8	74.9–80.7	86.0–90.5
8th	%	3.2	5.6	10.3	80.9	91.1
(n=9,276)	CI	2.9–3.6	5.2–6.1	9.6–10.9	80.0–81.7	90.5–91.7
10th	%	5.6	8.8	12.0	73.7	85.6
(n=6,654)	CI	4.9–6.2	8.1–9.6	11.2–12.9	72.5–74.8	84.7–86.6
12th	%	9.1	11.8	13.6	65.4	79.0
(n=5,096)	CI	8.3–10.1	10.8–12.9	12.6–14.6	63.9–66.9	77.7–80.3

Note: The confidence interval (CI) is 95 percent.

Table TN.46. Parents Feel Smoking Tobacco is Wrong

		How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.9	4.1	9.8	83.2	93.0
(n=21,023)	CI	2.6–3.2	3.8–4.4	9.4–10.3	82.6–83.8	92.6–93.5
Male	%	3.5	4.5	10.2	81.9	92.0
(n=10,234)	CI	3.0–3.9	4.1–5.0	9.5–10.9	81.0–82.7	91.4–92.6
Female	%	1.9	3.5	9.2	85.3	94.6
(n=10,216)	CI	1.6–2.3	3.1–4.0	8.6–9.9	84.5–86.1	94.0–95.1
White	%	2.7	4.4	10.4	82.5	92.9
(n=16,197)	CI	2.4–3.1	4.0–4.8	9.9–11.0	81.8–83.2	92.4–93.4
Black	%	3.4	2.8	7.9	85.9	93.8
(n=1,358)	CI	2.5–4.6	1.9–4.0	6.4–9.6	83.7–87.8	92.2–95.1
Hispanic	%	2.3	2.1	6.0	89.5	95.6
(n=935)	CI	1.4–3.7	1.4–3.3	4.5–8.0	87.1–91.5	93.9–96.8
8th	%	1.3	2.0	7.1	89.6	96.7
(n=9,276)	CI	1.1–1.6	1.7–2.3	6.5–7.6	88.9–90.2	96.3–97.0
10th	%	2.5	3.7	9.2	84.7	93.9
(n=6,649)	CI	2.1–2.9	3.2–4.2	8.4–10.0	83.7–85.7	93.2–94.5
12th	%	5.1	6.9	13.8	74.2	88.0
(n=5,095)	CI	4.4–5.9	6.2–7.7	12.7–14.9	72.8–75.6	86.9–89.0

Note: The confidence interval (CI) is 95 percent.

Table TN.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.9	1.9	6.1	90.1	96.2
(n=21,015)	CI	1.7–2.1	1.7–2.2	5.7–6.5	89.6–90.6	95.8–96.5
Male	%	2.2	2.0	6.5	89.3	95.7
(n=10,220)	CI	1.9–2.6	1.7–2.4	5.9–7.1	88.5–90.0	95.2–96.2
Female	%	1.2	1.7	5.5	91.6	97.1
(n=10,224)	CI	1.0–1.4	1.4–2.0	5.0–6.1	90.9–92.2	96.7–97.5
White	%	1.6	2.0	6.3	90.1	96.4
(n=16,198)	CI	1.4–1.8	1.7–2.3	5.9–6.8	89.5–90.6	96.0–96.8
Black	%	3.3	1.3	5.3	90.1	95.4
(n=1,356)	CI	2.4–4.5	0.8–2.3	4.1–6.7	88.2–91.7	93.9–96.5
Hispanic	%	2.2	0.7	3.9	93.2	97.2
(n=932)	CI	1.3–3.5	0.3–1.5	2.8–5.5	91.3–94.8	95.7–98.1
8th	%	0.9	1.0	3.7	94.4	98.1
(n=9,272)	CI	0.8–1.1	0.8–1.2	3.3–4.2	93.9–94.8	97.8–98.4
10th	%	1.9	1.6	5.1	91.4	96.5
(n=6,648)	CI	1.6–2.3	1.3–2.0	4.6–5.7	90.6–92.1	95.9–97.0
12th	%	3.0	3.3	9.8	83.8	93.7
(n=5,095)	CI	2.6–3.6	2.7–4.0	8.9–10.8	82.6–85.0	92.8–94.4

Note: The confidence interval (CI) is 95 percent.

Table TN.48. Parents Feel Using Electronic Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.1	7.8	12.2	74.9	87.0
(n=21,005)	CI	4.8–5.5	7.4–8.3	11.7–12.7	74.2–75.5	86.5–87.6
Male	%	5.7	8.0	12.4	73.9	86.2
(n=10,218)	CI	5.2–6.3	7.4–8.7	11.7–13.1	72.9–74.8	85.4–87.0
Female	%	4.0	7.6	11.7	76.7	88.4
(n=10,215)	CI	3.6–4.5	7.0–8.2	11.0–12.4	75.8–77.6	87.7–89.1
White	%	5.0	8.2	12.8	74.0	86.8
(n=16,191)	CI	4.6–5.4	7.8–8.8	12.3–13.4	73.2–74.7	86.2–87.4
Black	%	5.4	4.6	8.5	81.5	90.0
(n=1,355)	CI	4.1–7.0	3.6–6.0	6.9–10.4	79.0–83.7	88.0–91.6
Hispanic	%	3.7	4.1	9.0	83.2	92.1
(n=932)	CI	2.6–5.3	2.8–6.0	7.1–11.2	80.2–85.7	89.9–93.9
8th	%	3.0	5.3	10.2	81.6	91.7
(n=9,266)	CI	2.6–3.3	4.9–5.8	9.5–10.8	80.7–82.4	91.1–92.3
10th	%	4.5	7.7	12.8	75.0	87.8
(n=6,643)	CI	4.0–5.1	7.0–8.5	11.9–13.7	73.8–76.1	86.8–88.6
12 th	%	8.3	10.9	13.8	67.1	80.9
(n=5,096)	CI	7.4–9.2	9.9–11.9	12.8–14.8	65.6–68.6	79.6–82.1

Note: The confidence interval (CI) is 95 percent.

Table TN.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.9	5.2	8.2	81.7	89.9
(n=21,012)	CI	4.6–5.3	4.8–5.5	7.8–8.7	81.1–82.3	89.4–90.4
Male	%	5.0	5.0	8.4	81.6	89.9
(n=10,221)	CI	4.6–5.6	4.5–5.5	7.8–9.0	80.7–82.4	89.2–90.6
Female	%	4.2	5.2	7.9	82.8	90.7
(n=10,218)	CI	3.7–4.7	4.7–5.7	7.3–8.5	81.9–83.6	90.0–91.3
White	%	4.3	5.0	8.3	82.3	90.6
(n=16,192)	CI	4.0–4.7	4.7–5.5	7.8–8.8	81.6–83.0	90.1–91.2
Black	%	7.9	5.2	8.7	78.2	86.9
(n=1,353)	CI	6.4–9.7	4.0–6.7	7.0–10.7	75.6–80.7	84.7–88.8
Hispanic	%	3.1	3.0	6.5	87.4	93.9
(n=935)	CI	2.1–4.6	1.9–4.7	4.9–8.6	84.7–89.6	91.8–95.4
8 th	%	2.4	2.6	5.5	89.5	94.9
(n=9,274)	CI	2.1–2.8	2.3–3.0	5.0–6.0	88.8–90.1	94.4–95.4
10 th	%	5.2	6.1	8.4	80.3	88.7
(n=6,645)	CI	4.6–5.9	5.5–6.7	7.7–9.2	79.2–81.4	87.9–89.6
12 th	%	7.4	7.0	11.1	74.4	85.6
(n=5,093)	CI	6.7–8.2	6.3–7.9	10.2–12.2	73.0–75.8	84.4–86.6

Note: The confidence interval (CI) is 95 percent.

Table TN.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.5	3.7	7.1	84.7	91.8
(n=20,982)	CI	4.2–4.9	3.4–4.0	6.7–7.5	84.1–85.2	91.3–92.2
Male	%	4.6	3.7	7.3	84.4	91.7
(n=10,213)	CI	4.1–5.1	3.3–4.2	6.7–7.8	83.6–85.2	91.0–92.3
Female	%	4.0	3.5	6.7	85.8	92.5
(n=10,199)	CI	3.5–4.4	3.2–4.0	6.1–7.3	85.0–86.6	91.9–93.1
White	%	3.9	3.6	7.1	85.4	92.5
(n=16,174)	CI	3.6–4.3	3.3–3.9	6.6–7.5	84.8–86.1	92.0–93.0
Black	%	8.0	4.1	7.5	80.4	87.9
(n=1,355)	CI	6.5–9.8	3.1–5.4	5.9–9.4	77.8–82.8	85.8–89.7
Hispanic	%	3.3	2.0	4.9	89.8	94.7
(n=931)	CI	2.2–4.8	1.2–3.3	3.5–6.9	87.3–91.8	92.9–96.1
8 th	%	2.2	1.8	4.7	91.3	96.1
(n=9,260)	CI	1.9–2.5	1.5–2.1	4.3–5.2	90.7–91.9	95.6–96.5
10 th	%	4.8	4.2	7.5	83.6	91.0
(n=6,638)	CI	4.2–5.4	3.7–4.7	6.8–8.2	82.6–84.6	90.2–91.8
12 th	%	7.0	5.3	9.3	78.4	87.7
(n=5,084)	CI	6.3–7.8	4.7–6.1	8.4–10.2	77.1–79.6	86.7–88.7

Note: The confidence interval (CI) is 95 percent.

Table TN.51. Friends Feel Misusing Prescription Drugs is Wrong

How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.8	6.9	14.9	70.4	85.4
(n=20,962)	CI	7.4–8.2	6.5–7.3	14.4–15.5	69.7–71.1	84.8–85.9
Male	%	8.5	7.1	15.5	68.9	84.4
(n=10,186)	CI	7.9–9.2	6.5–7.6	14.8–16.4	67.9–69.9	83.6–85.2
Female	%	6.4	6.6	14.5	72.6	87.0
(n=10,205)	CI	5.9–6.9	6.1–7.2	13.7–15.3	71.6–73.5	86.3–87.7
White	%	7.2	6.8	15.2	70.9	86.0
(n=16,159)	CI	6.7–7.6	6.4–7.2	14.6–15.8	70.1–71.7	85.4–86.6
Black	%	10.0	6.8	13.1	70.2	83.2
(n=1,351)	CI	8.4–11.9	5.4–8.4	11.3–15.1	67.5–72.8	80.9–85.3
Hispanic	%	7.7	6.4	15.5	70.4	85.9
(n=931)	CI	6.0–9.9	4.9–8.3	13.0–18.3	67.0–73.6	83.2–88.2
8th	%	5.6	5.5	13.5	75.5	88.9
(n=9,250)	CI	5.1–6.1	5.0–6.0	12.7–14.2	74.5–76.4	88.3–89.6
10th	%	8.8	7.5	15.8	67.9	83.7
(n=6,626)	CI	8.1–9.6	6.8–8.2	14.8–16.7	66.7–69.2	82.7–84.7
12th	%	9.1	7.7	15.7	67.5	83.2
(n=5,086)	CI	8.2–10.0	7.0–8.5	14.6–16.9	66.0–68.9	82.1–84.3

Note: The confidence interval (CI) is 95 percent.

Table TN.52. Friends Feel Drinking Alcohol is Wrong

How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	15.5	15.1	18.4	51.0	69.4
(n=20,992)	CI	15.0–16.1	14.5–15.6	17.8–19.0	50.2–51.7	68.7–70.1
Male	%	17.1	14.8	18.5	49.6	68.1
(n=10,207)	CI	16.3–18.0	14.0–15.6	17.7–19.4	48.5–50.7	67.1–69.2
Female	%	13.4	15.4	18.5	52.7	71.2
(n=10,213)	CI	12.6–14.3	14.6–16.2	17.7–19.4	51.6–53.8	70.2–72.2
White	%	15.3	15.4	19.0	50.4	69.3
(n=16,180)	CI	14.6–16.0	14.8–16.0	18.3–19.7	49.5–51.2	68.5–70.1
Black	%	15.6	11.2	16.2	57.0	73.2
(n=1,354)	CI	13.6–17.9	9.4–13.2	14.2–18.4	54.1–59.9	70.5–75.7
Hispanic	%	13.5	14.3	18.4	53.8	72.2
(n=930)	CI	11.1–16.2	12.0–17.1	15.8–21.3	50.1–57.4	68.8–75.3
8th	%	8.2	11.1	18.2	62.4	80.6
(n=9,263)	CI	7.7–8.8	10.5–11.8	17.4–19.1	61.4–63.5	79.8–81.5
10th	%	17.1	17.0	19.5	46.4	65.9
(n=6,635)	CI	16.1–18.1	16.0–18.0	18.5–20.6	45.1–47.8	64.6–67.2
12th	%	22.1	17.4	17.4	43.0	60.4
(n=5,094)	CI	20.8–23.5	16.3–18.7	16.3–18.6	41.5–44.6	58.9–62.0

Note: The confidence interval (CI) is 95 percent.

Table TN.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.8	12.2	18.2	54.8	73.0
(n=20,965)	CI	14.2–15.3	11.7–12.8	17.6–18.8	54.1–55.6	72.3–73.7
Male	%	16.7	12.2	18.4	52.7	71.1
(n=10,190)	CI	15.9–17.6	11.5–12.9	17.5–19.2	51.6–53.8	70.1–72.1
Female	%	12.2	12.3	18.3	57.2	75.5
(n=10,204)	CI	11.5–13.0	11.6–13.1	17.4–19.2	56.1–58.3	74.5–76.4
White	%	15.0	12.8	18.8	53.5	72.2
(n=16,163)	CI	14.3–15.7	12.2–13.4	18.1–19.5	52.6–54.4	71.4–73.0
Black	%	13.4	7.6	14.3	64.6	79.0
(n=1,352)	CI	11.5–15.6	6.2–9.3	12.4–16.5	61.8–67.4	76.4–81.3
Hispanic	%	10.1	10.2	19.8	59.9	79.7
(n=926)	CI	8.1–12.5	8.1–12.6	17.1–22.9	56.3–63.4	76.7–82.5
8th	%	8.0	9.0	16.9	66.1	83.0
(n=9,253)	CI	7.4–8.6	8.4–9.6	16.1–17.7	65.1–67.1	82.2–83.8
10th	%	15.3	13.2	19.3	52.1	71.4
(n=6,625)	CI	14.4–16.3	12.3–14.2	18.3–20.4	50.7–53.5	70.2–72.7
12th	%	21.8	14.8	18.2	45.1	63.3
(n=5,087)	CI	20.6–23.2	13.8–16.0	17.0–19.5	43.6–46.7	61.8–64.8

Note: The confidence interval (CI) is 95 percent.

Table TN.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	26.5	14.6	13.8	45.0	58.9
(n=20,918)	CI	25.8–27.2	14.1–15.1	13.3–14.4	44.3–45.8	58.1–59.6
Male	%	27.7	14.0	14.0	44.3	58.3
(n=10,163)	CI	26.7–28.7	13.3–14.8	13.2–14.7	43.2–45.4	57.2–59.4
Female	%	24.8	15.2	14.0	46.0	60.0
(n=10,185)	CI	23.8–25.8	14.4–16.0	13.2–14.8	44.9–47.1	58.9–61.1
White	%	27.4	15.0	14.0	43.5	57.5
(n=16,135)	CI	26.6–28.2	14.4–15.7	13.4–14.6	42.7–44.4	56.7–58.4
Black	%	19.5	8.9	13.6	58.0	71.6
(n=1,345)	CI	17.3–22.0	7.4–10.6	11.6–15.8	55.0–60.9	68.9–74.2
Hispanic	%	18.7	12.8	16.1	52.4	68.5
(n=921)	CI	15.9–21.8	10.5–15.5	13.7–18.9	48.8–56.1	65.0–71.9
8th	%	15.7	13.0	15.0	56.3	71.3
(n=9,238)	CI	15.0–16.5	12.3–13.7	14.2–15.8	55.2–57.4	70.3–72.3
10th	%	29.0	16.3	13.9	40.9	54.8
(n=6,604)	CI	27.7–30.2	15.3–17.3	13.0–14.9	39.5–42.2	53.4–56.1
12th	%	36.1	14.5	12.4	37.0	49.4
(n=5,076)	CI	34.6–37.6	13.5–15.6	11.4–13.5	35.5–38.5	47.8–50.9

Note: The confidence interval (CI) is 95 percent.

Table TN.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	21.4	12.3	14.1	52.2	66.3
(n=20,955)	CI	20.7–22.0	11.8–12.8	13.6–14.7	51.4–53.0	65.6–67.1
Male	%	22.0	11.9	14.7	51.4	66.1
(n=10,189)	CI	21.1–22.9	11.2–12.7	14.0–15.6	50.2–52.5	65.0–67.2
Female	%	19.9	12.8	13.8	53.5	67.3
(n=10,197)	CI	19.0–20.8	12.0–13.5	13.0–14.6	52.4–54.7	66.2–68.4
White	%	20.5	12.3	14.6	52.6	67.2
(n=16,155)	CI	19.8–21.2	11.7–12.9	14.0–15.2	51.7–53.5	66.4–68.0
Black	%	26.7	10.1	11.7	51.5	63.2
(n=1,354)	CI	24.1–29.4	8.5–11.9	9.8–13.8	48.6–54.5	60.3–66.0
Hispanic	%	15.9	12.1	15.9	56.1	72.0
(n=929)	CI	13.4–18.8	9.8–14.8	13.5–18.7	52.4–59.6	68.5–75.2
8th	%	10.7	8.0	13.8	67.4	81.2
(n=9,245)	CI	10.1–11.4	7.5–8.7	13.1–14.6	66.4–68.4	80.4–82.1
10th	%	24.8	13.9	15.0	46.3	61.3
(n=6,623)	CI	23.6–26.0	12.9–14.8	14.1–16.0	44.9–47.7	60.0–62.7
12th	%	29.6	15.4	13.5	41.5	55.0
(n=5,087)	CI	28.2–31.1	14.3–16.5	12.4–14.6	40.0–43.1	53.5–56.6

Note: The confidence interval (CI) is 95 percent.

Table TN.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	88.2	5.4	2.4	0.9	3.2	11.8
(n=20,966)	CI	87.7–88.6	5.1–5.7	2.2–2.6	0.8–1.1	2.9–3.5	11.4–12.3
Male	%	89.5	4.5	2.2	0.8	3.0	10.5
(n=10,201)	CI	88.7–90.1	4.1–5.0	1.9–2.6	0.6–1.0	2.6–3.4	9.9–11.3
Female	%	87.5	6.3	2.5	1.0	2.8	12.5
(n=10,197)	CI	86.8–88.2	5.8–6.8	2.2–2.8	0.8–1.3	2.4–3.2	11.8–13.2
White	%	88.4	5.5	2.4	0.9	2.8	11.6
(n=16,167)	CI	87.8–89.0	5.1–5.9	2.1–2.6	0.7–1.1	2.5–3.2	11.0–12.2
Black	%	90.0	3.5	1.6	1.3	3.6	10.0
(n=1,360)	CI	88.1–91.5	2.6–4.7	1.1–2.5	0.7–2.2	2.7–4.8	8.5–11.9
Hispanic	%	90.3	4.6	2.2	0.8	2.2	9.7
(n=926)	CI	87.9–92.3	3.2–6.4	1.4–3.5	0.4–1.7	1.3–3.5	7.7–12.1
8th	%	91.8	4.3	1.6	0.6	1.7	8.2
(n=9,231)	CI	91.2–92.4	3.8–4.7	1.4–2.0	0.5–0.8	1.5–2.0	7.6–8.8
10th	%	86.0	6.3	2.8	1.2	3.8	14.0
(n=6,628)	CI	85.0–86.9	5.7–7.0	2.4–3.2	0.9–1.5	3.3–4.4	13.1–15.0
12th	%	86.5	5.6	2.7	1.0	4.2	13.5
(n=5,107)	CI	85.5–87.5	5.0–6.3	2.3–3.2	0.7–1.4	3.6–4.8	12.5–14.5

Note: The confidence interval (CI) is 95 percent.

Table TN.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	54.6	12.8	10.2	5.0	17.3	45.4
(n=21,105)	CI	53.8–55.4	12.3–13.3	9.8–10.7	4.7–5.4	16.7–18.0	44.6–46.2
Male	%	59.1	10.9	8.7	4.1	17.2	40.9
(n=10,262)	CI	58.0–60.2	10.2–11.6	8.1–9.3	3.7–4.6	16.3–18.1	39.8–42.0
Female	%	50.5	14.8	11.8	6.0	16.9	49.5
(n=10,266)	CI	49.4–51.6	14.0–15.6	11.1–12.5	5.5–6.6	16.1–17.8	48.4–50.6
White	%	53.3	12.9	10.6	5.4	17.8	46.7
(n=16,267)	CI	52.4–54.2	12.4–13.5	10.1–11.1	5.0–5.8	17.1–18.5	45.8–47.6
Black	%	66.6	9.2	7.9	3.0	13.3	33.4
(n=1,371)	CI	63.8–69.4	7.7–11.0	6.4–9.6	2.2–4.1	11.3–15.6	30.6–36.2
Hispanic	%	62.5	14.8	7.7	2.9	12.2	37.5
(n=935)	CI	58.9–65.9	12.3–17.7	6.0–9.8	1.9–4.2	10.0–14.7	34.1–41.1
8th	%	70.2	12.1	7.5	2.8	7.4	29.8
(n=9,286)	CI	69.2–71.2	11.4–12.8	7.0–8.1	2.5–3.2	6.9–8.0	28.8–30.8
10th	%	48.8	14.6	11.7	5.8	19.1	51.2
(n=6,681)	CI	47.4–50.2	13.7–15.6	10.8–12.5	5.2–6.5	18.0–20.2	49.8–52.6
12th	%	43.5	11.5	11.6	6.7	26.6	56.5
(n=5,138)	CI	42.0–45.1	10.5–12.6	10.7–12.6	5.9–7.6	25.2–28.0	54.9–58.0

Note: The confidence interval (CI) is 95 percent.

Table TN.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	69.0	9.4	6.3	3.7	11.6	31.0
(n=21,049)	CI	68.3–69.7	9.0–9.9	5.9–6.7	3.4–4.0	11.1–12.2	30.3–31.7
Male	%	70.9	8.4	5.5	3.3	11.9	29.1
(n=10,233)	CI	69.9–71.9	7.8–9.0	5.0–6.1	2.9–3.7	11.1–12.7	28.1–30.1
Female	%	67.6	10.4	7.0	4.2	10.8	32.4
(n=10,243)	CI	66.5–68.7	9.8–11.1	6.5–7.6	3.7–4.7	10.1–11.6	31.3–33.5
White	%	67.5	9.9	6.7	3.9	12.0	32.5
(n=16,224)	CI	66.7–68.4	9.4–10.4	6.3–7.2	3.5–4.3	11.4–12.6	31.6–33.3
Black	%	79.6	5.8	3.8	2.4	8.5	20.4
(n=1,369)	CI	77.0–81.9	4.5–7.3	2.8–5.0	1.7–3.5	6.8–10.5	18.1–23.0
Hispanic	%	78.3	8.4	3.8	2.3	7.2	21.7
(n=931)	CI	75.2–81.2	6.5–10.7	2.7–5.4	1.5–3.4	5.4–9.4	18.8–24.8
8th	%	86.0	6.4	3.0	1.2	3.4	14.0
(n=9,262)	CI	85.3–86.7	5.9–6.9	2.6–3.4	1.0–1.5	3.0–3.8	13.3–14.7
10th	%	64.4	11.3	7.4	4.6	12.4	35.6
(n=6,653)	CI	63.0–65.6	10.5–12.1	6.8–8.2	4.0–5.2	11.5–13.3	34.4–37.0
12th	%	55.0	10.7	8.6	5.5	20.1	45.0
(n=5,134)	CI	53.4–56.5	9.8–11.7	7.8–9.6	4.9–6.3	18.8–21.4	43.5–46.6

Note: The confidence interval (CI) is 95 percent.

Table TN.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have smoked part, or all of a cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	71.7	10.3	5.8	3.0	9.1	28.3
(n=21,000)	CI	71.0–72.4	9.9–10.8	5.4–6.2	2.8–3.3	8.6–9.6	27.6–29.0
Male	%	73.6	9.6	5.0	2.6	9.2	26.4
(n=10,223)	CI	72.6–74.6	9.0–10.2	4.6–5.5	2.2–2.9	8.6–10.0	25.4–27.4
Female	%	70.5	11.2	6.5	3.5	8.3	29.5
(n=10,208)	CI	69.4–71.5	10.5–12.0	6.0–7.1	3.1–4.0	7.7–8.9	28.5–30.6
White	%	70.2	11.0	6.3	3.2	9.2	29.8
(n=16,181)	CI	69.4–71.0	10.5–11.6	5.9–6.8	2.9–3.6	8.7–9.8	29.0–30.6
Black	%	83.4	6.0	2.6	1.6	6.3	16.6
(n=1,366)	CI	81.1–85.5	4.9–7.4	1.8–3.7	1.0–2.6	4.9–8.0	14.5–18.9
Hispanic	%	81.1	7.0	3.6	1.7	6.5	18.9
(n=931)	CI	78.2–83.7	5.4–9.1	2.5–5.2	1.0–2.9	5.0–8.4	16.3–21.8
8th	%	80.7	8.9	3.7	1.8	4.9	19.3
(n=9,257)	CI	79.8–81.5	8.3–9.6	3.3–4.1	1.5–2.1	4.4–5.4	18.5–20.2
10th	%	69.3	11.4	6.1	3.2	10.0	30.7
(n=6,622)	CI	68.0–70.5	10.5–12.3	5.5–6.8	2.8–3.7	9.2–10.9	29.5–32.0
12th	%	64.4	10.8	7.8	4.3	12.8	35.6
(n=5,121)	CI	62.8–65.8	9.9–11.8	7.0–8.7	3.7–5.0	11.8–13.9	34.2–37.2

Note: The confidence interval (CI) is 95 percent.

Table TN.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used an e-cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	58.3	10.0	7.2	4.8	19.7	41.7
(n=20,995)	CI	57.6–59.1	9.5–10.4	6.8–7.6	4.4–5.1	19.1–20.3	40.9–42.4
Male	%	59.3	9.1	6.3	4.6	20.7	40.7
(n=10,212)	CI	58.2–60.4	8.5–9.7	5.8–6.9	4.1–5.1	19.8–21.7	39.6–41.8
Female	%	57.7	11.0	8.2	5.0	18.1	42.3
(n=10,209)	CI	56.6–58.8	10.4–11.7	7.6–8.8	4.6–5.5	17.2–19.0	41.2–43.4
White	%	55.7	10.6	7.8	5.2	20.6	44.3
(n=16,179)	CI	54.9–56.6	10.1–11.2	7.4–8.3	4.8–5.6	19.9–21.4	43.4–45.1
Black	%	78.4	5.3	3.5	2.0	10.8	21.6
(n=1,367)	CI	75.9–80.7	4.2–6.7	2.6–4.7	1.3–2.9	9.1–12.8	19.3–24.1
Hispanic	%	71.2	7.7	5.3	3.0	12.7	28.8
(n=926)	CI	67.9–74.4	6.1–9.8	3.9–7.3	2.0–4.3	10.5–15.3	25.6–32.1
8th	%	69.6	10.1	5.8	3.0	11.5	30.4
(n=9,246)	CI	68.6–70.6	9.5–10.7	5.3–6.3	2.6–3.4	10.8–12.2	29.4–31.4
10th	%	53.3	10.8	8.2	5.4	22.3	46.7
(n=6,636)	CI	51.9–54.7	10.0–11.6	7.5–9.0	4.9–6.1	21.2–23.5	45.3–48.1
12th	%	51.2	8.9	7.8	6.0	26.1	48.8
(n=5,113)	CI	49.6–52.7	8.1–9.8	7.0–8.6	5.3–6.8	24.7–27.5	47.3–50.4

Note: The confidence interval (CI) is 95 percent.

Table TN.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	66.9	9.5	6.4	3.7	13.5	33.1
(n=21,065)	CI	66.1–67.6	9.0–10.0	6.0–6.8	3.4–4.0	12.9–14.1	32.4–33.9
Male	%	69.3	8.3	5.7	3.6	13.1	30.7
(n=10,240)	CI	68.3–70.4	7.7–8.9	5.2–6.2	3.2–4.0	12.3–13.9	29.6–31.7
Female	%	65.1	10.7	7.1	3.9	13.2	34.9
(n=10,249)	CI	64.0–66.1	10.0–11.4	6.6–7.8	3.5–4.3	12.4–14.0	33.9–36.0
White	%	67.2	9.8	6.5	3.8	12.7	32.8
(n=16,237)	CI	66.4–68.1	9.3–10.4	6.0–6.9	3.5–4.1	12.1–13.3	31.9–33.6
Black	%	65.3	6.9	6.4	3.5	17.9	34.7
(n=1,366)	CI	62.4–68.1	5.6–8.5	5.1–8.0	2.6–4.7	15.6–20.4	31.9–37.6
Hispanic	%	76.0	6.4	4.4	2.9	10.3	24.0
(n=935)	CI	72.8–78.9	4.9–8.3	3.1–6.2	1.9–4.4	8.3–12.8	21.1–27.2
8th	%	81.8	7.5	3.6	2.0	5.1	18.2
(n=9,286)	CI	80.9–82.6	6.9–8.1	3.2–4.0	1.7–2.3	4.7–5.6	17.4–19.1
10th	%	61.3	10.8	7.1	4.7	16.2	38.7
(n=6,659)	CI	59.9–62.6	10.0–11.6	6.4–7.8	4.2–5.2	15.2–17.2	37.4–40.1
12th	%	56.3	10.3	8.8	4.6	20.0	43.7
(n=5,120)	CI	54.8–57.9	9.4–11.3	7.9–9.7	4.0–5.3	18.7–21.3	42.1–45.2

Note: The confidence interval (CI) is 95 percent.

Table TN.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	9.5	5.6	13.0	71.9	84.9
(n=20,785)	CI	9.1–10.0	5.3–5.9	12.5–13.6	71.2–72.5	84.3–85.4
Male	%	11.5	5.2	13.6	69.8	83.4
(n=10,110)	CI	10.8–12.2	4.7–5.6	12.9–14.4	68.8–70.8	82.5–84.2
Female	%	7.0	6.0	12.5	74.5	87.0
(n=10,107)	CI	6.5–7.6	5.5–6.5	11.8–13.2	73.5–75.4	86.2–87.7
White	%	7.7	5.4	13.0	74.0	87.0
(n=16,032)	CI	7.2–8.1	5.0–5.8	12.4–13.6	73.2–74.8	86.4–87.5
Black	%	25.3	6.5	11.5	56.7	68.2
(n=1,346)	CI	22.8–28.0	5.2–8.0	9.8–13.5	53.8–59.6	65.4–70.9
Hispanic	%	14.3	7.0	14.1	64.6	78.7
(n=913)	CI	11.8–17.1	5.4–9.0	11.8–16.7	61.0–68.0	75.5–81.6
8th	%	8.3	6.5	13.1	72.1	85.2
(n=9,162)	CI	7.7–8.9	6.0–7.0	12.4–13.8	71.1–73.1	84.4–86.0
10th	%	10.0	5.9	13.0	71.1	84.1
(n=6,569)	CI	9.3–10.9	5.3–6.6	12.1–13.9	69.8–72.3	83.0–85.0
12th	%	10.3	4.2	13.0	72.5	85.5
(n=5,054)	CI	9.4–11.2	3.7–4.9	12.0–14.1	71.1–73.8	84.4–86.5

Note: The confidence interval (CI) is 95 percent.

Table TN.63. Perceived Risk of Regularly Drinking Alcohol

How much do you think people risk harming themselves physically and in other ways when they take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	13.3	15.6	23.8	47.3	71.1
(n=20,871)	CI	12.8–13.8	15.1–16.2	23.1–24.5	46.5–48.0	70.3–71.8
Male	%	16.1	16.6	23.9	43.3	67.3
(n=10,149)	CI	15.3–16.9	15.8–17.5	23.0–24.9	42.2–44.4	66.2–68.3
Female	%	10.0	14.5	23.9	51.6	75.4
(n=10,150)	CI	9.4–10.7	13.7–15.4	22.9–24.8	50.5–52.7	74.5–76.4
White	%	11.8	16.2	24.6	47.4	72.0
(n=16,106)	CI	11.3–12.4	15.6–16.9	23.8–25.3	46.5–48.3	71.2–72.8
Black	%	26.2	13.2	18.8	41.7	60.6
(n=1,344)	CI	23.7–29.0	11.3–15.3	16.7–21.2	38.9–44.7	57.7–63.4
Hispanic	%	17.3	12.0	19.8	50.9	70.7
(n=916)	CI	14.6–20.3	10.0–14.4	17.1–22.9	47.2–54.6	67.3–73.9
8th	%	11.6	14.3	22.0	52.1	74.1
(n=9,190)	CI	10.9–12.3	13.6–15.1	21.1–22.9	51.0–53.1	73.1–75.0
10th	%	14.0	15.8	24.4	45.8	70.2
(n=6,608)	CI	13.1–14.9	14.9–16.8	23.3–25.6	44.4–47.2	68.9–71.4
12th	%	14.5	16.9	25.1	43.5	68.6
(n=5,073)	CI	13.5–15.6	15.7–18.2	23.7–26.4	42.0–45.1	67.1–70.0

Note: The confidence interval (CI) is 95 percent.

Table TN.64. Perceived Risk of Binge Drinking

How much do you think people risk harming themselves physically and in other ways when they take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	11.4	10.3	21.4	56.9	78.3
(n=20,884)	CI	10.9–11.9	9.9–10.8	20.7–22.0	56.1–57.7	77.6–78.9
Male	%	13.9	11.1	21.9	53.1	75.0
(n=10,158)	CI	13.1–14.7	10.4–11.8	21.0–22.8	52.0–54.2	74.1–76.0
Female	%	8.3	9.6	20.8	61.3	82.0
(n=10,155)	CI	7.8–9.0	9.0–10.3	19.9–21.7	60.2–62.3	81.2–82.9
White	%	9.7	10.6	22.0	57.7	79.7
(n=16,121)	CI	9.2–10.2	10.1–11.2	21.3–22.8	56.8–58.5	78.9–80.4
Black	%	26.0	9.2	16.3	48.5	64.8
(n=1,341)	CI	23.5–28.7	7.7–11.1	14.2–18.6	45.5–51.4	61.9–67.6
Hispanic	%	15.9	9.0	17.2	57.8	75.0
(n=918)	CI	13.4–18.9	7.1–11.4	14.7–19.9	54.2–61.4	71.7–78.1
8th	%	9.8	8.6	19.3	62.3	81.7
(n=9,194)	CI	9.1–10.4	8.0–9.2	18.5–20.2	61.3–63.4	80.8–82.5
10th	%	11.7	10.7	22.0	55.6	77.6
(n=6,607)	CI	10.8–12.6	9.9–11.6	20.9–23.2	54.2–56.9	76.5–78.7
12th	%	12.9	11.9	23.0	52.2	75.2
(n=5,083)	CI	11.9–13.9	11.0–13.0	21.7–24.3	50.7–53.8	73.9–76.5

Note: The confidence interval (CI) is 95 percent.

Table TN.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.0	6.0	16.0	68.0	84.0
(n=20,923)	CI	9.6–10.5	5.6–6.3	15.5–16.6	67.2–68.7	83.4–84.6
Male	%	12.0	6.1	17.2	64.6	81.8
(n=10,173)	CI	11.3–12.8	5.7–6.7	16.4–18.1	63.5–65.7	81.0–82.7
Female	%	7.5	5.8	14.8	71.9	86.7
(n=10,175)	CI	6.9–8.1	5.3–6.3	14.1–15.6	70.9–72.8	86.0–87.4
White	%	8.3	5.7	16.3	69.7	86.0
(n=16,147)	CI	7.8–8.8	5.4–6.1	15.6–16.9	68.9–70.5	85.4–86.6
Black	%	25.1	7.2	12.7	55.0	67.7
(n=1,349)	CI	22.6–27.8	5.8–8.9	10.9–14.8	52.0–57.9	64.8–70.4
Hispanic	%	15.3	6.8	17.5	60.4	77.9
(n=919)	CI	12.8–18.2	5.3–8.6	15.0–20.4	56.7–63.9	74.7–80.8
8th	%	8.7	6.4	13.6	71.4	84.9
(n=9,225)	CI	7.9–9.6	5.8–7.1	12.7–14.5	70.1–72.6	83.9–85.9
10th	%	10.7	5.9	13.5	69.8	83.4
(n=6,617)	CI	9.5–12.0	5.2–6.8	12.3–14.9	67.9–71.7	81.8–84.8
12th	%	10.8	4.7	13.4	71.1	84.5
(n=5,081)	CI	9.4–12.5	3.7–5.9	11.9–15.1	68.9–73.2	82.6–86.2

Note: The confidence interval (CI) is 95 percent.

Table TN.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	13.6	20.4	26.3	39.6	66.0
(n=20,879)	CI	13.1–14.2	19.8–21.0	25.7–27.0	38.9–40.4	65.2–66.7
Male	%	16.3	22.2	24.9	36.6	61.5
(n=10,154)	CI	15.5–17.2	21.3–23.1	24.0–25.9	35.5–37.6	60.4–62.6
Female	%	10.4	18.4	27.9	43.3	71.2
(n=10,154)	CI	9.7–11.1	17.6–19.3	26.9–28.9	42.2–44.4	70.2–72.2
White	%	12.1	20.9	27.3	39.7	67.0
(n=16,123)	CI	11.5–12.6	20.2–21.7	26.5–28.1	38.9–40.6	66.2–67.8
Black	%	28.3	17.2	18.9	35.6	54.5
(n=1,344)	CI	25.7–31.1	15.2–19.5	16.7–21.3	32.9–38.5	51.6–57.4
Hispanic	%	17.1	16.2	24.7	42.0	66.7
(n=914)	CI	14.4–20.1	13.8–19.0	21.7–28.0	38.4–45.7	63.2–70.1
8th	%	12.0	18.3	24.1	45.6	69.7
(n=9,208)	CI	11.3–12.7	17.5–19.2	23.2–25.1	44.5–46.6	68.7–70.6
10th	%	13.6	21.8	27.0	37.6	64.6
(n=6,595)	CI	12.7–14.6	20.7–23.0	25.8–28.2	36.3–38.9	63.3–65.9
12th	%	15.6	21.1	28.1	35.2	63.3
(n=5,076)	CI	14.5–16.7	19.9–22.4	26.7–29.5	33.7–36.8	61.8–64.8

Note: The confidence interval (CI) is 95 percent.

Table TN.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	29.8	21.2	15.1	33.9	48.9
(n=20,791)	CI	29.1–30.6	20.6–21.9	14.5–15.7	33.1–34.6	48.2–49.7
Male	%	32.6	20.6	14.6	32.3	46.8
(n=10,108)	CI	31.5–33.7	19.7–21.5	13.8–15.4	31.2–33.3	45.7–48.0
Female	%	26.2	22.1	15.8	35.9	51.7
(n=10,113)	CI	25.2–27.2	21.2–23.0	15.1–16.7	34.8–36.9	50.6–52.8
White	%	28.7	21.5	15.0	34.8	49.8
(n=16,045)	CI	27.9–29.5	20.8–22.2	14.4–15.6	34.0–35.7	48.9–50.7
Black	%	40.6	19.6	13.4	26.4	39.8
(n=1,344)	CI	37.8–43.6	17.4–22.0	11.5–15.4	23.9–29.1	37.0–42.7
Hispanic	%	26.6	16.6	18.3	38.5	56.8
(n=910)	CI	23.4–30.0	14.1–19.4	15.6–21.3	35.0–42.2	53.2–60.5
8th	%	19.6	18.9	16.6	44.9	61.5
(n=9,147)	CI	18.7–20.4	18.1–19.8	15.8–17.4	43.8–46.0	60.5–62.6
10th	%	32.1	22.7	15.3	29.9	45.2
(n=6,576)	CI	30.8–33.4	21.6–23.9	14.3–16.3	28.6–31.2	43.8–46.6
12th	%	38.9	22.2	13.1	25.8	38.9
(n=5,068)	CI	37.4–40.5	20.9–23.5	12.1–14.2	24.4–27.2	37.4–40.4

Note: The confidence interval (CI) is 95 percent.

Table TN.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	22.2	15.8	19.2	42.8	62.0
(n=20,817)	CI	21.6–22.9	15.2–16.4	18.6–19.9	42.0–43.5	61.3–62.8
Male	%	25.0	15.9	18.9	40.3	59.1
(n=10,125)	CI	24.0–25.9	15.1–16.7	18.0–19.7	39.2–41.4	58.0–60.2
Female	%	18.8	15.5	19.8	45.9	65.8
(n=10,124)	CI	17.9–19.7	14.7–16.3	19.0–20.7	44.8–47.0	64.7–66.8
White	%	20.3	15.7	19.6	44.4	64.0
(n=16,079)	CI	19.6–21.1	15.0–16.3	18.9–20.3	43.5–45.3	63.1–64.8
Black	%	37.3	16.2	16.4	30.1	46.5
(n=1,338)	CI	34.4–40.2	14.1–18.6	14.4–18.6	27.5–32.9	43.6–49.5
Hispanic	%	22.8	13.7	21.3	42.2	63.5
(n=910)	CI	19.8–26.2	11.3–16.4	18.4–24.5	38.6–45.9	59.9–67.0
8th	%	15.2	11.2	18.0	55.5	73.6
(n=9,175)	CI	14.4–16.0	10.6–11.9	17.2–18.9	54.4–56.6	72.6–74.5
10th	%	23.8	17.2	20.1	38.9	59.0
(n=6,581)	CI	22.7–25.0	16.2–18.3	19.0–21.2	37.6–40.3	57.6–60.3
12th	%	28.3	19.4	19.7	32.6	52.3
(n=5,061)	CI	26.9–29.8	18.2–20.6	18.4–20.9	31.2–34.1	50.7–53.8

Note: The confidence interval (CI) is 95 percent.

Table TN.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.3	4.4	11.3	81.0	92.3
(n=20,914)	CI	3.0–3.5	4.1–4.7	10.8–11.8	80.4–81.6	91.9–92.7
Male	%	3.6	4.3	11.1	81.0	92.1
(n=10,175)	CI	3.2–4.1	3.9–4.7	10.4–11.8	80.1–81.9	91.5–92.7
Female	%	2.3	4.4	11.4	81.8	93.3
(n=10,168)	CI	2.0–2.6	4.0–4.9	10.7–12.2	80.9–82.6	92.7–93.8
White	%	2.6	4.2	11.1	82.1	93.2
(n=16,152)	CI	2.3–2.9	3.8–4.5	10.6–11.7	81.4–82.8	92.8–93.6
Black	%	6.3	5.1	11.1	77.5	88.6
(n=1,344)	CI	5.0–8.0	3.9–6.6	9.3–13.1	74.9–79.9	86.5–90.4
Hispanic	%	3.7	5.2	11.8	79.3	91.1
(n=910)	CI	2.5–5.3	3.8–7.1	9.7–14.2	76.2–82.1	88.8–92.9
8th	%	2.1	3.4	9.2	85.4	94.6
(n=9,219)	CI	1.8–2.4	3.0–3.8	8.6–9.8	84.6–86.1	94.1–95.0
10th	%	3.5	5.0	12.1	79.3	91.4
(n=6,621)	CI	3.1–4.0	4.5–5.6	11.2–13.0	78.2–80.4	90.7–92.1
12th	%	4.3	4.9	12.8	78.1	90.8
(n=5,074)	CI	3.8–4.9	4.3–5.6	11.8–13.8	76.8–79.3	89.9–91.6

Note: The confidence interval (CI) is 95 percent.

Table TN.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.0	10.6	19.0	64.5	83.4
(n=20,988)	CI	5.6–6.4	10.1–11.1	18.3–19.6	63.7–65.2	82.8–84.0
Male	%	6.9	10.7	18.1	64.3	82.4
(n=10,209)	CI	6.4–7.5	10.0–11.4	17.3–19.0	63.2–65.3	81.5–83.2
Female	%	4.3	10.5	19.9	65.3	85.2
(n=10,204)	CI	3.9–4.8	9.8–11.2	19.0–20.8	64.2–66.4	84.4–86.0
White	%	5.6	10.8	19.1	64.5	83.6
(n=16,193)	CI	5.2–6.1	10.3–11.4	18.4–19.8	63.6–65.3	82.9–84.2
Black	%	7.2	8.9	14.8	69.0	83.8
(n=1,359)	CI	5.8–9.0	7.3–10.8	12.8–17.0	66.2–71.7	81.5–86.0
Hispanic	%	5.1	9.2	18.8	66.9	85.7
(n=918)	CI	3.7–6.9	7.4–11.4	16.1–21.8	63.4–70.2	83.0–88.0
8th	%	2.7	6.5	14.2	76.6	90.8
(n=9,259)	CI	2.4–3.1	6.0–7.0	13.5–15.0	75.7–77.5	90.2–91.4
10th	%	6.4	11.8	20.9	60.9	81.8
(n=6,638)	CI	5.8–7.1	11.0–12.7	19.8–22.0	59.5–62.2	80.7–82.8
12th	%	9.2	14.0	22.1	54.7	76.8
(n=5,091)	CI	8.4–10.2	12.9–15.1	20.9–23.5	53.1–56.2	75.5–78.1

Note: The confidence interval (CI) is 95 percent.

Table TN.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.6	10.0	18.0	64.4	82.4
(n=20,982)	CI	7.2–8.1	9.5–10.4	17.4–18.6	63.7–65.2	81.8–83.0
Male	%	8.9	9.9	17.0	64.2	81.2
(n=10,207)	CI	8.3–9.6	9.2–10.6	16.2–17.8	63.1–65.3	80.3–82.1
Female	%	5.6	9.9	19.1	65.5	84.5
(n=10,202)	CI	5.1–6.1	9.3–10.6	18.2–20.0	64.4–66.5	83.7–85.3
White	%	7.5	10.4	18.5	63.6	82.1
(n=16,196)	CI	7.0–8.0	9.9–11.0	17.8–19.2	62.7–64.5	81.4–82.8
Black	%	7.1	6.8	15.3	70.8	86.1
(n=1,355)	CI	5.6–8.9	5.4–8.6	13.4–17.5	68.0–73.4	83.7–88.1
Hispanic	%	5.6	7.2	15.8	71.4	87.2
(n=917)	CI	4.2–7.5	5.5–9.4	13.3–18.6	68.0–74.6	84.6–89.4
8th	%	2.7	5.5	13.6	78.2	91.9
(n=9,256)	CI	2.3–3.0	5.0–6.0	12.9–14.4	77.3–79.1	91.3–92.4
10th	%	6.9	10.4	20.1	62.6	82.7
(n=6,634)	CI	6.3–7.7	9.6–11.2	19.0–21.2	61.3–63.9	81.6–83.7
12th	%	14.0	14.6	20.6	50.8	71.4
(n=5,092)	CI	12.9–15.1	13.5–15.7	19.4–21.9	49.3–52.4	70.0–72.8

Note: The confidence interval (CI) is 95 percent.

Table TN.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	13.9	16.0	16.8	53.2	70.0
(n=20,953)	CI	13.4–14.5	15.5–16.6	16.2–17.4	52.4–54.0	69.3–70.7
Male	%	15.0	15.1	16.4	53.5	69.9
(n=10,196)	CI	14.2–15.9	14.4–15.9	15.5–17.2	52.4–54.6	68.8–70.9
Female	%	12.1	16.8	17.6	53.5	71.1
(n=10,188)	CI	11.4–12.9	16.0–17.6	16.8–18.5	52.4–54.6	70.1–72.1
White	%	14.2	16.6	17.1	52.1	69.2
(n=16,176)	CI	13.6–14.8	16.0–17.3	16.4–17.8	51.2–53.0	68.3–70.0
Black	%	10.7	12.0	14.3	63.0	77.3
(n=1,350)	CI	8.9–12.7	10.2–14.1	12.3–16.5	60.1–65.9	74.7–79.7
Hispanic	%	9.4	12.3	17.0	61.3	78.3
(n=916)	CI	7.5–11.6	10.2–14.9	14.4–20.0	57.7–64.8	75.2–81.1
8th	%	7.3	11.5	14.4	66.7	81.1
(n=9,246)	CI	6.8–7.9	10.9–12.2	13.7–15.2	65.7–67.7	80.3–81.9
10th	%	14.0	18.0	18.2	49.8	68.0
(n=6,617)	CI	13.1–15.0	17.0–19.1	17.2–19.3	48.4–51.1	66.7–69.2
12th	%	21.4	19.0	18.0	41.6	59.7
(n=5,090)	CI	20.1–22.7	17.8–20.2	16.8–19.3	40.1–43.2	58.1–61.2

Note: The confidence interval (CI) is 95 percent.

Table TN.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	12.9	11.6	14.2	61.3	75.4
(n=20,945)	CI	12.4–13.5	11.1–12.1	13.6–14.7	60.5–62.0	74.8–76.1
Male	%	13.6	10.5	14.2	61.7	75.9
(n=10,193)	CI	12.8–14.4	9.9–11.2	13.4–15.0	60.6–62.8	74.9–76.8
Female	%	11.5	12.5	14.3	61.7	76.0
(n=10,180)	CI	10.8–12.2	11.7–13.2	13.5–15.1	60.6–62.8	75.0–77.0
White	%	12.2	11.4	14.1	62.3	76.4
(n=16,164)	CI	11.6–12.8	10.8–12.0	13.5–14.8	61.4–63.1	75.6–77.1
Black	%	16.4	13.0	12.9	57.7	70.6
(n=1,353)	CI	14.3–18.8	11.0–15.2	11.0–15.1	54.8–60.7	67.8–73.3
Hispanic	%	10.2	9.8	13.9	66.1	80.1
(n=916)	CI	8.0–12.8	7.8–12.1	11.6–16.6	62.6–69.5	76.9–82.9
8th	%	4.9	6.5	11.0	77.6	88.6
(n=9,237)	CI	4.4–5.4	6.0–7.0	10.4–11.7	76.7–78.5	88.0–89.3
10th	%	15.2	13.3	14.8	56.7	71.5
(n=6,625)	CI	14.3–16.2	12.4–14.2	13.9–15.9	55.3–58.0	70.3–72.7
12th	%	19.5	15.7	16.9	47.9	64.8
(n=5,083)	CI	18.4–20.8	14.5–16.8	15.8–18.1	46.3–49.5	63.3–66.3

Note: The confidence interval (CI) is 95 percent.

Table TN.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	64.0	25.1	10.9	36.0
(n=20,887)	CI	63.2–64.7	24.5–25.8	10.4–11.4	35.3–36.8
Male	%	65.3	24.1	10.6	34.7
(n=10,155)	CI	64.3–66.4	23.1–25.0	10.0–11.3	33.6–35.7
Female	%	62.5	26.7	10.9	37.5
(n=10,163)	CI	61.4–63.6	25.7–27.6	10.2–11.6	36.4–38.6
White	%	64.1	25.3	10.6	35.9
(n=16,120)	CI	63.2–64.9	24.6–26.1	10.0–11.1	35.1–36.8
Black	%	68.2	21.0	10.8	31.8
(n=1,343)	CI	65.4–70.9	18.6–23.5	9.1–12.8	29.1–34.6
Hispanic	%	60.0	26.6	13.4	40.0
(n=922)	CI	56.3–63.6	23.5–30.0	10.9–16.2	36.4–43.7
8th	%	60.7	27.7	11.5	39.3
(n=9,209)	CI	59.7–61.8	26.8–28.7	10.9–12.3	38.2–40.3
10th	%	64.1	24.5	11.4	35.9
(n=6,614)	CI	62.8–65.4	23.3–25.7	10.6–12.4	34.6–37.2
12th	%	67.5	22.9	9.6	32.5
(n=5,064)	CI	66.0–69.0	21.6–24.2	8.7–10.6	31.0–34.0

Note: The confidence interval (CI) is 95 percent.

Table TN.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	47.5	33.6	18.9	52.5
(n=20,887)	CI	46.8–48.3	32.9–34.3	18.3–19.5	51.7–53.2
Male	%	49.9	32.2	17.9	50.1
(n=10,164)	CI	48.8–51.0	31.2–33.2	17.1–18.8	49.0–51.2
Female	%	44.8	35.4	19.7	55.2
(n=10,155)	CI	43.7–45.9	34.4–36.5	18.9–20.6	54.1–56.3
White	%	46.9	34.3	18.8	53.1
(n=16,126)	CI	46.0–47.8	33.5–35.2	18.1–19.5	52.2–54.0
Black	%	57.7	25.5	16.8	42.3
(n=1,338)	CI	54.7–60.5	23.1–28.1	14.7–19.2	39.5–45.3
Hispanic	%	45.9	35.3	18.8	54.1
(n=921)	CI	42.2–49.5	31.9–38.8	16.1–22.0	50.5–57.8
8th	%	44.6	35.2	20.2	55.4
(n=9,206)	CI	43.5–45.7	34.1–36.2	19.3–21.1	54.3–56.5
10th	%	46.5	33.8	19.8	53.5
(n=6,616)	CI	45.1–47.9	32.5–35.1	18.7–20.9	52.1–54.9
12th	%	52.0	31.6	16.4	48.0
(n=5,065)	CI	50.4–53.6	30.2–33.0	15.2–17.6	46.4–49.6

Note: The confidence interval (CI) is 95 percent.

Table TN.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	38.6	32.2	29.2	61.4
(n=20,895)	CI	37.9–39.4	31.5–32.9	28.5–29.9	60.6–62.1
Male	%	41.5	29.8	28.8	58.5
(n=10,158)	CI	40.4–42.6	28.8–30.8	27.8–29.8	57.4–59.6
Female	%	35.4	35.1	29.5	64.6
(n=10,172)	CI	34.3–36.5	34.0–36.2	28.5–30.5	63.5–65.7
White	%	37.5	33.2	29.3	62.5
(n=16,133)	CI	36.6–38.4	32.4–34.1	28.5–30.1	61.6–63.4
Black	%	50.1	24.6	25.2	49.9
(n=1,342)	CI	47.2–53.1	22.2–27.2	22.7–27.9	46.9–52.8
Hispanic	%	40.9	31.8	27.3	59.1
(n=920)	CI	37.3–44.5	28.6–35.3	24.1–30.7	55.5–62.7
8th	%	34.5	32.1	33.4	65.5
(n=9,214)	CI	33.5–35.6	31.1–33.1	32.3–34.4	64.4–66.5
10th	%	38.8	32.3	28.9	61.2
(n=6,613)	CI	37.5–40.1	31.1–33.6	27.6–30.2	59.9–62.5
12th	%	43.2	32.1	24.8	56.8
(n=5,068)	CI	41.6–44.7	30.6–33.5	23.4–26.1	55.3–58.4

Note: The confidence interval (CI) is 95 percent.

Behavioral Health Planning Region 1 Results Summary

Table BH1.1. Lifetime Prescription Drug Misuse

On how many occasions (if any) have you ever <u>used prescription drugs not prescribed to you?</u>									
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	91.4	3.7	2.1	0.7	0.7	0.2	1.2	8.6
(n=2,057)	CI	89.7–92.9	2.7–4.9	1.4–3.1	0.4–1.3	0.4–1.4	0.1–0.5	0.7–2.1	7.1–10.3
Male	%	91.7	2.6	1.9	0.8	0.7	0.0	2.3	8.3
(n=970)	CI	89.0–93.7	1.6–4.5	1.0–3.6	0.3–1.8	0.2–1.8	–	1.3–4.1	6.3–11.0
Female	%	92.0	4.4	2.1	0.4	0.8	0.3	0.0	8.0
(n=1,028)	CI	89.6–93.8	3.1–6.3	1.3–3.6	0.2–1.2	0.3–1.9	0.1–1.0	–	6.2–10.4
White	%	–	–	–	–	–	–	–	–
(n=1,660)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	95.0	2.8	1.1	0.2	0.5	0.2	0.2	5.0
(n=1,308)	CI	93.6–96.0	2.1–3.9	0.7–1.9	0.0–0.6	0.3–1.1	0.0–0.6	0.1–0.7	4.0–6.4
10th	%	89.7	3.7	2.4	1.6	0.8	0.4	1.4	10.3
(n=505)	CI	86.8–92.1	2.4–5.8	1.3–4.1	0.8–3.1	0.3–2.1	0.1–1.6	0.7–2.9	7.9–13.2
12th	%	89.5	4.5	2.8	0.4	0.8	0.0	2.0	10.5
(n=244)	CI	85.0–92.8	2.5–7.9	1.3–5.9	0.1–2.9	0.2–3.2	–	0.8–4.8	7.2–15.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.2. Lifetime Alcohol Use

On how many occasions (if any) have you ever <u>had one or more drinks of an alcoholic beverage?</u>									
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	60.4	17.1	7.5	3.5	4.3	2.9	4.4	39.6
(n=2,071)	CI	57.7–63.0	15.2–19.2	6.1–9.2	2.6–4.7	3.3–5.5	2.0–4.1	3.3–5.8	37.0–42.3
Male	%	60.5	15.6	7.1	4.1	4.1	2.3	6.4	39.5
(n=980)	CI	56.6–64.3	13.1–18.5	5.1–9.6	2.7–6.2	2.7–6.0	1.2–4.1	4.5–9.0	35.7–43.4
Female	%	60.0	18.5	8.4	2.8	4.5	3.5	2.2	40.0
(n=1,031)	CI	56.2–63.7	15.8–21.7	6.4–11.0	1.8–4.5	3.1–6.4	2.2–5.5	1.3–3.8	36.3–43.8
White	%	–	–	–	–	–	–	–	–
(n=1,674)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	71.1	19.1	3.9	1.9	2.4	0.7	0.9	28.9
(n=1,317)	CI	68.6–73.5	17.1–21.4	3.0–5.1	1.3–2.8	1.7–3.3	0.4–1.3	0.5–1.6	26.5–31.4
10th	%	57.6	16.1	7.8	3.7	6.5	3.1	5.1	42.4
(n=508)	CI	53.3–61.9	13.1–19.5	5.8–10.5	2.4–5.8	4.6–9.0	1.9–5.1	3.5–7.4	38.1–46.7
12th	%	52.2	16.1	10.8	4.8	4.0	4.8	7.2	47.8
(n=246)	CI	46.0–58.4	12.0–21.2	7.5–15.4	2.7–8.3	2.2–7.3	2.7–8.3	4.6–11.2	41.6–54.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.3. Lifetime Binge Drinking

On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>									
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	80.3	6.3	3.4	2.8	2.5	1.8	2.8	19.7
(n=2,070)	CI	77.9–82.6	5.1–7.9	2.5–4.7	2.0–3.9	1.7–3.7	1.1–2.9	1.9–4.0	17.4–22.1
Male	%	78.3	7.0	3.5	2.5	2.2	1.5	5.0	21.7
(n=980)	CI	74.5–81.7	5.1–9.6	2.2–5.4	1.5–4.2	1.2–4.1	0.7–3.2	3.3–7.4	18.3–25.5
Female	%	82.3	5.5	3.6	3.2	2.9	1.6	0.8	17.7
(n=1,031)	CI	79.0–85.2	4.0–7.6	2.3–5.5	2.1–5.1	1.8–4.8	0.7–3.2	0.3–2.0	14.8–21.0
White	%	–	–	–	–	–	–	–	–
(n=1,672)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	92.9	3.1	1.6	0.8	0.8	0.4	0.5	7.1
(n=1,319)	CI	91.4–94.1	2.3–4.2	1.0–2.4	0.5–1.5	0.4–1.4	0.2–0.9	0.2–1.0	5.9–8.6
10th	%	77.4	7.5	3.9	4.3	2.4	1.4	3.1	22.6
(n=506)	CI	73.5–80.8	5.5–10.1	2.5–6.0	2.9–6.5	1.3–4.1	0.7–2.9	1.9–5.1	19.2–26.5
12th	%	70.6	8.5	4.8	3.2	4.4	3.6	4.8	29.4
(n=245)	CI	64.6–75.9	5.6–12.7	2.8–8.4	1.6–6.3	2.5–7.9	1.9–6.9	2.8–8.4	24.1–35.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.4. Lifetime Cigarette Use

On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>									
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.2	8.3	3.4	2.0	1.2	1.0	5.0	20.8
(n=2,073)	CI	76.9–81.4	6.9–9.8	2.5–4.6	1.3–2.9	0.7–2.0	0.6–1.7	3.9–6.5	18.6–23.1
Male	%	77.4	9.2	3.2	1.9	1.2	1.4	5.7	22.6
(n=981)	CI	73.8–80.7	7.2–11.8	2.0–5.0	1.1–3.4	0.5–2.7	0.6–2.9	4.0–8.1	19.3–26.2
Female	%	81.5	7.1	3.2	2.0	1.2	0.6	4.4	18.5
(n=1,032)	CI	78.3–84.3	5.5–9.2	2.1–4.8	1.1–3.5	0.6–2.4	0.3–1.2	2.9–6.5	15.7–21.7
White	%	–	–	–	–	–	–	–	–
(n=1,675)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	86.0	8.1	2.4	0.8	0.5	1.1	1.1	14.0
(n=1,319)	CI	84.0–87.7	6.8–9.7	1.7–3.4	0.4–1.4	0.3–1.1	0.7–1.9	0.6–1.8	12.3–16.0
10th	%	74.9	9.8	3.7	3.1	1.8	0.6	6.1	25.1
(n=508)	CI	70.9–78.5	7.5–12.7	2.4–5.8	1.9–5.1	0.9–3.4	0.2–1.8	4.3–8.5	21.5–29.1
12th	%	76.7	6.8	4.0	2.0	1.2	1.2	8.0	23.3
(n=246)	CI	71.0–81.6	4.3–10.7	2.2–7.3	0.8–4.8	0.4–3.7	0.4–3.7	5.2–12.2	18.4–29.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	66.9	10.5	4.8	2.3	2.6	2.1	10.7	33.1
(n=2,068)	CI	64.3–69.5	9.0–12.3	3.8–6.1	1.6–3.2	1.9–3.7	1.5–3.0	9.0–12.7	30.5–35.7
Male	%	64.7	10.2	5.4	2.4	3.1	1.8	12.4	35.3
(n=979)	CI	60.8–68.4	8.1–12.8	3.9–7.5	1.4–4.0	1.9–5.0	1.1–2.9	9.8–15.6	31.6–39.2
Female	%	69.3	10.9	4.3	2.3	2.2	2.2	8.8	30.7
(n=1,029)	CI	65.6–72.7	8.7–13.6	3.0–6.2	1.5–3.6	1.4–3.6	1.4–3.6	6.6–11.5	27.3–34.4
White	%	–	–	–	–	–	–	–	–
(n=1,671)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	78.4	9.6	3.3	1.9	1.9	2.0	2.9	21.6
(n=1,318)	CI	76.1–80.5	8.2–11.4	2.5–4.5	1.3–2.8	1.3–2.8	1.3–2.9	2.1–3.9	19.5–23.9
10th	%	59.4	11.0	6.7	3.4	3.2	3.2	13.2	40.6
(n=505)	CI	55.0–63.6	8.6–14.1	4.8–9.2	2.1–5.3	1.9–5.1	1.9–5.1	10.5–16.5	36.4–45.0
12th	%	62.9	10.9	4.4	1.6	2.8	1.2	16.1	37.1
(n=245)	CI	56.7–68.7	7.6–15.4	2.5–7.9	0.6–4.2	1.3–5.8	0.4–3.7	12.0–21.3	31.3–43.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	85.9	5.3	1.3	0.9	0.9	0.9	4.8	14.1
(n=2,061)	CI	83.8–87.7	4.2–6.7	0.8–2.1	0.5–1.7	0.5–1.6	0.5–1.4	3.6–6.3	12.3–16.2
Male	%	78.5	7.3	1.9	0.6	1.5	1.3	8.9	21.5
(n=976)	CI	74.9–81.7	5.4–9.8	1.1–3.4	0.2–1.8	0.7–2.9	0.7–2.4	6.7–11.8	18.3–25.1
Female	%	92.9	3.6	0.6	0.9	0.4	0.5	1.1	7.1
(n=1,025)	CI	90.7–94.6	2.4–5.4	0.3–1.4	0.4–2.1	0.1–1.1	0.2–1.1	0.5–2.5	5.4–9.3
White	%	–	–	–	–	–	–	–	–
(n=1,666)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=90)	CI	–	–	–	–	–	–	–	–
8th	%	91.9	4.0	1.0	0.5	0.6	0.6	1.4	8.1
(n=1,312)	CI	90.3–93.3	3.1–5.3	0.6–1.7	0.2–1.0	0.3–1.2	0.3–1.2	0.9–2.2	6.7–9.7
10th	%	83.0	5.9	1.8	1.2	1.2	2.0	4.9	17.0
(n=505)	CI	79.5–86.1	4.2–8.3	0.9–3.4	0.5–2.6	0.5–2.6	1.1–3.6	3.3–7.2	13.9–20.5
12th	%	82.6	6.1	1.2	1.2	0.8	0.0	8.1	17.4
(n=244)	CI	77.3–86.9	3.7–9.9	0.4–3.7	0.4–3.7	0.2–3.2	–	5.3–12.3	13.1–22.7

Note: The confidence interval (CI) is 95 percent.

Table BH1.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	79.4	7.1	2.4	2.3	1.7	1.6	5.4	20.6
(n=2,065)	CI	77.0-81.7	5.8-8.8	1.7-3.5	1.5-3.3	1.1-2.6	1.0-2.6	4.1-6.9	18.3-23.0
Male	%	79.3	5.4	2.9	1.9	2.0	1.3	7.1	20.7
(n=978)	CI	75.7-82.6	3.8-7.6	1.8-4.8	1.0-3.6	1.1-3.6	0.6-2.9	5.1-9.8	17.4-24.3
Female	%	80.4	8.1	1.9	2.7	1.6	1.7	3.6	19.6
(n=1,027)	CI	77.0-83.4	6.1-10.6	1.1-3.4	1.7-4.3	0.9-2.9	1.0-3.1	2.3-5.5	16.6-23.0
White	%	-	-	-	-	-	-	-	-
(n=1,671)	CI	-	-	-	-	-	-	-	-
Black	%	-	-	-	-	-	-	-	-
(n=53)	CI	-	-	-	-	-	-	-	-
Hispanic	%	-	-	-	-	-	-	-	-
(n=91)	CI	-	-	-	-	-	-	-	-
8th	%	91.4	3.7	1.1	0.6	1.0	1.1	1.0	8.6
(n=1,314)	CI	89.8-92.8	2.8-4.9	0.7-1.9	0.3-1.2	0.6-1.7	0.7-1.9	0.6-1.7	7.2-10.2
10th	%	74.2	8.9	3.3	3.7	2.2	1.4	6.3	25.8
(n=506)	CI	70.2-77.8	6.7-11.7	2.1-5.3	2.4-5.8	1.2-3.9	0.7-2.9	4.5-8.8	22.2-29.8
12th	%	72.6	8.9	2.8	2.4	2.0	2.4	8.9	27.4
(n=245)	CI	66.7-77.8	5.9-13.1	1.3-5.8	1.1-5.3	0.8-4.8	1.1-5.3	5.9-13.1	22.2-33.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	98.4	0.7	0.0	0.1	0.3	0.0	0.5	1.6
(n=2,072)	CI	97.5-99.0	0.3-1.3	0.0-0.2	0.0-0.9	0.1-0.9	0.0-0.2	0.2-1.0	1.0-2.5
Male	%	98.4	0.6	0.0	0.0	0.5	0.0	0.5	1.6
(n=980)	CI	96.8-99.2	0.2-2.2	-	-	0.1-1.8	-	0.2-1.4	0.8-3.2
Female	%	99.0	0.7	0.0	0.0	0.2	0.0	0.1	1.0
(n=1,032)	CI	98.0-99.5	0.3-1.5	-	-	0.0-0.8	-	0.0-0.9	0.5-2.0
White	%	-	-	-	-	-	-	-	-
(n=1,674)	CI	-	-	-	-	-	-	-	-
Black	%	-	-	-	-	-	-	-	-
(n=55)	CI	-	-	-	-	-	-	-	-
Hispanic	%	-	-	-	-	-	-	-	-
(n=91)	CI	-	-	-	-	-	-	-	-
8th	%	99.1	0.4	0.1	0.0	0.2	0.1	0.2	0.9
(n=1,318)	CI	98.4-99.5	0.2-0.9	0.0-0.5	-	0.0-0.6	0.0-0.5	0.1-0.7	0.5-1.6
10th	%	98.0	0.8	0.0	0.0	0.4	0.0	0.8	2.0
(n=508)	CI	96.4-98.9	0.3-2.1	-	-	0.1-1.6	-	0.3-2.1	1.1-3.6
12th	%	98.0	0.8	0.0	0.4	0.4	0.0	0.4	2.0
(n=246)	CI	95.2-99.2	0.2-3.2	-	0.1-2.8	0.1-2.8	-	0.1-2.8	0.8-4.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	96.7	1.0	0.9	0.1	0.5	0.2	0.7	3.3
(n=2,069)	CI	95.5–97.6	0.6–1.7	0.4–1.7	0.0–0.2	0.2–1.2	0.0–0.8	0.4–1.4	2.4–4.5
Male	%	95.7	0.8	1.2	0.1	1.0	0.3	1.0	4.3
(n=978)	CI	93.5–97.2	0.4–1.7	0.5–2.8	0.0–0.4	0.4–2.6	0.0–2.0	0.5–2.2	2.8–6.5
Female	%	98.0	1.3	0.4	0.1	0.1	0.0	0.3	2.0
(n=1,031)	CI	96.6–98.8	0.6–2.6	0.1–1.1	0.0–0.4	0.0–0.4	–	0.1–1.0	1.2–3.4
White	%	–	–	–	–	–	–	–	–
(n=1,672)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	98.3	0.8	0.2	0.2	0.2	0.1	0.2	1.7
(n=1,316)	CI	97.4–98.8	0.5–1.5	0.1–0.7	0.0–0.6	0.1–0.7	0.0–0.5	0.1–0.7	1.2–2.6
10th	%	96.7	1.4	0.8	0.0	0.0	0.0	1.2	3.3
(n=508)	CI	94.7–97.9	0.7–2.9	0.3–2.1	–	–	–	0.5–2.6	2.1–5.3
12th	%	95.2	0.8	1.6	0.0	1.2	0.4	0.8	4.8
(n=245)	CI	91.6–97.2	0.2–3.2	0.6–4.2	–	0.4–3.7	0.1–2.8	0.2–3.2	2.8–8.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.10. Age of First Prescription Drug Misuse

		How old were you when you first <u>used prescription drugs not prescribed to you?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	93.2	0.9	0.3	0.7	0.8	1.5	0.7	1.2	0.6	13.67
(n=2,074)	CI	91.6–94.5	0.6–1.6	0.2–0.7	0.4–1.2	0.5–1.4	0.9–2.4	0.3–1.4	0.7–2.1	0.2–1.4	13.2–14.2
Male	%	94.7	0.6	0.3	0.4	0.7	1.1	0.6	1.0	0.7	14.01
(n=982)	CI	92.4–96.3	0.2–1.3	0.1–1.1	0.1–1.0	0.3–1.7	0.5–2.6	0.1–2.2	0.4–2.5	0.2–2.3	13.2–14.8
Female	%	92.6	0.7	0.3	1.0	1.0	1.9	0.8	1.2	0.5	13.71
(n=1,032)	CI	90.3–94.4	0.4–1.5	0.1–0.7	0.5–1.8	0.5–2.0	1.0–3.4	0.3–2.0	0.5–2.7	0.1–2.1	13.1–14.3
White	%	–	–	–	–	–	–	–	–	–	–
(n=1,676)	CI	–	–	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–	–	–
8th	%	96.1	1.1	0.6	0.8	0.9	0.6	0.0	0.0	0.0	11.85
(n=1,320)	CI	94.9–97.0	0.6–1.8	0.3–1.2	0.4–1.4	0.5–1.6	0.3–1.2	–	–	–	11.4–12.2
10th	%	92.0	1.0	0.4	1.4	1.2	2.0	0.8	1.2	0.2	13.34
(n=508)	CI	89.3–94.0	0.4–2.3	0.1–1.6	0.7–2.9	0.5–2.6	1.1–3.6	0.3–2.1	0.5–2.6	0.0–1.4	12.7–14.0
12th	%	91.6	0.8	0.0	0.0	0.4	2.0	1.2	2.4	1.6	14.86
(n=246)	CI	87.4–94.5	0.2–3.2	–	–	0.1–2.8	0.8–4.8	0.4–3.7	1.1–5.3	0.6–4.2	13.9–15.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.11. Age of First Alcohol Use

How old were you when you first had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	63.5	4.3	2.4	3.5	5.8	5.5	6.5	5.8	2.9	13.72
(n=2,069)	CI	60.8–66.1	3.3–5.5	1.8–3.1	2.8–4.3	4.8–7.0	4.4–7.0	5.1–8.1	4.4–7.5	2.0–4.3	13.5–13.9
Male	%	64.4	4.4	3.2	4.1	5.5	5.0	6.7	4.6	2.3	13.49
(n=981)	CI	60.5–68.1	3.1–6.2	2.2–4.5	3.0–5.5	4.1–7.5	3.4–7.3	4.7–9.4	2.9–7.1	1.2–4.3	13.2–13.8
Female	%	62.3	3.5	1.6	3.2	6.2	6.4	6.2	7.0	3.7	13.99
(n=1,029)	CI	58.4–66.0	2.3–5.2	1.0–2.7	2.3–4.3	4.8–8.0	4.8–8.6	4.4–8.5	4.9–9.8	2.2–6.0	13.7–14.3
White	%	–	–	–	–	–	–	–	–	–	–
(n=1,675)	CI	–	–	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–	–	–
8th	%	74.7	5.0	3.9	6.2	7.8	2.4	0.0	0.0	0.0	11.95
(n=1,318)	CI	72.3–77.0	4.0–6.3	3.0–5.1	5.0–7.7	6.5–9.4	1.7–3.3	–	–	–	11.8–12.1
10th	%	61.6	3.3	2.4	3.7	6.7	8.3	9.8	3.0	1.2	13.60
(n=505)	CI	57.3–65.8	2.1–5.3	1.3–4.1	2.4–5.8	4.8–9.2	6.2–11.0	7.5–12.8	1.8–4.8	0.5–2.6	13.3–13.9
12th	%	53.8	4.4	0.8	0.4	2.8	6.0	9.6	14.5	7.6	14.82
(n=246)	CI	47.6–60.0	2.5–7.8	0.2–3.2	0.1–2.8	1.3–5.8	3.7–9.8	6.5–14.0	10.6–19.4	4.9–11.7	14.4–15.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.12. Age of First Cigarette Use

		How old were you when you first <u>smoked a cigarette, even just a puff?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	79.8	4.2	1.2	3.0	2.6	3.0	2.8	2.1	1.2	13.09
(n=2,072)	CI	77.5–82.0	3.3–5.3	0.8–1.9	2.2–4.0	2.0–3.5	2.2–4.2	1.9–4.0	1.4–3.3	0.7–2.2	12.8–13.4
Male	%	78.2	4.3	1.8	3.5	2.4	2.1	3.7	2.4	1.6	13.13
(n=981)	CI	74.6–81.4	3.1–6.1	1.1–3.1	2.3–5.4	1.5–3.7	1.2–3.7	2.3–5.8	1.3–4.4	0.7–3.5	12.7–13.6
Female	%	82.0	3.6	0.7	2.6	2.9	3.9	2.0	1.6	0.7	13.03
(n=1,032)	CI	78.9–84.8	2.5–5.1	0.4–1.4	1.7–4.1	1.9–4.3	2.6–5.7	1.1–3.7	0.8–3.2	0.2–2.1	12.6–13.4
White	%	–	–	–	–	–	–	–	–	–	–
(n=1,675)	CI	–	–	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–	–	–
8th	%	86.7	4.3	1.7	3.0	3.0	1.2	0.1	0.0	0.1	11.67
(n=1,391)	CI	84.7–88.4	3.3–5.6	1.2–2.6	2.2–4.0	2.2–4.0	0.7–2.0	0.0–0.5	–	0.0–0.5	11.5–11.9
10th	%	75.3	5.7	1.6	2.7	3.7	4.3	4.3	2.0	0.4	12.90
(n=507)	CI	71.4–78.9	4.0–8.1	0.8–3.1	1.6–4.6	2.4–5.8	2.9–6.5	2.9–6.5	1.1–3.6	0.1–1.6	12.5–13.3
12th	%	77.5	2.4	0.4	3.2	1.2	3.6	4.0	4.4	3.2	14.18
(n=246)	CI	71.9–82.3	1.1–5.3	0.1–2.8	1.6–6.3	0.4–3.7	1.9–6.8	2.2–7.3	2.5–7.8	1.6–6.3	13.6–14.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	68.3	1.7	1.4	2.6	5.0	5.0	6.2	5.9	3.9	14.27
(n=2,063)	CI	65.7–70.8	1.2–2.5	1.0–1.9	1.9–3.4	4.1–6.0	3.9–6.4	5.0–7.7	4.5–7.6	2.8–5.5	14.1–14.5
Male	%	66.6	1.3	1.5	3.3	5.3	5.7	7.4	6.7	2.3	14.19
(n=973)	CI	62.7–70.2	0.8–2.1	0.9–2.4	2.2–4.8	4.0–7.1	4.0–8.0	5.5–9.9	4.7–9.5	1.2–4.5	13.9–14.4
Female	%	70.1	1.9	1.4	1.7	4.7	4.7	5.1	4.7	5.7	14.39
(n=1,030)	CI	66.4–73.5	1.1–3.2	0.9–2.2	1.1–2.6	3.6–6.2	3.4–6.6	3.7–7.1	3.1–7.1	3.8–8.4	14.1–14.7
White	%	–	–	–	–	–	–	–	–	–	–
(n=1,667)	CI	–	–	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–	–	–
8th	%	79.8	2.4	2.5	4.5	8.1	2.5	0.2	0.0	0.0	12.31
(n=1,315)	CI	77.6–81.9	1.7–3.4	1.8–3.5	3.5–5.8	6.7–9.7	1.8–3.5	0.0–0.6	–	–	12.2–12.5
10th	%	61.6	1.6	1.6	1.6	5.5	8.1	14.1	5.7	0.2	14.16
(n=502)	CI	57.2–65.7	0.8–3.1	0.8–3.1	0.8–3.1	3.9–7.9	6.0–10.9	11.3–17.4	4.0–8.2	0.0–1.4	13.9–14.4
12th	%	63.5	1.2	0.0	1.6	1.2	4.4	4.4	12.0	11.6	15.48
(n=246)	CI	57.2–69.2	0.4–3.7	–	0.6–4.2	0.4–3.7	2.5–7.8	2.5–7.8	8.5–16.7	8.2–16.3	15.1–15.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	79.3	1.3	0.7	2.4	3.1	3.0	5.1	3.2	1.9	14.11
(n=2,075)	CI	76.9–81.6	0.8–2.0	0.4–1.3	1.7–3.3	2.3–4.2	2.2–4.1	4.0–6.6	2.2–4.6	1.1–3.1	13.9–14.4
Male	%	79.3	1.6	0.9	2.3	2.9	2.3	5.0	4.1	1.6	14.05
(n=983)	CI	75.7–82.5	0.9–3.0	0.4–2.1	1.4–3.8	1.8–4.6	1.4–3.9	3.4–7.2	2.6–6.5	0.7–3.5	13.7–14.4
Female	%	80.2	0.6	0.4	2.3	3.3	3.4	5.2	2.6	2.0	14.26
(n=1,032)	CI	76.8–83.2	0.3–1.2	0.2–0.9	1.5–3.5	2.3–4.9	2.2–5.1	3.6–7.5	1.5–4.6	1.0–4.0	14.0–14.6
White	%	–	–	–	–	–	–	–	–	–	–
(n=1,678)	CI	–	–	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–	–	–
8th	%	91.6	1.2	0.7	2.4	3.0	1.1	0.0	0.0	0.1	12.28
(n=1,320)	CI	90.0–93.0	0.7–2.0	0.4–1.3	1.7–3.4	2.2–4.0	0.6–1.8	–	–	0.0–0.5	12.0–12.5
10th	%	74.0	1.4	1.0	3.1	3.5	5.5	8.6	2.5	0.4	13.87
(n=509)	CI	70.0–77.6	0.7–2.8	0.4–2.3	1.9–5.0	2.2–5.5	3.8–7.8	6.5–11.4	1.5–4.3	0.1–1.6	13.6–14.2
12th	%	72.3	1.2	0.4	1.6	2.8	2.4	6.8	7.2	5.2	14.90
(n=246)	CI	66.4–77.5	0.4–3.7	0.1–2.8	0.6–4.2	1.3–5.8	1.1–5.3	4.3–10.7	4.6–11.2	3.0–8.8	14.5–15.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.15. Past Month Prescription Drug Misuse

		During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	96.6	1.5	0.8	0.4	0.2	0.1	0.4	3.4
(n=2,072)	CI	95.4–97.5	0.9–2.4	0.4–1.5	0.2–1.0	0.1–0.5	0.0–0.9	0.1–1.1	2.5–4.6
Male	%	97.3	1.2	0.7	0.3	0.2	0.3	0.0	2.7
(n=979)	CI	95.5–98.4	0.5–2.8	0.2–1.9	0.1–1.1	0.0–0.9	0.0–2.0	–	1.6–4.5
Female	%	96.6	1.9	0.6	0.4	0.2	0.0	0.2	3.4
(n=1,033)	CI	95.0–97.7	1.1–3.3	0.3–1.4	0.1–1.6	0.0–0.8	–	0.1–0.8	2.3–5.0
White	%	–	–	–	–	–	–	–	–
(n=1,674)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	98.1	0.8	0.6	0.2	0.2	0.0	0.2	1.9
(n=1,317)	CI	97.2–98.7	0.4–1.4	0.3–1.2	0.1–0.7	0.0–0.6	–	0.0–0.6	1.3–2.8
10th	%	96.1	1.8	1.0	0.6	0.4	0.0	0.2	3.9
(n=510)	CI	94.0–97.5	0.9–3.3	0.4–2.3	0.2–1.8	0.1–1.6	–	0.0–1.4	2.5–6.0
12th	%	95.6	2.0	0.8	0.4	0.0	0.4	0.8	4.4
(n=245)	CI	92.1–97.5	0.8–4.8	0.2–3.2	0.1–2.8	–	0.1–2.8	0.2–3.2	2.5–7.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.16. Past Month Alcohol Use

		During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	81.2	9.7	4.3	1.5	1.9	0.7	0.8	18.8
(n=2,058)	CI	78.8–83.3	8.1–11.6	3.2–5.7	1.0–2.4	1.2–2.9	0.3–1.3	0.4–1.6	16.7–21.2
Male	%	82.3	7.2	4.3	2.3	2.7	0.8	0.5	17.7
(n=972)	CI	78.8–85.3	5.3–9.7	2.8–6.5	1.3–3.8	1.6–4.5	0.3–2.3	0.1–1.8	14.7–21.2
Female	%	80.3	12.6	4.2	0.9	0.9	0.4	0.7	19.7
(n=1,026)	CI	76.9–83.3	10.1–15.6	2.9–6.2	0.4–2.1	0.4–2.0	0.1–1.1	0.3–1.4	16.7–23.1
White	%	–	–	–	–	–	–	–	–
(n=1,665)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=53)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=89)	CI	–	–	–	–	–	–	–	–
8th	%	89.9	5.7	2.1	0.4	1.1	0.4	0.5	10.1
(n=1,307)	CI	88.1–91.4	4.6–7.1	1.4–3.0	0.2–0.9	0.6–1.8	0.2–0.9	0.2–1.0	8.6–11.9
10th	%	80.2	9.2	3.9	2.9	2.2	0.8	0.8	19.8
(n=508)	CI	76.5–83.5	7.0–12.0	2.5–6.0	1.8–4.8	1.2–3.9	0.3–2.1	0.3–2.1	16.5–23.5
12th	%	73.2	14.2	6.9	1.2	2.4	0.8	1.2	26.8
(n=243)	CI	67.2–78.4	10.4–19.2	4.3–10.9	0.4–3.7	1.1–5.3	0.2–3.2	0.4–3.7	21.6–32.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.17. Past Month Binge Drinking

During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>									
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	89.0	5.0	3.1	0.8	1.0	0.4	0.6	11.0
(n=2,070)	CI	87.0–90.8	3.9–6.5	2.2–4.4	0.4–1.6	0.6–1.7	0.1–0.9	0.3–1.4	9.2–13.0
Male	%	87.8	5.8	2.6	1.4	1.3	0.6	0.6	12.2
(n=980)	CI	84.6–90.4	4.0–8.3	1.5–4.4	0.7–2.9	0.6–2.7	0.2–1.9	0.2–1.9	9.6–15.4
Female	%	90.3	4.7	3.6	0.4	0.8	0.1	0.2	9.7
(n=1,031)	CI	87.6–92.5	3.2–6.8	2.3–5.6	0.1–1.7	0.3–1.6	0.0–0.9	0.0–0.5	7.5–12.4
White	%	–	–	–	–	–	–	–	–
(n=1,676)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=90)	CI	–	–	–	–	–	–	–	–
8th	%	96.6	1.2	1.3	0.2	0.5	0.1	0.2	3.4
(n=1,316)	CI	95.4–97.4	0.7–2.0	0.8–2.1	0.0–0.6	0.2–1.0	0.0–0.5	0.1–0.7	2.6–4.6
10th	%	87.3	6.2	2.5	1.2	1.8	0.6	0.4	12.7
(n=509)	CI	84.1–89.9	4.4–8.7	1.5–4.3	0.5–2.6	0.9–3.4	0.2–1.8	0.1–1.6	10.1–15.9
12th	%	83.1	7.7	5.6	1.2	0.8	0.4	1.2	16.9
(n=245)	CI	77.8–87.3	4.9–11.7	3.4–9.3	0.4–3.7	0.2–3.2	0.1–2.8	0.4–3.7	12.7–22.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.18. Past Month Cigarette Use

During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>									
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	90.2	3.8	0.9	0.7	1.8	0.4	2.3	9.8
(n=2,069)	CI	88.4–91.7	2.9–5.0	0.5–1.6	0.4–1.4	1.2–2.7	0.2–0.8	1.5–3.4	8.3–11.6
Male	%	90.0	4.0	1.3	0.9	1.5	0.4	1.8	10.0
(n=979)	CI	87.3–92.2	2.8–5.9	0.6–2.8	0.4–2.3	0.8–2.8	0.1–1.3	1.0–3.4	7.8–12.7
Female	%	91.1	3.3	0.6	0.5	1.9	0.3	2.2	8.9
(n=1,030)	CI	88.7–93.1	2.2–4.9	0.3–1.3	0.2–1.3	1.1–3.5	0.1–0.8	1.2–3.8	6.9–11.3
White	%	–	–	–	–	–	–	–	–
(n=1,674)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=53)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	93.5	3.0	0.9	0.5	1.1	0.3	0.7	6.5
(n=1,314)	CI	92.0–94.7	2.2–4.0	0.5–1.6	0.3–1.1	0.7–1.9	0.1–0.8	0.4–1.3	5.3–8.0
10th	%	87.9	5.3	0.6	0.8	2.1	0.8	2.5	12.1
(n=510)	CI	84.8–90.5	3.6–7.6	0.2–1.8	0.3–2.1	1.2–3.8	0.3–2.1	1.5–4.3	9.5–15.2
12th	%	89.1	3.2	1.2	0.8	2.0	0.0	3.6	10.9
(n=245)	CI	84.6–92.4	1.6–6.3	0.4–3.7	0.2–3.2	0.8–4.8	–	1.9–6.9	7.6–15.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	79.2	6.0	3.5	1.6	2.3	1.6	5.7	20.8
(n=2,064)	CI	76.9–81.3	4.9–7.4	2.7–4.7	1.1–2.4	1.6–3.3	1.1–2.4	4.4–7.3	18.7–23.1
Male	%	77.3	6.7	4.5	2.1	1.9	1.7	5.8	22.7
(n=979)	CI	73.8–80.5	5.0–8.9	3.1–6.5	1.2–3.5	1.0–3.4	0.9–3.1	4.0–8.4	19.5–26.2
Female	%	81.4	5.8	2.4	1.1	2.7	1.3	5.2	18.6
(n=1,026)	CI	78.2–84.2	4.3–7.8	1.5–3.8	0.6–2.2	1.7–4.3	0.7–2.4	3.6–7.4	15.8–21.8
White	%	–	–	–	–	–	–	–	–
(n=1,670)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	87.0	5.0	2.4	0.8	2.2	1.6	0.8	13.0
(n=1,311)	CI	85.1–88.7	4.0–6.4	1.7–3.4	0.5–1.5	1.5–3.2	1.0–2.4	0.5–1.5	11.3–14.9
10th	%	72.9	8.2	4.9	3.1	2.3	2.0	6.6	27.1
(n=509)	CI	68.8–76.5	6.1–10.9	3.3–7.1	1.9–5.0	1.3–4.1	1.1–3.6	4.8–9.2	23.5–31.2
12th	%	77.7	4.9	3.2	0.8	2.4	1.2	9.7	22.3
(n=244)	CI	72.1–82.5	2.8–8.4	1.6–6.4	0.2–3.2	1.1–5.3	0.4–3.7	6.6–14.1	17.5–27.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	93.5	1.5	0.8	0.6	0.4	0.4	2.9	6.5
(n=2,073)	CI	91.9–94.8	1.0–2.3	0.4–1.5	0.3–1.3	0.2–0.8	0.1–1.1	2.0–4.1	5.2–8.1
Male	%	88.6	2.5	1.4	0.8	0.5	0.7	5.4	11.4
(n=979)	CI	85.6–91.1	1.5–4.2	0.6–2.9	0.4–2.0	0.2–1.4	0.2–2.3	3.7–7.8	8.9–14.4
Female	%	98.0	0.6	0.2	0.5	0.3	0.1	0.4	2.0
(n=1,034)	CI	96.8–98.8	0.3–1.3	0.1–0.8	0.1–1.6	0.1–1.0	0.0–0.4	0.1–1.1	1.2–3.2
White	%	–	–	–	–	–	–	–	–
(n=1,678)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=89)	CI	–	–	–	–	–	–	–	–
8th	%	96.1	1.3	0.7	0.5	0.2	0.1	1.1	3.9
(n=1,316)	CI	94.9–97.0	0.8–2.1	0.4–1.3	0.3–1.1	0.0–0.6	0.0–0.5	0.7–1.9	3.0–5.1
10th	%	93.2	1.9	0.4	0.6	1.0	0.2	2.7	6.8
(n=510)	CI	90.6–95.1	1.0–3.6	0.1–1.6	0.2–1.8	0.4–2.3	0.0–1.4	1.6–4.6	4.9–9.4
12th	%	91.2	1.2	1.2	0.8	0.0	0.8	4.8	8.8
(n=247)	CI	87.0–94.2	0.4–3.7	0.4–3.7	0.2–3.2	–	0.2–3.2	2.7–8.3	5.8–13.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	88.2	4.4	2.3	0.6	1.0	1.3	2.2	11.8
(n=2,073)	CI	86.3–90.0	3.3–5.7	1.6–3.3	0.3–1.2	0.5–1.7	0.8–2.2	1.5–3.3	10.0–13.7
Male	%	88.0	3.6	2.6	0.5	1.4	1.5	2.5	12.0
(n=981)	CI	85.0–90.4	2.4–5.4	1.5–4.4	0.2–1.3	0.6–2.9	0.7–3.0	1.5–4.3	9.6–15.0
Female	%	89.1	4.8	2.1	0.8	0.7	1.0	1.5	10.9
(n=1,032)	CI	86.3–91.3	3.4–6.9	1.3–3.6	0.3–1.8	0.3–1.3	0.4–2.3	0.8–3.0	8.7–13.7
White	%	–	–	–	–	–	–	–	–
(n=1,677)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=54)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	95.1	1.7	1.0	0.5	0.9	0.5	0.3	4.9
(n=1,316)	CI	93.8–96.2	1.1–2.5	0.6–1.7	0.2–1.0	0.5–1.6	0.3–1.1	0.1–0.8	3.8–6.2
10th	%	84.4	6.6	3.5	1.0	0.8	1.0	2.7	15.6
(n=511)	CI	81.0–87.3	4.8–9.1	2.2–5.5	0.4–2.3	0.3–2.1	0.4–2.3	1.6–4.6	12.7–19.0
12th	%	85.1	4.8	2.4	0.4	1.2	2.4	3.6	14.9
(n=246)	CI	80.1–89.1	2.7–8.3	1.1–5.3	0.1–2.8	0.4–3.7	1.1–5.3	1.9–6.8	10.9–19.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.6	0.3	0.3	0.2	0.2	0.0	0.4	1.4
(n=2,080)	CI	97.7–99.2	0.1–0.9	0.1–0.9	0.0–0.8	0.1–0.8	–	0.1–1.1	0.8–2.3
Male	%	98.2	0.5	0.4	0.1	0.4	0.0	0.3	1.8
(n=984)	CI	96.6–99.1	0.1–1.8	0.1–1.8	0.0–1.0	0.1–1.8	–	0.1–1.1	0.9–3.4
Female	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=1,035)	CI	99.0–99.9	0.0–0.9	0.0–0.9	–	–	–	–	0.1–1.0
White	%	–	–	–	–	–	–	–	–
(n=1,681)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	99.8	0.1	0.0	0.0	0.1	0.0	0.0	0.2
(n=1,321)	CI	99.4–100.0	0.0–0.5	–	–	0.0–0.5	–	–	0.0–0.6
10th	%	98.4	0.4	0.4	0.2	0.2	0.0	0.4	1.6
(n=512)	CI	96.9–99.2	0.1–1.5	0.1–1.5	0.0–1.4	0.0–1.4	–	0.1–1.5	0.8–3.1
12th	%	97.6	0.4	0.4	0.4	0.4	0.0	0.8	2.4
(n=247)	CI	94.7–98.9	0.1–2.8	0.1–2.8	0.1–2.8	0.1–2.8	–	0.2–3.2	1.1–5.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.1	0.3	0.2	0.0	0.0	0.0	0.3	0.9
(n=2,080)	CI	98.3–99.5	0.1–0.8	0.1–0.8	0.0–0.2	–	0.0–0.2	0.1–1.1	0.5–1.7
Male	%	99.1	0.3	0.5	0.0	0.0	0.0	0.1	0.9
(n=984)	CI	97.9–99.6	0.1–0.9	0.1–1.8	–	–	–	0.0–1.0	0.4–2.1
Female	%	99.8	0.1	0.0	0.1	0.0	0.0	0.0	0.2
(n=1,035)	CI	99.5–100.0	0.0–0.4	–	0.0–0.4	–	–	–	0.0–0.5
White	%	–	–	–	–	–	–	–	–
(n=1,681)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	99.5	0.3	0.1	0.1	0.0	0.1	0.0	0.5
(n=1,321)	CI	98.9–99.7	0.1–0.8	0.0–0.5	0.0–0.5	–	0.0–0.5	–	0.3–1.1
10th	%	99.4	0.2	0.2	0.0	0.0	0.0	0.2	0.6
(n=512)	CI	98.2–99.8	0.0–1.4	0.0–1.4	–	–	–	0.0–1.4	0.2–1.8
12th	%	98.4	0.4	0.4	0.0	0.0	0.0	0.8	1.6
(n=247)	CI	95.8–99.4	0.1–2.8	0.1–2.8	–	–	–	0.2–3.2	0.6–4.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.4	0.5	0.3	0.2	0.0	0.1	0.5	1.6
(n=2,078)	CI	97.4–99.1	0.2–1.2	0.1–1.0	0.0–0.8	–	0.0–0.9	0.2–1.2	0.9–2.6
Male	%	97.4	1.0	0.6	0.4	0.0	0.3	0.3	2.6
(n=984)	CI	95.4–98.5	0.4–2.5	0.1–2.2	0.1–1.8	–	0.0–2.0	0.1–1.1	1.5–4.6
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.1	0.2
(n=1,034)	CI	99.2–100.0	0.0–0.4	–	–	–	–	0.0–0.9	0.0–0.8
White	%	–	–	–	–	–	–	–	–
(n=1,680)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	99.8	0.2	0.0	0.0	0.0	0.0	0.0	0.2
(n=1,319)	CI	99.4–100.0	0.0–0.6	–	–	–	–	–	0.0–0.6
10th	%	98.6	0.6	0.0	0.2	0.0	0.0	0.6	1.4
(n=512)	CI	97.2–99.4	0.2–1.8	–	0.0–1.4	–	–	0.2–1.8	0.6–2.8
12th	%	96.8	0.8	0.8	0.4	0.0	0.4	0.8	3.2
(n=247)	CI	93.7–98.4	0.2–3.2	0.2–3.2	0.1–2.8	–	0.1–2.8	0.2–3.2	1.6–6.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.1	0.1	0.0	0.1	0.1	0.4	0.8
(n=2,079)	CI	98.4–99.6	0.0–0.4	0.0–0.9	–	0.0–0.9	0.0–0.5	0.1–1.1	0.4–1.6
Male	%	99.2	0.1	0.3	0.0	0.3	0.0	0.1	0.8
(n=984)	CI	97.6–99.7	0.0–1.0	0.0–2.0	–	0.0–2.0	–	0.0–1.0	0.3–2.4
Female	%	99.8	0.1	0.0	0.0	0.0	0.1	0.1	0.2
(n=1,035)	CI	99.2–99.9	0.0–0.4	–	–	–	0.0–0.9	0.0–0.4	0.1–0.8
White	%	–	–	–	–	–	–	–	–
(n=1,681)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	99.8	0.1	0.0	0.0	0.0	0.0	0.1	0.2
(n=1,320)	CI	99.4–100.0	0.0–0.5	–	–	–	–	0.0–0.5	0.0–0.6
10th	%	99.4	0.2	0.0	0.0	0.0	0.2	0.2	0.6
(n=512)	CI	98.2–99.8	0.0–1.4	–	–	–	0.0–1.4	0.0–1.4	0.2–1.8
12th	%	98.4	0.0	0.4	0.0	0.4	0.0	0.8	1.6
(n=247)	CI	95.8–99.4	–	0.1–2.8	–	0.1–2.8	–	0.2–3.2	0.6–4.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.1	0.1	0.2	0.1	0.1	0.0	0.3	0.9
(n=2,078)	CI	98.3–99.5	0.0–0.4	0.1–0.8	0.0–0.9	0.0–0.4	–	0.1–1.1	0.5–1.7
Male	%	99.3	0.0	0.4	0.0	0.1	0.0	0.1	0.7
(n=983)	CI	98.0–99.7	–	0.1–1.8	–	0.0–1.0	–	0.0–1.0	0.3–2.0
Female	%	99.7	0.2	0.1	0.0	0.1	0.0	0.0	0.3
(n=1,035)	CI	99.1–99.9	0.1–0.8	0.0–0.4	–	0.0–0.4	–	–	0.1–0.9
White	%	–	–	–	–	–	–	–	–
(n=1,680)	CI	–	–	–	–	–	–	–	–
Black	%	–	–	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–	–	–
(n=91)	CI	–	–	–	–	–	–	–	–
8th	%	99.7	0.2	0.1	0.0	0.1	0.0	0.0	0.3
(n=1,319)	CI	99.2–99.9	0.0–0.6	0.0–0.5	–	0.0–0.5	–	–	0.1–0.8
10th	%	99.2	0.2	0.2	0.0	0.2	0.0	0.2	0.8
(n=512)	CI	97.9–99.7	0.0–1.4	0.0–1.4	–	0.0–1.4	–	0.0–1.4	0.3–2.1
12th	%	98.4	0.0	0.4	0.4	0.0	0.0	0.8	1.6
(n=247)	CI	95.8–99.4	–	0.1–2.8	0.1–2.8	–	–	0.2–3.2	0.6–4.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you use ecstasy (MDMA)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	98.8	0.1	0.2	0.3	0.1	0.1	0.3	1.2
(n=2,078)	CI	97.9-99.4	0.0-0.4	0.0-0.8	0.1-1.1	0.0-0.5	0.0-0.9	0.1-1.1	0.6-2.1
Male	%	98.5	0.2	0.4	0.3	0.1	0.3	0.1	1.5
(n=984)	CI	96.9-99.3	0.0-0.9	0.1-1.8	0.0-2.0	0.0-1.0	0.0-2.0	0.0-1.0	0.7-3.1
Female	%	99.8	0.1	0.0	0.1	0.0	0.0	0.0	0.2
(n=1,034)	CI	99.2-100.0	0.0-0.4	-	0.0-0.9	-	-	-	0.0-0.8
White	%	-	-	-	-	-	-	-	-
(n=1,680)	CI	-	-	-	-	-	-	-	-
Black	%	-	-	-	-	-	-	-	-
(n=55)	CI	-	-	-	-	-	-	-	-
Hispanic	%	-	-	-	-	-	-	-	-
(n=91)	CI	-	-	-	-	-	-	-	-
8th	%	99.8	0.2	0.0	0.0	0.0	0.0	0.0	0.2
(n=1,319)	CI	99.4-100.0	0.0-0.6	-	-	-	-	-	0.0-0.6
10th	%	99.0	0.2	0.2	0.2	0.2	0.0	0.2	1.0
(n=512)	CI	97.7-99.6	0.0-1.4	0.0-1.4	0.0-1.4	0.0-1.4	-	0.0-1.4	0.4-2.3
12th	%	97.6	0.0	0.4	0.8	0.0	0.4	0.8	2.4
(n=247)	CI	94.7-98.9	-	0.1-2.8	0.2-3.2	-	0.1-2.8	0.2-3.2	1.1-5.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you use methamphetamines (crank, etc.)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.3	0.1	0.2	0.0	0.1	0.0	0.3	0.7
(n=2,079)	CI	98.5-99.6	0.0-0.4	0.0-0.8	-	0.0-0.4	-	0.1-1.1	0.4-1.5
Male	%	99.2	0.1	0.4	0.0	0.1	0.0	0.1	0.8
(n=984)	CI	98.0-99.7	0.0-1.0	0.1-1.8	-	0.0-0.4	-	0.0-1.0	0.3-2.0
Female	%	99.8	0.0	0.0	0.0	0.2	0.0	0.0	0.2
(n=1,036)	CI	99.2-100.0	-	-	-	0.0-0.8	-	-	0.0-0.8
White	%	97.1	1.6	0.6	0.2	0.2	0.1	0.2	2.9
(n=1,682)	CI	96.6-97.5	1.3-2.0	0.4-0.9	0.1-0.3	0.1-0.4	0.1-0.3	0.1-0.3	2.5-3.4
Black	%	95.5	2.3	1.2	0.3	0.3	0.1	0.2	4.5
(n=55)	CI	94.7-96.2	1.9-2.9	0.9-1.7	0.2-0.6	0.2-0.5	0.1-0.3	0.1-0.4	3.8-5.3
Hispanic	%	95.2	3.0	0.8	0.3	0.2	0.2	0.3	4.8
(n=91)	CI	93.8-96.2	2.0-4.3	0.5-1.3	0.2-0.6	0.1-0.4	0.1-0.5	0.2-0.5	3.8-6.2
8th	%	99.8	0.1	0.0	0.0	0.2	0.0	0.0	0.2
(n=1,320)	CI	99.3-99.9	0.0-0.5	-	-	0.0-0.6	-	-	0.1-0.7
10th	%	99.2	0.2	0.2	0.0	0.2	0.0	0.2	0.8
(n=512)	CI	97.9-99.7	0.0-1.4	0.0-1.4	-	0.0-1.4	-	0.0-1.4	0.3-2.1
12th	%	98.8	0.0	0.4	0.0	0.0	0.0	0.8	1.2
(n=247)	CI	96.3-99.6	-	0.1-2.8	-	-	-	0.2-3.2	0.4-3.7

Note: The confidence interval (CI) is 95 percent.

Table BH1.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	25.7	22.9	16.4	34.9	51.4
(n=742)	CI	21.9–29.9	19.3–27.1	13.3–20.2	30.8–39.3	46.8–55.9
Male	%	26.7	18.0	17.6	37.7	55.3
(n=339)	CI	21.1–33.3	13.2–24.0	13.0–23.4	31.6–44.2	48.5–61.9
Female	%	22.7	28.8	16.6	31.9	48.5
(n=376)	CI	18.0–28.3	23.3–35.1	12.1–22.2	26.3–38.1	42.1–54.9
White	%	–	–	–	–	–
(n=598)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=16)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=29)	CI	–	–	–	–	–
8th	%	21.5	18.5	12.9	47.0	60.0
(n=502)	CI	18.1–25.3	15.4–22.2	10.3–16.2	42.7–51.4	55.6–64.2
10th	%	32.9	26.7	18.5	21.9	40.4
(n=145)	CI	25.7–41.0	20.1–34.6	12.9–25.7	15.9–29.4	32.7–48.7
12th	%	24.5	24.5	18.4	32.7	51.0
(n=95)	CI	16.9–34.1	16.9–34.1	11.8–27.5	24.0–42.7	41.0–60.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	27.1	33.1	16.5	23.3	39.8
(n=1,058)	CI	23.8–30.7	29.7–36.7	13.9–19.4	20.4–26.5	36.2–43.4
Male	%	28.0	31.5	13.5	27.0	40.5
(n=481)	CI	23.1–33.5	26.6–36.9	10.2–17.6	22.6–32.0	35.3–45.9
Female	%	24.8	35.9	19.6	19.6	39.3
(n=542)	CI	20.5–29.7	31.1–41.1	15.8–24.1	16.0–23.9	34.3–44.4
White	%	–	–	–	–	–
(n=864)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=23)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=44)	CI	–	–	–	–	–
8th	%	19.1	30.4	17.5	33.0	50.5
(n=634)	CI	16.2–22.3	27.0–34.1	14.7–20.7	29.4–36.7	46.6–54.4
10th	%	27.9	36.2	16.2	19.6	35.8
(n=264)	CI	22.8–33.7	30.6–42.2	12.2–21.2	15.2–24.9	30.3–41.8
12th	%	32.5	32.5	16.0	19.0	35.0
(n=160)	CI	25.7–40.2	25.7–40.2	11.1–22.5	13.7–25.9	28.0–42.7

Note: The confidence interval (CI) is 95 percent.

Table BH1.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	41.6	26.5	11.9	20.1	32.0
(n=1,048)	CI	37.8–45.4	23.3–29.9	9.8–14.3	17.4–23.1	28.7–35.4
Male	%	42.4	26.2	11.3	20.1	31.4
(n=500)	CI	37.0–47.9	21.8–31.2	8.5–14.9	16.4–24.5	26.8–36.4
Female	%	39.6	27.6	13.3	19.5	32.8
(n=512)	CI	34.3–45.2	23.1–32.6	10.2–17.2	15.8–23.8	28.1–37.8
White	%	–	–	–	–	–
(n=857)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=24)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=38)	CI	–	–	–	–	–
8th	%	26.1	24.7	15.7	33.5	49.2
(n=632)	CI	22.8–29.7	21.5–28.2	13.0–18.7	30.0–37.3	45.3–53.1
10th	%	41.3	33.7	12.5	12.5	25.0
(n=263)	CI	35.5–47.4	28.2–39.7	9.0–17.1	9.0–17.1	20.1–30.6
12th	%	53.8	21.8	8.3	16.0	24.4
(n=153)	CI	45.9–61.6	15.9–29.0	4.9–13.9	11.0–22.7	18.2–31.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	30.1	23.4	12.9	33.6	46.5
(n=866)	CI	26.3–34.2	20.0–27.3	10.5–15.7	29.9–37.4	42.4–50.5
Male	%	28.2	22.3	13.7	35.8	49.5
(n=407)	CI	22.8–34.3	17.5–28.0	10.3–18.1	30.5–41.4	43.5–55.5
Female	%	30.9	24.8	12.1	32.1	44.3
(n=426)	CI	25.5–36.9	19.9–30.5	8.9–16.3	27.1–37.7	38.6–50.2
White	%	–	–	–	–	–
(n=688)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=22)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=41)	CI	–	–	–	–	–
8th	%	13.9	13.1	13.7	59.3	73.0
(n=518)	CI	11.2–17.2	10.5–16.3	11.0–17.0	55.0–63.4	69.0–76.6
10th	%	35.5	26.5	19.0	19.0	37.9
(n=210)	CI	29.3–42.3	21.0–33.0	14.2–24.9	14.2–24.9	31.6–44.7
12th	%	37.6	28.4	7.8	26.2	34.0
(n=138)	CI	29.9–46.0	21.5–36.5	4.3–13.6	19.6–34.2	26.6–42.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	1.8	98.2
(n=2,073)	CI	1.2–2.6	97.4–98.8
Male	%	1.4	98.6
(n=981)	CI	0.7–2.8	97.2–99.3
Female	%	1.9	98.1
(n=1,032)	CI	1.1–3.2	96.8–98.9
White	%	–	–
(n=1,675)	CI	–	–
Black	%	–	–
(n=54)	CI	–	–
Hispanic	%	–	–
(n=92)	CI	–	–
8th	%	1.1	98.9
(n=1,318)	CI	0.7–1.9	98.1–99.3
10th	%	2.5	97.5
(n=512)	CI	1.5–4.3	95.7–98.5
12th	%	1.6	98.4
(n=243)	CI	0.6–4.3	95.7–99.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	16.4	83.6
(n=2,060)	CI	14.3–18.7	81.3–85.7
Male	%	14.5	85.5
(n=973)	CI	11.7–17.7	82.3–88.3
Female	%	18.1	81.9
(n=1,026)	CI	15.2–21.5	78.5–84.8
White	%	–	–
(n=1,667)	CI	–	–
Black	%	–	–
(n=53)	CI	–	–
Hispanic	%	–	–
(n=90)	CI	–	–
8th	%	7.6	92.4
(n=1,306)	CI	6.3–9.1	90.9–93.7
10th	%	18.1	81.9
(n=512)	CI	15.0–21.6	78.4–85.0
12th	%	23.7	76.3
(n=242)	CI	18.7–29.4	70.6–81.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=32)	Male (n=10)	Female (n=18)	White (n=24)	Black (n=2)	Hispanic (n=1)	8th (n=15)	10th (n=13)	12th (n=4)
Got a prescription from one doctor	%	18.7	41.9	6.9	–	–	–	0.0	23.1	25.0
	CI	6.5–43.1	10.4–81.8	0.8–41.5	–	–	–	–	6.3–57.2	0.5–95.9
Got a prescription from more than one doctor	%	0.0	0.0	0.0	–	–	–	0.0	0.0	0.0
	CI	–	–	–	–	–	–	–	–	–
Bought on the Internet	%	5.2	14.5	0.0	–	–	–	6.7	7.7	0.0
	CI	1.0–22.3	2.0–57.9	–	–	–	–	0.7–41.6	0.8–46.9	–
Bought from a friend, relative, or someone you know	%	41.4	54.0	41.0	–	–	–	53.3	30.8	50.0
	CI	22.7–63.0	16.4–87.6	16.8–70.5	–	–	–	26.6–78.3	10.2–63.5	2.5–97.5
Got from a friend, relative, or someone you know for free	%	33.9	39.5	27.1	–	–	–	53.3	30.8	25.0
	CI	17.5–55.4	9.5–80.4	10.2–55.0	–	–	–	26.6–78.3	10.2–63.5	0.5–95.9
Took from a friend, relative, or someone you know without asking	%	30.5	21.0	33.0	–	–	–	20.0	23.1	50.0
	CI	14.0–54.2	2.0–77.4	11.4–65.4	–	–	–	5.6–51.2	6.3–57.2	2.5–97.5
Took from someone's house or workplace without asking	%	7.5	10.5	0.0	–	–	–	0.0	15.4	0.0
	CI	1.7–27.2	0.9–60.3	–	–	–	–	–	3.1–51.0	–
Got it at a party	%	28.2	43.6	9.6	–	–	–	26.7	30.8	25.0
	CI	13.1–50.5	11.4–82.2	1.7–39.0	–	–	–	9.0–57.1	10.2–63.5	0.5–95.9
Got it some other way	%	37.9	56.5	26.0	–	–	–	20.0	38.5	50.0
	CI	19.7–60.4	19.0–87.8	8.9–55.8	–	–	–	5.6–51.2	14.6–69.5	2.5–97.5

Note: The confidence interval (CI) is 95 percent.

Table BH1.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=248)	Male (n=99)	Female (n=140)	White (n=201)	Black (n=11)	Hispanic (n=10)	8th (n=99)	10th (n=92)	12th (n=57)
I bought it in a store	%	3.9	4.9	1.0	–	–	–	2.0	7.5	1.7
	CI	2.0–7.6	2.0–11.6	0.2–4.6	–	–	–	0.5–7.9	3.6–15.1	0.2–11.9
I bought it in a gas station	%	3.8	6.3	0.0	–	–	–	1.0	5.4	3.4
	CI	1.8–7.9	2.6–14.5	–	–	–	–	0.1–7.0	2.2–12.5	0.8–13.3
I bought it in a restaurant	%	2.8	2.0	1.5	–	–	–	0.0	1.1	5.2
	CI	1.0–7.5	0.3–13.1	0.2–9.8	–	–	–	–	0.1–7.5	1.6–15.3
I bought it in a bar or club	%	2.0	2.0	0.0	–	–	–	0.0	3.2	1.7
	CI	0.7–5.6	0.5–7.8	–	–	–	–	–	1.0–9.7	0.2–11.9
I bought it at a public event	%	1.9	1.4	0.3	–	–	–	2.0	2.2	1.7
	CI	0.7–5.3	0.3–6.2	0.0–2.0	–	–	–	0.5–7.9	0.5–8.4	0.2–11.9
I gave someone money to buy it for me	%	29.6	27.6	29.7	–	–	–	21.2	37.6	25.9
	CI	23.3–36.6	18.6–38.8	21.5–39.5	–	–	–	14.2–30.5	28.2–48.1	16.0–39.0
I got it from a sibling (under 21)	%	4.5	2.1	3.9	–	–	–	11.1	5.4	1.7
	CI	2.6–7.9	0.7–6.4	1.8–7.9	–	–	–	6.2–19.1	2.2–12.5	0.2–11.9
I got it from a friend (under 21)	%	19.5	14.3	21.3	–	–	–	29.3	31.2	6.9
	CI	14.8–25.2	8.5–23.0	15.0–29.4	–	–	–	21.1–39.2	22.5–41.5	2.5–17.4
I got it at a party	%	42.4	48.0	38.1	–	–	–	18.2	49.5	44.8
	CI	35.3–49.9	36.6–59.5	29.0–48.2	–	–	–	11.7–27.2	39.3–59.7	32.3–58.1
I took it from a store or family member	%	8.1	7.2	7.4	–	–	–	18.2	7.5	5.2
	CI	5.1–12.5	3.7–13.6	3.9–13.9	–	–	–	11.7–27.2	3.6–15.1	1.6–15.3
I took it home without permission	%	15.6	9.4	19.4	–	–	–	30.3	18.3	8.6
	CI	11.3–21.1	5.2–16.1	13.1–27.7	–	–	–	21.9–40.2	11.6–27.7	3.5–19.5
I got it from my parents with permission	%	17.2	15.5	16.3	–	–	–	27.3	15.1	15.5
	CI	12.4–23.2	9.3–24.7	10.3–24.7	–	–	–	19.3–37.0	9.0–24.0	8.1–27.6
I got it from an older sibling (over 21)	%	9.0	9.8	7.5	–	–	–	11.1	12.9	5.2
	CI	5.8–13.7	5.1–18.0	4.0–13.6	–	–	–	6.2–19.1	7.4–21.6	1.6–15.3
I got it from an older friend (over 21)	%	25.8	26.8	24.5	–	–	–	15.2	21.5	32.8
	CI	19.7–33.0	17.5–38.7	16.7–34.4	–	–	–	9.3–23.8	14.2–31.2	21.7–46.2
I got it from some other adult	%	16.9	15.3	16.9	–	–	–	20.2	23.7	10.3
	CI	12.3–22.7	8.8–25.2	11.2–24.7	–	–	–	13.3–29.4	16.0–33.5	4.6–21.6
I got it some other way	%	26.0	36.1	19.5	–	–	–	19.2	24.7	29.3
	CI	20.0–33.1	25.5–48.3	13.0–28.3	–	–	–	12.5–28.3	16.9–34.7	18.8–42.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?				
		I bought or tried to buy alcohol	Yes, asked for proof of age	No, not asked for proof of age
Total	%	2.2	19.1	80.9
(n=38)	CI	1.6–3.2	8.8–36.9	63.1–91.2
Male	%	3.7	17.1	82.9
(n=25)	CI	2.4–5.6	6.2–39.2	60.8–93.8
Female	%	0.9	14.1	85.9
(n=11)	CI	0.5–1.6	2.5–51.5	48.5–97.5
White	%	–	–	–
(n=32)	CI	–	–	–
Black	%	–	–	–
(n=0)	CI	–	–	–
Hispanic	%	–	–	–
(n=2)	CI	–	–	–
8th	%	1.9	22.2	77.8
(n=18)	CI	1.2–2.9	7.7–49.5	50.5–92.3
10th	%	3.9	22.2	77.8
(n=18)	CI	2.5–6.1	7.7–49.5	50.5–92.3
12th	%	0.9	0.0	100.0
(n=2)	CI	0.2–3.5	–	–

Note: The confidence interval (CI) is 95 percent.

Table BH1.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?					
		I bought or tried to buy alcohol	No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	2.2	47.5	8.3	44.2
(n=38)	CI	1.6–3.2	29.3–66.4	2.6–23.4	25.9–64.3
Male	%	3.7	45.0	7.5	47.5
(n=25)	CI	2.4–5.6	22.8–69.4	1.4–31.5	24.0–72.2
Female	%	0.9	65.5	6.2	28.4
(n=11)	CI	0.5–1.6	29.6–89.5	0.6–41.6	7.1–67.3
White	%	–	–	–	–
(n=32)	CI	–	–	–	–
Black	%	–	–	–	–
(n=0)	CI	–	–	–	–
Hispanic	%	–	–	–	–
(n=2)	CI	–	–	–	–
8th	%	1.9	66.7	14.3	19.0
(n=18)	CI	1.2–2.9	42.6–84.3	4.2–38.7	6.7–43.6
10th	%	3.9	50.0	7.1	42.9
(n=18)	CI	2.5–6.1	23.2–76.8	0.7–44.1	18.3–71.6
12th	%	0.9	0.0	0.0	100.0
(n=2)	CI	0.2–3.5	–	–	–

Note: The confidence interval (CI) is 95 percent.

Table BH1.39. Ever Ridden with a Prescription Drugs Impaired Driver

		Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		
		No	Yes	Not sure
Total	%	83.8	6.9	9.3
(n=1,954)	CI	81.8–85.6	5.6–8.3	8.0–10.9
Male	%	84.5	6.1	9.4
(n=909)	CI	81.4–87.2	4.4–8.4	7.4–11.9
Female	%	83.7	7.4	8.9
(n=988)	CI	80.9–86.1	5.8–9.6	7.1–11.0
White	%	–	–	–
(n=1,588)	CI	–	–	–
Black	%	–	–	–
(n=49)	CI	–	–	–
Hispanic	%	–	–	–
(n=83)	CI	–	–	–
8th	%	81.1	6.9	12.1
(n=1,236)	CI	78.8–83.2	5.6–8.4	10.4–14.0
10th	%	81.2	9.1	9.7
(n=482)	CI	77.5–84.5	6.8–12.0	7.4–12.7
12th	%	89.1	4.6	6.3
(n=236)	CI	84.5–92.5	2.6–8.2	3.8–10.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.40. Frequency Riding with a Prescription Drugs Impaired Driver

		During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?					
		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	95.6	1.0	1.7	0.5	1.1	4.4
(n=2,017)	CI	94.5–96.6	0.7–1.6	1.1–2.5	0.3–1.1	0.7–1.8	3.4–5.5
Male	%	95.4	1.1	2.2	0.6	0.8	4.6
(n=951)	CI	93.4–96.8	0.5–2.2	1.2–3.8	0.2–1.8	0.4–1.7	3.2–6.6
Female	%	96.1	1.1	1.3	0.5	1.0	3.9
(n=1,009)	CI	94.7–97.1	0.6–1.8	0.7–2.2	0.2–1.2	0.6–1.9	2.9–5.3
White	%	–	–	–	–	–	–
(n=1,635)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=51)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=880)	CI	–	–	–	–	–	–
8th	%	95.2	1.7	1.3	0.5	1.2	4.8
(n=1,277)	CI	93.9–96.3	1.1–2.6	0.8–2.1	0.3–1.1	0.7–1.9	3.7–6.1
10th	%	94.2	1.0	2.4	0.6	1.8	5.8
(n=501)	CI	91.8–96.0	0.4–2.4	1.4–4.2	0.2–1.8	0.9–3.4	4.0–8.2
12th	%	97.5	0.4	1.2	0.4	0.4	2.5
(n=239)	CI	94.6–98.9	0.1–2.9	0.4–3.8	0.1–2.9	0.1–2.9	1.1–5.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	73.4	18.8	7.8
(n=2,011)	CI	71.0–75.7	16.7–21.0	6.5–9.2
Male	%	74.1	17.3	8.6
(n=948)	CI	70.5–77.5	14.4–20.6	6.6–11.1
Female	%	73.7	19.5	6.8
(n=1,004)	CI	70.3–76.8	16.7–22.7	5.3–8.6
White	%	–	–	–
(n=1,630)	CI	–	–	–
Black	%	–	–	–
(n=49)	CI	–	–	–
Hispanic	%	–	–	–
(n=88)	CI	–	–	–
8th	%	75.5	14.6	9.8
(n=1,271)	CI	73.1–77.8	12.8–16.7	8.3–11.6
10th	%	69.2	22.2	8.5
(n=501)	CI	65.1–73.1	18.8–26.1	6.4–11.3
12th	%	75.6	19.4	5.0
(n=239)	CI	69.8–80.7	14.9–24.9	2.8–8.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	89.5	3.4	4.0	0.7	2.4	10.5
(n=2,020)	CI	87.8–91.1	2.6–4.5	3.1–5.2	0.4–1.4	1.7–3.3	8.9–12.2
Male	%	90.8	2.9	3.4	0.8	2.1	9.2
(n=953)	CI	88.2–92.9	1.8–4.4	2.2–5.3	0.3–2.4	1.3–3.5	7.1–11.8
Female	%	89.2	3.8	4.5	0.5	2.0	10.8
(n=1,009)	CI	86.8–91.3	2.6–5.4	3.2–6.3	0.2–1.3	1.2–3.2	8.7–13.2
White	%	–	–	–	–	–	–
(n=1,636)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=50)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=88)	CI	–	–	–	–	–	–
8th	%	91.2	4.0	2.7	0.5	1.6	8.8
(n=1,279)	CI	89.6–92.7	3.0–5.2	1.9–3.7	0.2–1.0	1.1–2.5	7.3–10.4
10th	%	86.2	3.4	5.9	0.8	3.8	13.8
(n=503)	CI	82.9–88.9	2.1–5.3	4.2–8.4	0.3–2.1	2.4–5.8	11.1–17.1
12th	%	91.3	2.9	3.3	0.8	1.7	8.7
(n=238)	CI	87.0–94.3	1.4–6.0	1.7–6.5	0.2–3.3	0.6–4.4	5.7–13.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.5	1.6	4.2	92.7	96.9
(n=2,001)	CI	1.0–2.4	1.0–2.4	3.3–5.4	91.2–94.0	95.8–97.7
Male	%	1.9	0.8	3.3	94.1	97.3
(n=946)	CI	1.0–3.4	0.3–2.3	2.2–4.9	91.8–95.7	95.5–98.4
Female	%	0.8	2.3	4.8	92.1	96.9
(n=999)	CI	0.4–1.8	1.4–3.5	3.5–6.7	89.9–93.8	95.5–97.9
White	%	–	–	–	–	–
(n=1,621)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	1.1	1.4	4.0	93.4	97.5
(n=1,267)	CI	0.7–1.9	0.9–2.2	3.1–5.3	91.9–94.7	96.4–98.2
10th	%	1.8	2.0	5.2	91.0	96.2
(n=496)	CI	0.9–3.4	1.1–3.7	3.6–7.6	88.1–93.2	94.1–97.6
12th	%	1.7	1.2	3.3	93.8	97.1
(n=238)	CI	0.6–4.4	0.4–3.8	1.7–6.5	89.9–96.2	94.0–98.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.3	4.3	9.9	83.5	93.4
(n=2,011)	CI	1.6–3.4	3.3–5.5	8.3–11.7	81.3–85.5	91.9–94.7
Male	%	2.5	4.7	8.9	83.9	92.8
(n=949)	CI	1.6–4.1	3.2–6.9	6.8–11.7	80.6–86.7	90.3–94.7
Female	%	2.0	4.1	10.3	83.6	93.9
(n=1,006)	CI	1.1–3.8	2.9–5.8	8.1–13.0	80.4–86.3	91.7–95.5
White	%	–	–	–	–	–
(n=1,629)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=88)	CI	–	–	–	–	–
8th	%	1.5	3.5	7.6	87.5	95.1
(n=1,275)	CI	1.0–2.3	2.6–4.6	6.3–9.2	85.5–89.2	93.7–96.1
10th	%	2.6	5.2	8.8	83.4	92.2
(n=498)	CI	1.5–4.4	3.6–7.5	6.6–11.6	79.9–86.4	89.5–94.3
12th	%	2.9	4.1	13.3	79.7	92.9
(n=238)	CI	1.4–6.0	2.2–7.6	9.5–18.2	74.1–84.3	88.9–95.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.45. Parents Feel Regularly Drinking Alcohol is Wrong

How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.1	7.9	10.3	75.6	86.0
(n=2,005)	CI	4.9–7.7	6.5–9.6	8.8–12.1	73.1–77.9	83.8–87.9
Male	%	6.9	6.9	8.3	77.9	86.2
(n=948)	CI	5.0–9.5	5.1–9.3	6.4–10.6	74.4–81.1	83.0–88.8
Female	%	5.5	8.4	12.4	73.7	86.1
(n=1,002)	CI	3.9–7.7	6.4–11.0	10.0–15.2	70.0–77.0	83.0–88.7
White	%	–	–	–	–	–
(n=1,625)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	3.5	5.3	9.2	82.0	91.2
(n=1,271)	CI	2.7–4.7	4.2–6.6	7.7–10.9	79.8–84.0	89.5–92.6
10th	%	7.0	8.0	11.8	73.2	85.0
(n=497)	CI	5.1–9.6	5.9–10.7	9.2–14.9	69.1–76.9	81.6–87.9
12th	%	7.9	10.4	10.0	71.7	81.7
(n=237)	CI	5.1–12.1	7.1–15.0	6.8–14.5	65.6–77.0	76.2–86.1

Note: The confidence interval (CI) is 95 percent.

Table BH1.46. Parents Feel Smoking Tobacco is Wrong

How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.3	4.3	9.0	83.4	92.4
(n=2,012)	CI	2.4–4.5	3.2–5.6	7.6–10.7	81.2–85.4	90.8–93.8
Male	%	4.0	4.7	8.9	82.4	91.3
(n=950)	CI	2.7–6.1	3.1–7.0	6.8–11.5	79.0–85.4	88.5–93.5
Female	%	2.5	3.8	8.5	85.2	93.7
(n=1,006)	CI	1.5–4.3	2.6–5.4	6.6–10.9	82.2–87.7	91.5–95.3
White	%	–	–	–	–	–
(n=1,631)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	2.0	2.8	7.8	87.5	95.2
(n=1,275)	CI	1.3–2.9	2.0–3.9	6.4–9.4	85.5–89.2	93.9–96.3
10th	%	3.8	4.6	9.4	82.3	91.6
(n=499)	CI	2.4–5.9	3.1–6.8	7.1–12.3	78.7–85.4	88.9–93.8
12th	%	4.1	5.4	10.0	80.5	90.5
(n=238)	CI	2.2–7.6	3.1–9.1	6.7–14.5	75.0–85.1	86.0–93.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.8	2.6	6.6	89.0	95.6
(n=2,008)	CI	1.2–2.8	1.8–3.7	5.2–8.2	87.0–90.7	94.2–96.6
Male	%	2.2	2.6	6.4	88.8	95.2
(n=948)	CI	1.3–3.8	1.4–4.6	4.5–9.0	85.7–91.3	92.9–96.8
Female	%	1.4	2.6	6.1	90.0	96.0
(n=1,004)	CI	0.6–2.9	1.7–4.1	4.4–8.3	87.3–92.1	94.2–97.3
White	%	–	–	–	–	–
(n=1,628)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	0.8	1.9	4.4	92.9	97.3
(n=1,273)	CI	0.4–1.5	1.3–2.8	3.4–5.7	91.4–94.2	96.3–98.1
10th	%	2.6	2.6	4.6	90.2	94.8
(n=497)	CI	1.5–4.4	1.5–4.4	3.1–6.8	87.3–92.5	92.5–96.4
12th	%	2.1	3.3	10.8	83.8	94.6
(n=238)	CI	0.9–4.9	1.7–6.5	7.4–15.4	78.6–88.0	90.9–96.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.48. Parents Feel Using Electronic Cigarettes is Wrong

How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.1	7.5	10.9	75.5	86.4
(n=2,006)	CI	4.9–7.6	6.2–9.1	9.3–12.7	73.1–77.8	84.3–88.1
Male	%	6.9	7.8	9.9	75.3	85.2
(n=945)	CI	5.1–9.4	5.9–10.3	7.8–12.5	71.7–78.7	82.1–87.9
Female	%	5.5	6.9	11.4	76.2	87.6
(n=1,005)	CI	3.9–7.7	5.3–9.0	9.2–14.0	72.8–79.3	84.8–89.9
White	%	–	–	–	–	–
(n=1,626)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	4.1	6.7	10.1	79.1	89.2
(n=1,271)	CI	3.1–5.3	5.4–8.2	8.6–11.9	76.7–81.2	87.4–90.8
10th	%	6.8	9.2	12.4	71.7	84.0
(n=498)	CI	4.9–9.4	6.9–12.1	9.8–15.6	67.5–75.4	80.5–87.0
12th	%	7.5	6.7	10.0	75.8	85.8
(n=237)	CI	4.8–11.6	4.1–10.6	6.8–14.5	70.0–80.9	80.8–89.7

Note: The confidence interval (CI) is 95 percent.

Table BH1.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.3	5.0	7.0	82.7	89.7
(n=2,010)	CI	4.1–6.7	3.9–6.4	5.7–8.6	80.4–84.7	87.8–91.3
Male	%	5.5	4.0	5.3	85.2	90.5
(n=949)	CI	3.9–7.8	2.6–6.1	3.8–7.4	82.0–87.9	87.7–92.7
Female	%	4.5	5.6	8.5	81.3	89.9
(n=1,005)	CI	3.1–6.5	4.1–7.7	6.5–11.1	78.0–84.3	87.2–92.0
White	%	–	–	–	–	–
(n=1,630)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	2.6	2.8	5.7	88.9	94.6
(n=1,273)	CI	1.8–3.6	2.0–3.9	4.5–7.1	87.1–90.5	93.2–95.7
10th	%	6.6	6.8	6.8	79.9	86.7
(n=499)	CI	4.7–9.1	4.9–9.3	4.9–9.3	76.1–83.2	83.4–89.4
12th	%	6.6	5.4	8.7	79.3	88.0
(n=238)	CI	4.1–10.6	3.1–9.1	5.7–13.0	73.6–83.9	83.2–91.5

Note: The confidence interval (CI) is 95 percent.

Table BH1.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.5	4.1	6.2	85.1	91.3
(n=2,003)	CI	3.5–5.9	3.1–5.5	5.0–7.7	83.0–87.0	89.6–92.8
Male	%	4.7	4.5	3.6	87.1	90.7
(n=950)	CI	3.2–6.9	3.0–6.7	2.4–5.3	84.1–89.6	87.9–92.9
Female	%	4.0	3.6	8.3	84.0	92.3
(n=997)	CI	2.7–6.0	2.5–5.4	6.3–10.9	80.8–86.7	89.9–94.2
White	%	–	–	–	–	–
(n=1,625)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=50)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	2.4	2.1	5.2	90.4	95.6
(n=1,268)	CI	1.7–3.4	1.4–3.0	4.1–6.6	88.6–91.9	94.3–96.6
10th	%	5.8	5.8	6.0	82.4	88.4
(n=498)	CI	4.0–8.2	4.0–8.2	4.2–8.4	78.8–85.5	85.3–91.0
12th	%	5.4	4.6	7.5	82.5	90.0
(n=237)	CI	3.2–9.1	2.5–8.1	4.8–11.6	77.1–86.8	85.5–93.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.51. Friends Feel Misusing Prescription Drugs is Wrong

How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	8.4	5.8	12.8	73.0	85.8
(n=2,009)	CI	7.0–10.1	4.7–7.1	11.1–14.8	70.5–75.3	83.8–87.6
Male	%	10.5	6.6	12.8	70.2	82.9
(n=948)	CI	8.3–13.3	4.8–8.8	10.3–15.7	66.4–73.7	79.7–85.7
Female	%	5.9	5.0	13.0	76.1	89.1
(n=1,005)	CI	4.3–7.9	3.8–6.6	10.6–15.8	72.8–79.2	86.7–91.1
White	%	–	–	–	–	–
(n=1,630)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	6.0	6.2	12.9	74.8	87.8
(n=1,276)	CI	4.9–7.5	5.0–7.7	11.2–14.9	72.4–77.2	85.9–89.5
10th	%	11.6	7.4	11.8	69.1	81.0
(n=496)	CI	9.1–14.8	5.4–10.1	9.3–15.0	64.9–73.1	77.3–84.2
12th	%	7.5	3.8	13.8	75.0	88.8
(n=237)	CI	4.8–11.6	2.0–7.1	9.9–18.8	69.1–80.1	84.1–92.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.52. Friends Feel Drinking Alcohol is Wrong

How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	17.4	14.1	16.4	52.0	68.4
(n=2,010)	CI	15.4–19.7	12.3–16.1	14.5–18.5	49.3–54.7	65.8–71.0
Male	%	18.8	14.6	16.1	50.5	66.6
(n=949)	CI	15.8–22.3	11.9–17.7	13.4–19.2	46.6–54.4	62.7–70.3
Female	%	15.4	14.0	17.3	53.3	70.6
(n=1,005)	CI	12.7–18.5	11.6–16.8	14.6–20.4	49.5–57.1	67.0–74.1
White	%	–	–	–	–	–
(n=1,631)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	9.5	12.9	17.0	60.7	77.7
(n=1,276)	CI	8.0–11.2	11.1–14.8	15.0–19.2	57.9–63.3	75.3–79.9
10th	%	22.4	15.4	16.0	46.1	62.1
(n=496)	CI	19.0–26.3	12.5–18.9	13.1–19.5	41.7–50.5	57.8–66.3
12th	%	20.3	14.1	16.2	49.4	65.6
(n=238)	CI	15.7–25.9	10.2–19.1	12.0–21.4	43.1–55.7	59.3–71.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	16.0	11.9	16.9	55.1	72.0
(n=2,008)	CI	14.0–18.2	10.3–13.8	15.0–19.0	52.4–57.8	69.5–74.5
Male	%	18.9	12.6	15.9	52.6	68.6
(n=949)	CI	15.8–22.3	10.0–15.7	13.3–19.0	48.7–56.5	64.6–72.2
Female	%	12.5	11.6	18.6	57.3	75.9
(n=1,004)	CI	10.1–15.4	9.5–14.2	15.7–21.7	53.5–61.1	72.4–79.0
White	%	–	–	–	–	–
(n=1,630)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=86)	CI	–	–	–	–	–
8th	%	9.5	9.7	17.5	63.3	80.8
(n=1,276)	CI	8.0–11.2	8.2–11.5	15.5–19.7	60.6–65.9	78.5–82.9
10th	%	20.3	14.5	16.7	48.5	65.2
(n=494)	CI	17.0–24.1	11.6–17.9	13.7–20.3	44.1–52.9	60.9–69.3
12th	%	18.3	11.6	16.6	53.5	70.1
(n=238)	CI	13.8–23.7	8.1–16.4	12.4–21.9	47.2–59.8	64.0–75.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	27.6	13.7	12.5	46.3	58.7
(n=2,001)	CI	25.2–30.2	11.9–15.6	10.8–14.3	43.6–49.0	56.0–61.4
Male	%	29.8	13.0	12.2	45.0	57.2
(n=945)	CI	26.2–33.7	10.6–15.8	9.9–14.9	41.1–48.9	53.2–61.1
Female	%	25.1	14.4	13.3	47.2	60.5
(n=1,000)	CI	21.9–28.7	11.9–17.3	10.9–16.1	43.3–51.0	56.6–64.2
White	%	–	–	–	–	–
(n=1,623)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=87)	CI	–	–	–	–	–
8th	%	17.6	13.4	14.8	54.3	69.0
(n=1,273)	CI	15.6–19.8	11.6–15.3	12.9–16.8	51.5–57.0	66.5–71.5
10th	%	34.0	15.6	11.7	38.7	50.4
(n=491)	CI	29.9–38.3	12.6–19.1	9.2–14.9	34.5–43.1	46.0–54.8
12th	%	31.3	12.1	10.8	45.8	56.7
(n=237)	CI	25.7–37.4	8.5–16.9	7.5–15.5	39.6–52.2	50.3–62.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	21.1	12.9	14.1	51.9	66.0
(n=2,004)	CI	18.9–23.6	11.1–14.9	12.3–16.1	49.2–54.6	63.3–68.6
Male	%	21.9	13.1	14.1	51.0	65.0
(n=946)	CI	18.6–25.5	10.6–16.2	11.5–17.1	47.0–54.9	61.0–68.8
Female	%	19.6	13.0	14.5	52.9	67.4
(n=1,003)	CI	16.6–22.9	10.6–15.9	12.0–17.4	49.0–56.7	63.6–71.0
White	%	–	–	–	–	–
(n=1,627)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=86)	CI	–	–	–	–	–
8th	%	11.6	9.0	14.0	65.4	79.4
(n=1,271)	CI	9.9–13.4	7.6–10.8	12.2–16.0	62.7–68.0	77.1–81.5
10th	%	26.1	16.3	13.7	44.0	57.6
(n=495)	CI	22.4–30.2	13.3–19.8	10.9–17.0	39.7–48.4	53.2–61.9
12th	%	25.7	13.3	14.5	46.5	61.0
(n=238)	CI	20.6–31.7	9.5–18.2	10.6–19.6	40.2–52.8	54.6–67.0

Note: The confidence interval (CI) is 95 percent.

Table BH1.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	87.8	5.4	1.6	0.8	4.3	12.2
(n=2,003)	CI	85.9–89.5	4.3–6.8	1.1–2.3	0.5–1.5	3.3–5.7	10.5–14.1
Male	%	86.6	6.4	1.3	0.7	5.0	13.4
(n=943)	CI	83.5–89.3	4.6–8.8	0.6–2.6	0.2–2.3	3.4–7.3	10.7–16.5
Female	%	90.0	4.7	1.9	0.9	2.5	10.0
(n=1,002)	CI	87.7–92.0	3.4–6.6	1.2–3.0	0.4–1.8	1.6–3.9	8.0–12.3
White	%	–	–	–	–	–	–
(n=1,622)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=50)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=88)	CI	–	–	–	–	–	–
8th	%	90.8	4.3	2.3	0.5	2.2	9.2
(n=1,265)	CI	89.0–92.2	3.3–5.5	1.6–3.3	0.2–1.1	1.5–3.2	7.8–11.0
10th	%	85.3	6.6	1.6	1.2	5.4	14.7
(n=500)	CI	81.9–88.1	4.7–9.1	0.8–3.2	0.5–2.6	3.7–7.7	11.9–18.1
12th	%	87.6	5.4	0.8	0.8	5.4	12.4
(n=238)	CI	82.7–91.2	3.1–9.1	0.2–3.3	0.2–3.3	3.1–9.1	8.8–17.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	54.1	10.9	9.4	5.5	20.0	45.9
(n=2,030)	CI	51.4–56.8	9.5–12.6	7.9–11.2	4.4–7.0	17.8–22.4	43.2–48.6
Male	%	57.6	9.7	8.2	4.7	19.9	42.4
(n=955)	CI	53.6–61.5	7.7–12.2	6.2–10.7	3.2–6.8	16.7–23.6	38.5–46.4
Female	%	51.6	12.0	10.6	6.7	19.2	48.4
(n=1,016)	CI	47.8–55.4	9.9–14.5	8.3–13.3	4.9–9.0	16.2–22.5	44.6–52.2
White	%	–	–	–	–	–	–
(n=1,645)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=51)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=87)	CI	–	–	–	–	–	–
8th	%	66.6	12.3	8.2	3.1	9.9	33.4
(n=1,286)	CI	63.9–69.1	10.6–14.2	6.8–9.8	2.3–4.2	8.4–11.6	30.9–36.1
10th	%	48.7	14.6	7.9	6.9	21.9	51.3
(n=504)	CI	44.4–53.1	11.8–18.0	5.8–10.6	5.0–9.5	18.5–25.7	46.9–55.6
12th	%	46.9	5.8	12.3	6.6	28.4	53.1
(n=240)	CI	40.7–53.2	3.4–9.5	8.7–17.1	4.1–10.5	23.0–34.4	46.8–59.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	67.1	8.2	7.0	3.9	13.8	32.9
(n=2,015)	CI	64.4–69.7	6.9–9.8	5.7–8.7	3.0–5.2	11.8–16.0	30.3–35.6
Male	%	68.9	6.9	5.6	4.1	14.4	31.1
(n=948)	CI	64.9–72.7	5.2–9.3	3.9–8.1	2.7–6.2	11.5–17.8	27.3–35.1
Female	%	66.1	9.4	8.4	4.0	12.1	33.9
(n=1,009)	CI	62.3–69.7	7.4–11.9	6.4–11.0	2.7–5.8	9.6–15.1	30.3–37.7
White	%	–	–	–	–	–	–
(n=1,630)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=51)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=88)	CI	–	–	–	–	–	–
8th	%	82.9	6.9	3.8	1.9	4.5	17.1
(n=1,273)	CI	80.7–84.8	5.6–8.4	2.9–5.1	1.3–2.8	3.5–5.8	15.2–19.3
10th	%	61.2	10.3	7.7	5.7	15.0	38.8
(n=502)	CI	56.8–65.4	7.9–13.3	5.7–10.4	4.0–8.2	12.2–18.5	34.6–43.2
12th	%	57.2	7.4	9.5	4.1	21.8	42.8
(n=240)	CI	50.9–63.3	4.7–11.5	6.4–13.9	2.2–7.5	17.0–27.5	36.7–49.1

Note: The confidence interval (CI) is 95 percent.

Table BH1.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>smoked part, or all of a cigarette?</u>							
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	69.5	9.6	6.6	3.1	11.2	30.5
(n=2,015)	CI	66.9–71.9	8.1–11.2	5.4–8.2	2.3–4.2	9.6–13.1	28.1–33.1
Male	%	69.1	8.9	7.2	2.2	12.6	30.9
(n=949)	CI	65.2–72.7	6.9–11.3	5.3–9.8	1.4–3.5	10.0–15.6	27.3–34.8
Female	%	70.8	10.3	6.3	4.0	8.7	29.2
(n=1,007)	CI	67.2–74.1	8.3–12.8	4.7–8.3	2.7–5.9	6.8–11.0	25.9–32.8
White	%	–	–	–	–	–	–
(n=1,631)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=50)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=87)	CI	–	–	–	–	–	–
8th	%	75.5	10.3	4.2	2.7	7.4	24.5
(n=1,274)	CI	73.1–77.8	8.7–12.1	3.2–5.4	1.9–3.7	6.1–9.0	22.2–26.9
10th	%	61.9	11.3	8.7	4.2	13.9	38.1
(n=501)	CI	57.6–66.1	8.8–14.4	6.6–11.5	2.7–6.3	11.1–17.2	33.9–42.4
12th	%	71.2	7.0	7.0	2.5	12.3	28.8
(n=240)	CI	65.1–76.6	4.4–11.0	4.4–11.0	1.1–5.4	8.7–17.1	23.4–34.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used an e-cigarette?</u>							
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	56.5	9.5	6.5	5.5	21.9	43.5
(n=2,013)	CI	53.8–59.2	8.1–11.1	5.3–8.0	4.4–7.0	19.7–24.3	40.8–46.2
Male	%	55.7	9.0	5.1	5.2	25.0	44.3
(n=947)	CI	51.8–59.6	7.1–11.4	3.6–7.0	3.7–7.3	21.6–28.8	40.4–48.2
Female	%	57.3	10.4	8.1	6.1	18.1	42.7
(n=1,007)	CI	53.4–61.1	8.3–12.9	6.2–10.6	4.4–8.4	15.3–21.3	38.9–46.6
White	%	–	–	–	–	–	–
(n=1,627)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=51)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=88)	CI	–	–	–	–	–	–
8th	%	66.4	10.1	5.8	3.2	14.5	33.6
(n=1,273)	CI	63.7–68.9	8.5–11.8	4.7–7.2	2.4–4.3	12.7–16.6	31.1–36.3
10th	%	47.2	10.9	7.5	7.5	26.8	52.8
(n=501)	CI	42.9–51.6	8.5–14.0	5.5–10.2	5.5–10.2	23.1–30.8	48.4–57.1
12th	%	56.2	7.4	6.2	5.8	24.4	43.8
(n=239)	CI	49.8–62.4	4.7–11.5	3.8–10.1	3.4–9.6	19.3–30.2	37.6–50.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	66.2	9.0	5.6	3.2	16.0	33.8
(n=2,023)	CI	63.5–68.7	7.6–10.7	4.5–7.1	2.3–4.4	14.0–18.3	31.3–36.5
Male	%	67.1	7.6	5.3	3.1	16.9	32.9
(n=955)	CI	63.2–70.8	5.8–9.9	3.7–7.5	1.9–5.0	13.9–20.3	29.2–36.8
Female	%	65.8	10.7	6.0	3.4	14.2	34.2
(n=1,009)	CI	62.0–69.4	8.6–13.2	4.4–8.1	2.2–5.3	11.6–17.2	30.6–38.0
White	%	–	–	–	–	–	–
(n=1,636)	CI	–	–	–	–	–	–
Black	%	–	–	–	–	–	–
(n=55)	CI	–	–	–	–	–	–
Hispanic	%	–	–	–	–	–	–
(n=88)	CI	–	–	–	–	–	–
8th	%	79.0	7.7	4.0	2.5	6.9	21.0
(n=1,281)	CI	76.7–81.1	6.3–9.2	3.0–5.2	1.8–3.5	5.6–8.4	18.9–23.3
10th	%	58.6	11.9	6.3	2.6	20.6	41.4
(n=502)	CI	54.2–62.8	9.3–15.0	4.5–8.8	1.5–4.4	17.3–24.4	37.2–45.8
12th	%	60.9	7.4	6.6	4.5	20.6	39.1
(n=240)	CI	54.6–66.9	4.7–11.5	4.1–10.5	2.5–8.0	15.9–26.2	33.1–45.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	8.6	4.6	11.8	74.9	86.8
(n=1,937)	CI	7.2–10.3	3.7–5.7	10.2–13.7	72.5–77.2	84.9–88.5
Male	%	11.5	3.8	13.3	71.4	84.7
(n=931)	CI	9.1–14.4	2.7–5.4	10.8–16.3	67.6–74.9	81.6–87.3
Female	%	5.7	5.3	10.4	78.5	88.9
(n=989)	CI	4.3–7.7	4.0–7.1	8.3–13.0	75.3–81.4	86.5–91.0
White	%	–	–	–	–	–
(n=1,605)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=47)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=84)	CI	–	–	–	–	–
8th	%	7.1	6.4	12.6	73.8	86.5
(n=1,250)	CI	5.8–8.7	5.2–7.9	10.9–14.6	71.3–76.2	84.5–88.3
10th	%	10.7	5.2	9.9	74.2	84.1
(n=493)	CI	8.2–13.7	3.6–7.6	7.5–12.8	70.1–77.9	80.6–87.0
12th	%	8.0	2.1	13.1	76.8	89.9
(n=234)	CI	5.2–12.3	0.9–5.0	9.3–18.0	70.9–81.8	85.3–93.1

Note: The confidence interval (CI) is 95 percent.

Table BH1.63. Perceived Risk of Regularly Drinking Alcohol

		How much do you think people risk harming themselves physically and in other ways when they <u>take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	12.5	15.4	23.2	48.8	72.0
(n=1,993)	CI	10.8–14.5	13.6–17.5	21.0–25.5	46.1–51.5	69.5–74.4
Male	%	17.4	14.2	25.4	43.0	68.4
(n=935)	CI	14.5–20.7	11.7–17.2	22.0–29.0	39.2–46.9	64.5–72.0
Female	%	8.1	16.2	21.6	54.1	75.7
(n=1,000)	CI	6.3–10.2	13.5–19.3	18.7–24.9	50.2–57.9	72.3–78.9
White	%	–	–	–	–	–
(n=1,618)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=48)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	10.0	15.6	24.1	50.2	74.4
(n=1,260)	CI	8.5–11.8	13.7–17.7	21.8–26.6	47.5–53.0	71.9–76.7
10th	%	15.8	14.8	23.2	46.3	69.5
(n=498)	CI	12.8–19.2	11.9–18.2	19.7–27.1	42.0–50.7	65.3–73.4
12th	%	11.8	16.0	22.3	50.0	72.3
(n=235)	CI	8.2–16.6	11.8–21.2	17.4–28.0	43.6–56.4	66.2–77.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.64. Perceived Risk of Binge Drinking

		How much do you think people risk harming themselves physically and in other ways when they <u>take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.9	11.2	19.3	58.7	78.0
(n=1,992)	CI	9.2–12.7	9.5–13.1	17.3–21.5	56.0–61.3	75.6–80.2
Male	%	14.9	11.0	18.7	55.4	74.1
(n=934)	CI	12.2–18.1	8.7–13.9	15.9–22.0	51.4–59.3	70.3–77.5
Female	%	7.0	11.4	20.0	61.6	81.6
(n=1,000)	CI	5.3–9.1	9.1–14.2	17.1–23.2	57.8–65.3	78.4–84.5
White	%	–	–	–	–	–
(n=1,618)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=48)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	8.2	9.0	20.0	62.8	82.8
(n=1,259)	CI	6.8–9.8	7.5–10.7	17.9–22.3	60.1–65.5	80.7–84.8
10th	%	12.6	11.2	21.0	55.3	76.2
(n=498)	CI	9.9–15.8	8.7–14.3	17.6–24.8	50.9–59.6	72.3–79.8
12th	%	11.8	13.4	16.8	58.0	74.8
(n=235)	CI	8.2–16.6	9.6–18.4	12.5–22.1	51.6–64.1	68.8–79.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	9.9	6.5	15.0	68.5	83.5
(n=1,993)	CI	8.4–11.7	5.4–8.0	13.3–17.0	65.9–70.9	81.4–85.4
Male	%	13.8	7.2	14.9	64.2	79.1
(n=934)	CI	11.1–16.9	5.4–9.4	12.4–17.8	60.3–67.9	75.6–82.2
Female	%	6.3	6.1	15.0	72.6	87.6
(n=1,001)	CI	4.7–8.3	4.6–8.2	12.5–17.8	69.2–75.8	84.9–89.8
White	%	–	–	–	–	–
(n=1,618)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=48)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	8.1	7.1	17.6	67.2	84.8
(n=1,262)	CI	6.7–9.7	5.8–8.7	15.6–19.8	64.6–69.7	82.7–86.7
10th	%	10.8	7.4	16.4	65.3	81.8
(n=496)	CI	8.4–13.9	5.4–10.1	13.4–20.0	61.0–69.4	78.1–84.9
12th	%	10.9	5.0	10.9	73.1	84.0
(n=235)	CI	7.5–15.6	2.9–8.7	7.5–15.6	67.1–78.4	78.8–88.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	14.6	19.0	25.1	41.4	66.4
(n=1,987)	CI	12.7–16.7	17.0–21.1	22.8–27.5	38.7–44.1	63.8–68.9
Male	%	20.3	20.1	23.8	35.8	59.6
(n=932)	CI	17.2–23.9	17.2–23.4	20.6–27.2	32.1–39.6	55.6–63.4
Female	%	9.1	18.1	26.1	46.7	72.8
(n=997)	CI	7.2–11.5	15.4–21.2	22.8–29.6	42.9–50.6	69.3–76.0
White	%	–	–	–	–	–
(n=1,612)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=48)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	11.9	18.8	26.6	42.7	69.3
(n=1,257)	CI	10.3–13.8	16.7–21.0	24.2–29.1	40.0–45.5	66.7–71.8
10th	%	15.9	23.7	24.3	36.1	60.4
(n=495)	CI	12.9–19.4	20.2–27.6	20.7–28.3	32.0–40.5	56.1–64.7
12th	%	16.0	14.3	24.4	45.4	69.7
(n=235)	CI	11.8–21.2	10.4–19.4	19.3–30.3	39.1–51.8	63.6–75.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	27.5	22.2	14.7	35.7	50.4
(n=1,982)	CI	25.0–30.0	20.0–24.5	12.9–16.6	33.2–38.3	47.7–53.1
Male	%	30.8	23.4	13.2	32.7	45.9
(n=931)	CI	27.1–34.7	20.1–26.9	10.8–16.0	29.2–36.4	42.0–49.8
Female	%	24.1	20.9	16.4	38.6	55.0
(n=993)	CI	20.9–27.6	17.9–24.3	13.8–19.3	34.9–42.4	51.1–58.8
White	%	–	–	–	–	–
(n=1,609)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=47)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	19.4	20.9	17.4	42.4	59.7
(n=1,256)	CI	17.3–21.7	18.7–23.2	15.4–19.6	39.6–45.1	57.0–62.4
10th	%	30.5	22.0	14.7	32.7	47.5
(n=492)	CI	26.6–34.7	18.6–25.9	11.9–18.2	28.7–37.0	43.1–51.9
12th	%	32.5	23.6	11.8	32.1	43.9
(n=234)	CI	26.8–38.8	18.6–29.5	8.3–16.6	26.4–38.3	37.7–50.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	21.4	16.0	18.1	44.5	62.6
(n=1,984)	CI	19.2–23.8	14.0–18.1	16.1–20.3	41.8–47.2	59.9–65.3
Male	%	25.0	15.3	19.2	40.5	59.7
(n=930)	CI	21.6–28.8	12.6–18.5	16.3–22.5	36.7–44.3	55.6–63.6
Female	%	18.0	16.0	17.2	48.8	66.0
(n=996)	CI	15.2–21.2	13.3–19.1	14.5–20.3	45.0–52.7	62.2–69.6
White	%	–	–	–	–	–
(n=1,610)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=47)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=84)	CI	–	–	–	–	–
8th	%	14.8	13.0	19.0	53.1	72.2
(n=1,257)	CI	12.9–16.9	11.3–15.0	16.9–21.3	50.4–55.9	69.6–74.6
10th	%	25.4	16.3	17.9	40.3	58.3
(n=493)	CI	21.8–29.4	13.3–19.9	14.8–21.6	36.1–44.7	53.9–62.5
12th	%	24.1	18.6	17.3	40.1	57.4
(n=234)	CI	19.0–29.9	14.1–24.1	13.0–22.7	34.0–46.5	51.0–63.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.5	3.8	10.2	82.4	92.6
(n=2,001)	CI	2.6–4.8	2.9–5.0	8.6–12.0	80.2–84.4	91.0–93.9
Male	%	4.5	4.1	9.2	82.2	91.4
(n=941)	CI	3.0–6.7	2.7–6.1	7.1–11.9	78.8–85.2	88.7–93.5
Female	%	2.0	3.8	10.9	83.3	94.2
(n=1,004)	CI	1.2–3.4	2.6–5.3	8.7–13.5	80.4–85.9	92.3–95.7
White	%	–	–	–	–	–
(n=1,625)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	1.4	4.3	8.6	85.7	94.3
(n=1,270)	CI	0.9–2.2	3.3–5.6	7.2–10.3	83.6–87.5	92.8–95.4
10th	%	5.0	3.8	11.4	79.8	91.2
(n=498)	CI	3.4–7.3	2.4–5.9	8.9–14.5	76.1–83.1	88.4–93.4
12th	%	4.2	3.4	10.6	81.8	92.4
(n=233)	CI	2.3–7.7	1.7–6.7	7.2–15.2	76.3–86.2	88.2–95.2

Note: The confidence interval (CI) is 95 percent.

Table BH1.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.9	10.5	17.4	65.3	82.6
(n=2,002)	CI	5.5–8.6	8.9–12.3	15.4–19.5	62.6–67.9	80.4–84.7
Male	%	8.5	10.0	16.8	64.7	81.5
(n=942)	CI	6.3–11.4	7.8–12.6	13.9–20.2	60.7–68.5	78.0–84.5
Female	%	4.1	10.8	18.4	66.7	85.1
(n=1,003)	CI	2.8–6.0	8.6–13.6	15.7–21.5	62.9–70.2	82.0–87.7
White	%	–	–	–	–	–
(n=1,626)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=84)	CI	–	–	–	–	–
8th	%	2.4	8.9	14.3	74.4	88.7
(n=1,272)	CI	1.7–3.4	7.4–10.6	12.5–16.3	71.9–76.7	86.8–90.3
10th	%	8.2	11.2	20.1	60.4	80.5
(n=495)	CI	6.1–11.0	8.7–14.3	16.8–23.8	56.1–64.7	76.8–83.8
12th	%	10.1	11.3	17.6	60.9	78.6
(n=235)	CI	6.8–14.6	7.9–16.1	13.3–23.1	54.5–67.0	72.9–83.4

Note: The confidence interval (CI) is 95 percent.

Table BH1.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	8.6	10.9	17.1	63.4	80.4
(n=2,003)	CI	7.1–10.5	9.3–12.8	15.1–19.2	60.6–66.0	78.0–82.6
Male	%	10.1	11.3	16.9	61.8	78.6
(n=943)	CI	7.7–13.2	8.9–14.2	14.0–20.2	57.7–65.6	74.9–81.9
Female	%	6.0	10.8	17.6	65.5	83.2
(n=1,003)	CI	4.3–8.5	8.6–13.4	14.9–20.7	61.8–69.1	79.9–85.9
White	%	–	–	–	–	–
(n=1,626)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	2.6	7.8	15.2	74.4	89.6
(n=1,271)	CI	1.9–3.6	6.4–9.4	13.3–17.3	72.0–76.8	87.8–91.2
10th	%	8.8	13.2	19.6	58.4	78.0
(n=497)	CI	6.6–11.6	10.5–16.5	16.3–23.3	54.0–62.7	74.1–81.4
12th	%	14.7	11.8	16.4	57.1	73.5
(n=235)	CI	10.7–19.8	8.2–16.6	12.2–21.7	50.7–63.3	67.5–78.8

Note: The confidence interval (CI) is 95 percent.

Table BH1.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	15.0	15.7	15.9	53.3	69.3
(n=1,998)	CI	13.0–17.2	13.9–17.8	14.1–18.0	50.6–56.0	66.6–71.8
Male	%	16.7	15.6	15.6	52.1	67.7
(n=943)	CI	13.7–20.2	13.0–18.6	12.9–18.8	48.1–56.0	63.8–71.4
Female	%	12.3	16.0	16.6	55.1	71.7
(n=999)	CI	9.9–15.3	13.3–19.0	13.9–19.6	51.3–58.9	68.0–75.1
White	%	–	–	–	–	–
(n=1,622)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	7.5	14.0	15.6	62.9	78.5
(n=1,267)	CI	6.2–9.1	12.2–16.0	13.7–17.7	60.2–65.5	76.2–80.7
10th	%	16.2	19.2	17.0	47.5	64.5
(n=496)	CI	13.2–19.7	16.0–22.9	14.0–20.6	43.1–51.9	60.2–68.6
12th	%	21.4	13.9	15.1	49.6	64.7
(n=235)	CI	16.6–27.1	10.0–18.9	11.1–20.3	43.2–56.0	58.4–70.6

Note: The confidence interval (CI) is 95 percent.

Table BH1.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	12.7	11.5	14.4	61.5	75.8
(n=2,000)	CI	10.8–14.8	9.8–13.4	12.5–16.4	58.7–64.1	73.3–78.2
Male	%	13.1	10.8	14.2	61.8	76.1
(n=944)	CI	10.5–16.4	8.4–13.7	11.6–17.4	57.8–65.7	72.3–79.5
Female	%	11.1	12.0	14.9	61.9	76.9
(n=999)	CI	8.8–14.0	9.7–14.8	12.3–18.0	58.1–65.7	73.3–80.1
White	%	–	–	–	–	–
(n=1,623)	CI	–	–	–	–	–
Black	%	–	–	–	–	–
(n=49)	CI	–	–	–	–	–
Hispanic	%	–	–	–	–	–
(n=85)	CI	–	–	–	–	–
8th	%	4.5	8.6	11.9	75.0	86.9
(n=1,270)	CI	3.5–5.8	7.2–10.3	10.2–13.8	72.6–77.3	85.0–88.7
10th	%	15.7	13.3	14.9	56.2	71.1
(n=495)	CI	12.7–19.1	10.5–16.5	12.0–18.3	51.8–60.5	66.9–74.9
12th	%	18.1	12.6	16.4	52.9	69.3
(n=235)	CI	13.7–23.5	8.9–17.5	12.2–21.7	46.5–59.2	63.1–74.9

Note: The confidence interval (CI) is 95 percent.

Table BH1.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	62.7	24.9	12.4	37.3
(n=1,990)	CI	60.1–65.3	22.6–27.3	10.7–14.3	34.7–39.9
Male	%	63.6	24.5	11.9	36.4
(n=937)	CI	59.7–67.3	21.2–28.0	9.6–14.8	32.7–40.3
Female	%	61.6	26.0	12.4	38.4
(n=996)	CI	57.9–65.3	22.8–29.5	10.1–15.0	34.7–42.1
White	%	–	–	–	–
(n=1,611)	CI	–	–	–	–
Black	%	–	–	–	–
(n=49)	CI	–	–	–	–
Hispanic	%	–	–	–	–
(n=85)	CI	–	–	–	–
8th	%	59.1	28.1	12.8	40.9
(n=1,261)	CI	56.3–61.8	25.7–30.6	11.1–14.8	38.2–43.7
10th	%	65.9	21.2	12.8	34.1
(n=496)	CI	61.6–70.0	17.9–25.1	10.2–16.1	30.0–38.4
12th	%	63.1	25.4	11.4	36.9
(n=233)	CI	56.7–69.1	20.2–31.4	7.9–16.2	30.9–43.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	48.3	32.5	19.2	51.7
(n=1,989)	CI	45.6–51.1	30.0–35.1	17.1–21.4	48.9–54.4
Male	%	52.6	29.4	18.0	47.4
(n=941)	CI	48.7–56.5	26.0–33.1	15.1–21.3	43.5–51.3
Female	%	43.7	35.9	20.4	56.3
(n=991)	CI	39.9–47.6	32.3–39.7	17.5–23.6	52.4–60.1
White	%	–	–	–	–
(n=1,610)	CI	–	–	–	–
Black	%	–	–	–	–
(n=49)	CI	–	–	–	–
Hispanic	%	–	–	–	–
(n=85)	CI	–	–	–	–
8th	%	45.4	34.2	20.3	54.6
(n=1,263)	CI	42.7–48.2	31.6–36.9	18.2–22.7	51.8–57.3
10th	%	52.4	30.2	17.3	47.6
(n=493)	CI	48.0–56.8	26.3–34.4	14.2–20.9	43.2–52.0
12th	%	47.0	33.1	19.9	53.0
(n=233)	CI	40.7–53.5	27.3–39.3	15.3–25.5	46.5–59.3

Note: The confidence interval (CI) is 95 percent.

Table BH1.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	37.9	32.2	29.9	62.1
(n=1,993)	CI	35.2–40.6	29.7–34.8	27.5–32.4	59.4–64.8
Male	%	42.8	29.2	28.0	57.2
(n=942)	CI	38.9–46.7	25.8–32.9	24.7–31.6	53.3–61.1
Female	%	32.5	35.4	32.1	67.5
(n=995)	CI	28.9–36.3	31.8–39.2	28.7–35.8	63.7–71.1
White	%	–	–	–	–
(n=1,614)	CI	–	–	–	–
Black	%	–	–	–	–
(n=49)	CI	–	–	–	–
Hispanic	%	–	–	–	–
(n=85)	CI	–	–	–	–
8th	%	31.5	32.7	35.8	68.5
(n=1,266)	CI	29.0–34.1	30.2–35.3	33.2–38.5	65.9–71.0
10th	%	39.4	31.0	29.6	60.6
(n=494)	CI	35.2–43.8	27.1–35.2	25.7–33.8	56.2–64.8
12th	%	42.8	33.1	24.2	57.2
(n=233)	CI	36.6–49.2	27.3–39.3	19.1–30.1	50.8–63.4

Note: The confidence interval (CI) is 95 percent.

Behavioral Health Planning Region 2 Results Summary

Table BH2.1. Lifetime Prescription Drug Misuse

		On how many occasions (if any) have you ever <u>used prescription drugs not prescribed to you?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	92.8	3.8	1.3	0.8	0.5	0.2	0.7	7.2
(n=6,369)	CI	92.1-93.4	3.3-4.3	1.1-1.6	0.6-1.0	0.3-0.7	0.1-0.3	0.5-0.9	6.6-7.9
Male	%	93.7	3.3	1.1	0.6	0.3	0.2	0.8	6.3
(n=3,079)	CI	92.7-94.5	2.7-4.0	0.8-1.5	0.4-1.0	0.2-0.6	0.1-0.5	0.5-1.2	5.5-7.3
Female	%	92.3	4.1	1.6	0.9	0.6	0.2	0.4	7.7
(n=3,138)	CI	91.3-93.2	3.5-4.9	1.2-2.1	0.6-1.3	0.4-0.9	0.1-0.4	0.2-0.7	6.8-8.7
White	%	93.1	3.5	1.4	0.7	0.4	0.2	0.5	6.9
(n=5,332)	CI	92.4-93.8	3.1-4.1	1.1-1.7	0.5-1.0	0.3-0.7	0.1-0.4	0.4-0.8	6.2-7.6
Black	%	85.0	8.7	2.1	0.7	2.1	0.0	1.4	15.0
(n=142)	CI	78.0-90.1	5.0-14.8	0.7-6.4	0.1-4.6	0.7-6.4	-	0.3-5.5	9.9-22.0
Hispanic	%	95.6	1.8	0.0	0.9	0.4	0.0	1.3	4.4
(n=226)	CI	92.0-97.6	0.7-4.7	-	0.2-3.5	0.1-3.1	-	0.4-3.9	2.4-8.0
8th	%	94.6	3.3	0.7	0.5	0.3	0.1	0.5	5.4
(n=2,367)	CI	93.6-95.4	2.7-4.1	0.4-1.2	0.3-0.8	0.1-0.6	0.0-0.3	0.3-0.9	4.6-6.4
10th	%	91.5	4.2	1.5	1.1	0.6	0.2	0.8	8.5
(n=2,194)	CI	90.3-92.6	3.5-5.2	1.1-2.2	0.7-1.6	0.4-1.1	0.1-0.5	0.5-1.2	7.4-9.7
12th	%	92.1	3.7	1.8	0.8	0.6	0.3	0.8	7.9
(n=1,808)	CI	90.8-93.3	2.9-4.7	1.3-2.5	0.5-1.3	0.3-1.0	0.1-0.7	0.5-1.3	6.7-9.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.2. Lifetime Alcohol Use

		On how many occasions (if any) have you ever <u>had one or more drinks of an alcoholic beverage?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	63.7	16.7	6.5	4.1	3.2	2.0	3.8	36.3
(n=6,416)	CI	62.5-64.8	15.8-17.6	6.0-7.2	3.6-4.6	2.8-3.7	1.7-2.4	3.3-4.3	35.2-37.5
Male	%	64.2	15.2	6.4	4.0	3.2	2.0	4.9	35.8
(n=3,107)	CI	62.5-65.9	14.0-16.5	5.6-7.3	3.4-4.8	2.6-3.9	1.6-2.6	4.2-5.8	34.1-37.5
Female	%	63.3	18.2	6.8	4.1	3.4	1.9	2.4	36.7
(n=3,156)	CI	61.6-65.0	16.8-19.5	6.0-7.8	3.4-4.8	2.8-4.1	1.4-2.4	1.9-3.0	35.0-38.4
White	%	64.5	16.2	6.4	4.0	3.5	1.9	3.5	35.5
(n=5362)	CI	63.2-65.8	15.3-17.3	5.7-7.1	3.5-4.5	3.0-4.0	1.6-2.4	3.0-4.0	34.2-36.8
Black	%	55.3	22.3	8.5	3.5	2.9	3.0	4.5	44.7
(n=143)	CI	47.0-63.4	16.1-29.9	4.8-14.4	1.4-8.2	1.1-7.6	1.1-7.9	2.0-9.8	36.6-53.0
Hispanic	%	60.9	23.4	4.9	4.5	1.8	2.3	2.3	39.1
(n=227)	CI	54.3-67.1	18.3-29.4	2.7-8.6	2.4-8.2	0.7-4.9	0.9-5.3	1.0-5.5	32.9-45.7
8th	%	77.0	15.1	3.9	1.7	1.1	0.6	0.5	23.0
(n=2,382)	CI	75.3-78.7	13.8-16.6	3.2-4.8	1.3-2.3	0.7-1.6	0.4-1.0	0.3-0.9	21.3-24.7
10th	%	59.8	17.8	7.7	5.4	3.4	2.3	3.5	40.2
(n=2,214)	CI	57.8-61.9	16.2-19.4	6.6-8.9	4.5-6.4	2.7-4.3	1.8-3.1	2.8-4.4	38.1-42.2
12th	%	53.2	17.3	8.2	5.2	5.4	3.1	7.6	46.8
(n=1,820)	CI	50.9-55.5	15.6-19.1	7.0-9.6	4.3-6.3	4.4-6.5	2.4-4.0	6.4-8.9	44.5-49.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.3. Lifetime Binge Drinking

		On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	83.0	6.4	3.1	2.3	2.1	1.2	1.9	17.0
(n=6,402)	CI	82.0–83.9	5.8–7.0	2.7–3.6	2.0–2.7	1.8–2.5	0.9–1.5	1.6–2.2	16.1–18.0
Male	%	81.8	6.0	3.1	2.7	2.3	1.5	2.6	18.2
(n=3,097)	CI	80.4–83.1	5.2–6.9	2.6–3.8	2.1–3.3	1.8–2.9	1.1–2.0	2.1–3.3	16.9–19.6
Female	%	84.3	7.0	3.1	2.1	1.9	0.8	0.8	15.7
(n=3,153)	CI	82.9–85.5	6.1–7.9	2.6–3.8	1.6–2.7	1.5–2.5	0.5–1.2	0.6–1.2	14.5–17.1
White	%	83.3	6.3	3.1	2.3	2.2	1.1	1.7	16.7
(n=5,351)	CI	82.2–84.3	5.7–7.0	2.7–3.6	1.9–2.7	1.8–2.6	0.8–1.4	1.4–2.1	15.7–17.8
Black	%	80.5	6.3	2.9	3.5	1.6	2.9	2.3	19.5
(n=143)	CI	73.0–86.4	3.3–11.8	1.1–7.5	1.5–8.2	0.4–6.1	1.1–7.7	0.7–6.8	13.6–27.0
Hispanic	%	83.9	6.9	3.2	2.3	1.4	1.0	1.3	16.1
(n=226)	CI	78.3–88.2	4.2–11.2	1.5–6.6	0.9–5.4	0.5–4.3	0.2–3.9	0.4–4.1	11.8–21.7
8th	%	94.4	3.4	0.7	0.6	0.5	0.1	0.3	5.6
(n=2,377)	CI	93.4–95.2	2.8–4.3	0.4–1.1	0.4–1.0	0.3–0.8	0.0–0.3	0.2–0.7	4.8–6.6
10th	%	81.2	7.4	3.5	2.8	2.3	1.2	1.5	18.8
(n=2,211)	CI	79.5–82.8	6.4–8.6	2.8–4.4	2.2–3.6	1.8–3.0	0.8–1.8	1.1–2.1	17.2–20.5
12th	%	72.5	8.6	5.4	3.6	3.6	2.3	4.0	27.5
(n=1,814)	CI	70.4–74.5	7.4–10.0	4.4–6.5	2.9–4.6	2.9–4.6	1.7–3.1	3.2–5.0	25.5–29.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.4. Lifetime Cigarette Use

		On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	80.5	9.1	2.8	1.7	1.6	0.9	3.5	19.5
(n=6,402)	CI	79.5–81.4	8.4–9.8	2.4–3.2	1.4–2.1	1.3–1.9	0.7–1.2	3.1–4.0	18.6–20.5
Male	%	80.2	8.9	3.0	1.4	1.5	1.0	4.0	19.8
(n=3,101)	CI	78.7–81.5	7.9–9.9	2.5–3.7	1.1–1.9	1.1–2.0	0.7–1.4	3.4–4.8	18.5–21.3
Female	%	81.2	9.1	2.7	2.0	1.6	0.7	2.8	18.8
(n=3,150)	CI	79.8–82.6	8.1–10.1	2.1–3.3	1.6–2.6	1.2–2.1	0.4–1.0	2.3–3.4	17.4–20.2
White	%	80.6	9.0	2.9	1.7	1.5	0.8	3.5	19.4
(n=5,352)	CI	79.5–81.6	8.3–9.8	2.5–3.4	1.3–2.0	1.2–1.9	0.6–1.1	3.1–4.1	18.4–20.5
Black	%	81.9	7.1	2.3	2.3	2.1	1.5	2.8	18.1
(n=143)	CI	74.5–87.5	3.8–12.8	0.7–7.1	0.7–6.8	0.7–6.5	0.4–5.8	1.0–7.3	12.5–25.5
Hispanic	%	85.2	5.5	1.8	2.7	1.0	1.8	2.1	14.8
(n=228)	CI	79.9–89.3	3.1–9.4	0.7–4.8	1.2–5.9	0.2–3.8	0.7–4.6	0.9–5.0	10.7–20.1
8th	%	87.7	7.5	2.0	0.9	0.5	0.3	1.1	12.3
(n=2,378)	CI	86.3–89.0	6.5–8.6	1.5–2.7	0.6–1.4	0.3–0.9	0.1–0.6	0.8–1.7	11.0–13.7
10th	%	78.6	10.0	2.6	1.8	1.5	1.4	4.1	21.4
(n=2,212)	CI	76.9–80.3	8.8–11.3	2.0–3.4	1.3–2.4	1.1–2.1	1.0–2.0	3.3–5.0	19.7–23.1
12th	%	74.5	9.8	3.9	2.6	2.8	1.0	5.4	25.5
(n=1,814)	CI	72.4–76.4	8.5–11.2	3.1–4.8	2.0–3.5	2.1–3.6	0.7–1.6	4.5–6.6	23.6–27.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	72.4	9.5	4.0	2.3	2.1	1.9	7.8	27.6
(n=6,375)	CI	71.3–73.5	8.8–10.2	3.5–4.5	2.0–2.7	1.8–2.5	1.6–2.3	7.2–8.5	26.5–28.7
Male	%	70.4	9.4	3.9	2.1	2.1	2.2	9.9	29.6
(n=3,086)	CI	68.7–72.0	8.4–10.5	3.3–4.7	1.6–2.7	1.7–2.7	1.7–2.8	8.9–11.0	28.0–31.3
Female	%	74.6	9.6	3.9	2.6	2.0	1.7	5.6	25.4
(n=3,137)	CI	73.1–76.1	8.6–10.7	3.3–4.7	2.1–3.2	1.6–2.6	1.3–2.2	4.8–6.5	23.9–26.9
White	%	72.4	9.3	4.0	2.4	2.1	2.0	7.7	27.6
(n=5,331)	CI	71.2–73.6	8.6–10.1	3.5–4.6	2.0–2.9	1.7–2.5	1.7–2.5	7.0–8.4	26.4–28.8
Black	%	68.4	11.3	3.5	3.6	0.7	0.8	11.7	31.6
(n=141)	CI	60.1–75.7	7.0–17.8	1.5–8.3	1.5–8.5	0.1–4.9	0.1–5.5	7.2–18.3	24.3–39.9
Hispanic	%	81.9	6.5	1.8	2.2	2.1	2.3	3.2	18.1
(n=226)	CI	76.2–86.4	4.0–10.6	0.7–4.7	0.9–5.2	0.9–5.0	1.0–5.5	1.5–6.5	13.6–23.8
8th	%	81.0	8.5	3.4	1.6	1.0	1.3	3.1	19.0
(n=2,370)	CI	79.4–82.6	7.5–9.7	2.7–4.2	1.2–2.2	0.7–1.5	1.0–1.9	2.5–3.9	17.4–20.6
10th	%	69.4	10.1	4.1	2.9	3.0	1.9	8.5	30.6
(n=2,196)	CI	67.5–71.3	9.0–11.5	3.3–5.0	2.3–3.7	2.4–3.8	1.4–2.6	7.4–9.7	28.7–32.5
12th	%	66.2	9.8	4.5	2.5	2.3	2.5	12.1	33.8
(n=1,809)	CI	64.0–68.3	8.5–11.3	3.7–5.6	1.9–3.4	1.7–3.1	1.9–3.3	10.7–13.7	31.7–36.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	88.4	4.1	1.7	1.0	0.8	0.5	3.5	11.6
(n=6,390)	CI	87.6–89.2	3.7–4.6	1.4–2.1	0.8–1.3	0.6–1.1	0.4–0.7	3.0–3.9	10.8–12.4
Male	%	82.9	5.2	2.3	1.6	1.2	0.8	6.0	17.1
(n=3,103)	CI	81.5–84.2	4.5–6.1	1.8–2.9	1.2–2.2	0.9–1.7	0.5–1.2	5.2–6.9	15.8–18.5
Female	%	94.0	3.1	1.1	0.4	0.3	0.2	0.9	6.0
(n=3,134)	CI	93.1–94.8	2.5–3.8	0.8–1.5	0.2–0.7	0.2–0.6	0.1–0.4	0.6–1.3	5.2–6.9
White	%	88.3	4.0	1.8	1.0	0.8	0.5	3.7	11.7
(n=5,342)	CI	87.4–89.1	3.5–4.5	1.5–2.2	0.8–1.3	0.6–1.1	0.3–0.7	3.2–4.2	10.9–12.6
Black	%	87.9	4.9	2.1	0.8	0.7	1.4	2.2	12.1
(n=142)	CI	81.3–92.4	2.3–10.1	0.7–6.4	0.1–5.5	0.1–4.6	0.4–5.7	0.7–6.6	7.6–18.7
Hispanic	%	92.0	5.3	0.9	0.5	0.4	0.0	0.8	8.0
(n=225)	CI	87.7–94.9	3.0–9.1	0.2–3.7	0.1–3.5	0.1–3.1	–	0.2–3.3	5.1–12.3
8th	%	93.1	3.2	0.9	0.8	0.5	0.2	1.3	6.9
(n=2,374)	CI	92.0–94.1	2.6–4.0	0.6–1.4	0.5–1.2	0.3–0.8	0.1–0.5	0.9–1.9	5.9–8.0
10th	%	86.7	4.8	2.1	1.0	0.9	0.5	3.9	13.3
(n=2,202)	CI	85.2–88.1	4.0–5.8	1.6–2.8	0.7–1.6	0.6–1.3	0.3–1.0	3.2–4.8	11.9–14.8
12th	%	85.0	4.3	2.3	1.2	1.1	0.8	5.3	15.0
(n=1,814)	CI	83.3–86.6	3.5–5.4	1.7–3.0	0.8–1.8	0.7–1.7	0.5–1.3	4.3–6.4	13.4–16.7

Note: The confidence interval (CI) is 95 percent.

Table BH2.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	83.1	5.5	2.4	1.4	1.5	1.3	4.9	16.9
(n=6,388)	CI	82.1–84.0	5.0–6.1	2.0–2.8	1.1–1.7	1.2–1.8	1.0–1.6	4.4–5.5	16.0–17.9
Male	%	83.1	5.6	2.2	1.2	1.5	1.1	5.3	16.9
(n=3,096)	CI	81.8–84.4	4.8–6.4	1.7–2.8	0.8–1.6	1.1–2.0	0.8–1.6	4.5–6.1	15.6–18.2
Female	%	83.5	5.3	2.6	1.6	1.3	1.4	4.3	16.5
(n=3,138)	CI	82.2–84.8	4.6–6.2	2.1–3.2	1.2–2.1	1.0–1.8	1.0–1.8	3.6–5.0	15.2–17.8
White	%	84.4	5.2	2.1	1.5	1.3	1.2	4.4	15.6
(n=5,343)	CI	83.3–85.3	4.6–5.8	1.8–2.6	1.2–1.8	1.0–1.6	0.9–1.6	3.9–5.0	14.7–16.7
Black	%	66.4	9.8	2.8	1.4	2.8	1.5	15.3	33.6
(n=142)	CI	58.1–73.8	5.9–16.0	1.0–7.3	0.3–5.5	1.1–7.4	0.4–5.8	10.2–22.4	26.2–41.9
Hispanic	%	83.7	6.0	3.1	1.4	1.4	0.9	3.6	16.3
(n=226)	CI	78.2–88.0	3.5–10.1	1.5–6.4	0.4–4.2	0.5–4.3	0.2–3.6	1.8–7.0	12.0–21.8
8th	%	92.1	3.8	1.1	0.7	0.6	0.4	1.4	7.9
(n=2,368)	CI	90.9–93.1	3.1–4.6	0.7–1.6	0.4–1.1	0.4–1.0	0.2–0.8	1.0–2.0	6.9–9.1
10th	%	80.3	6.5	2.9	1.4	1.9	1.6	5.3	19.7
(n=2,213)	CI	78.6–81.9	5.6–7.7	2.3–3.7	1.0–2.0	1.4–2.5	1.2–2.2	4.5–6.3	18.1–21.4
12th	%	76.3	6.4	3.3	2.2	2.0	1.8	8.2	23.7
(n=1,807)	CI	74.2–78.2	5.4–7.6	2.5–4.2	1.6–2.9	1.4–2.7	1.3–2.5	7.0–9.5	21.8–25.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	99.0	0.4	0.1	0.1	0.1	0.0	0.2	1.0
(n=6,398)	CI	98.7–99.2	0.3–0.6	0.1–0.2	0.1–0.3	0.0–0.2	0.0–0.1	0.1–0.3	0.8–1.3
Male	%	98.8	0.5	0.2	0.2	0.1	0.0	0.1	1.2
(n=3,100)	CI	98.4–99.1	0.3–0.8	0.1–0.4	0.1–0.5	0.0–0.3	0.0–0.2	0.0–0.3	0.9–1.6
Female	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7
(n=3,144)	CI	–	–	–	–	–	–	–	0.4–1.0
White	%	99.1	0.4	0.1	0.1	0.1	0.0	0.1	0.9
(n=5,351)	CI	98.8–99.3	0.3–0.7	0.1–0.2	0.1–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.7–1.2
Black	%	97.8	2.2	0.0	0.0	0.0	0.0	0.0	2.2
(n=142)	CI	93.4–99.3	0.7–6.6	–	–	–	–	–	0.7–6.6
Hispanic	%	99.2	0.0	0.4	0.0	0.0	0.0	0.4	0.8
(n=226)	CI	96.6–99.8	–	0.1–3.1	–	–	–	0.1–2.9	0.2–3.4
8th	%	99.5	0.3	0.0	0.0	0.0	0.0	0.1	0.5
(n=2,371)	CI	99.1–99.7	0.2–0.7	0.0–0.3	0.0–0.3	–	–	0.0–0.3	0.3–0.9
10th	%	98.8	0.5	0.3	0.1	0.0	0.0	0.3	1.2
(n=2,213)	CI	98.3–99.2	0.2–0.8	0.1–0.6	0.0–0.4	0.0–0.3	0.0–0.3	0.1–0.6	0.8–1.7
12th	%	98.6	0.6	0.1	0.3	0.2	0.1	0.2	1.4
(n=1,814)	CI	97.9–99.0	0.3–1.0	0.0–0.4	0.1–0.7	0.1–0.6	0.0–0.4	0.1–0.6	1.0–2.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	97.6	1.1	0.4	0.3	0.2	0.1	0.3	2.4
(n=6,399)	CI	97.2-98.0	0.9-1.4	0.3-0.6	0.2-0.5	0.1-0.3	0.0-0.2	0.2-0.4	2.0-2.8
Male	%	97.4	1.3	0.4	0.4	0.2	0.0	0.2	2.6
(n=3,096)	CI	96.8-97.9	0.9-1.8	0.3-0.7	0.2-0.7	0.1-0.5	0.0-0.2	0.1-0.5	2.1-3.2
Female	%	98.1	0.9	0.4	0.1	0.2	0.1	0.2	1.9
(n=3,149)	CI	97.6-98.5	0.6-1.3	0.2-0.7	0.0-0.4	0.1-0.4	0.0-0.3	0.1-0.4	1.5-2.4
White	%	98.0	1.0	0.3	0.2	0.2	0.1	0.2	2.0
(n=5,347)	CI	97.6-98.3	0.8-1.3	0.2-0.6	0.1-0.4	0.1-0.3	0.0-0.2	0.1-0.4	1.7-2.4
Black	%	95.6	2.2	0.7	0.8	0.0	0.0	0.8	4.4
(n=142)	CI	90.4-98.0	0.7-6.6	0.1-4.9	0.1-5.5	-	-	0.1-5.5	2.0-9.6
Hispanic	%	96.1	1.8	1.3	0.5	0.0	0.0	0.4	3.9
(n=228)	CI	92.5-97.9	0.7-4.6	0.4-4.0	0.1-3.4	-	-	0.1-2.9	2.1-7.5
8th	%	98.9	0.5	0.1	0.1	0.1	0.0	0.2	1.1
(n=2,372)	CI	98.4-99.3	0.3-0.8	0.0-0.4	0.0-0.4	0.0-0.3	0.0-0.3	0.1-0.5	0.7-1.6
10th	%	97.1	1.5	0.6	0.3	0.2	0.2	0.1	2.9
(n=2,211)	CI	96.3-97.7	1.1-2.1	0.4-1.1	0.1-0.6	0.1-0.5	0.1-0.5	0.0-0.4	2.3-3.7
12th	%	96.8	1.4	0.5	0.5	0.3	0.1	0.4	3.2
(n=1,816)	CI	95.8-97.5	1.0-2.1	0.3-0.9	0.3-1.0	0.1-0.7	0.0-0.4	0.2-0.9	2.5-4.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.10. Age of First Prescription Drug Misuse

		How old were you when you first used prescription drugs not prescribed to you?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	94.0	1.1	0.5	0.6	0.9	0.9	1.1	0.5	0.4	13.23
(n=6,423)	CI	93.4–94.5	0.9–1.4	0.3–0.7	0.4–0.8	0.7–1.2	0.7–1.2	0.9–1.4	0.3–0.7	0.3–0.7	13.0–13.5
Male	%	94.9	1.2	0.4	0.3	0.7	0.6	1.0	0.4	0.5	13.28
(n=3,109)	CI	94.1–95.6	0.9–1.6	0.2–0.7	0.1–0.5	0.5–1.1	0.4–1.0	0.7–1.4	0.3–0.8	0.3–0.9	12.9–13.7
Female	%	93.4	0.8	0.5	0.8	1.0	1.2	1.3	0.5	0.4	13.34
(n=3,159)	CI	92.5–94.2	0.6–1.2	0.3–0.9	0.6–1.2	0.7–1.5	0.9–1.7	0.9–1.8	0.3–0.8	0.2–0.7	13.1–13.6
White	%	94.3	1.1	0.5	0.5	0.8	0.9	1.2	0.4	0.4	13.22
(n=5,368)	CI	93.7–94.9	0.8–1.4	0.3–0.7	0.4–0.7	0.6–1.1	0.7–1.2	0.9–1.5	0.3–0.6	0.2–0.6	13.0–13.5
Black	%	87.9	2.8	1.3	0.0	2.8	2.3	0.0	1.5	1.6	13.16
(n=143)	CI	81.2–92.4	1.0–7.3	0.3–5.1	–	1.0–7.2	0.7–6.8	–	0.4–5.8	0.4–6.1	11.9–14.5
Hispanic	%	96.1	0.8	0.0	0.0	0.5	1.3	1.3	0.0	0.0	13.35
(n=229)	CI	92.6–98.0	0.2–3.3	–	–	0.1–3.4	0.4–4.0	0.4–4.0	–	–	11.9–14.8
8th	%	95.8	1.4	0.5	0.7	1.1	0.4	0.0	0.0	0.0	11.68
(n=2,385)	CI	94.9–96.5	1.0–1.9	0.3–0.9	0.4–1.1	0.8–1.6	0.2–0.8	–	–	–	11.4–12.0
10th	%	92.8	1.2	0.3	0.6	1.1	1.5	2.2	0.3	0.0	13.28
(n=2,221)	CI	91.7–93.8	0.8–1.8	0.1–0.6	0.4–1.1	0.7–1.6	1.1–2.1	1.7–2.9	0.2–0.7	–	13.0–13.6
12th	%	93.2	0.8	0.5	0.3	0.5	0.8	1.2	1.1	1.4	14.24
(n=1,817)	CI	92.0–94.3	0.5–1.4	0.3–1.0	0.1–0.7	0.3–0.9	0.5–1.4	0.8–1.8	0.7–1.7	1.0–2.1	13.8–14.7

Note: The confidence interval (CI) is 95 percent.

Table BH2.11. Age of First Alcohol Use

		How old were you when you first had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	67.0	4.2	1.9	2.7	4.5	5.2	6.2	4.8	3.6	13.83
(n=6,420)	CI	65.9–68.2	3.7–4.7	1.6–2.2	2.3–3.1	4.0–5.0	4.7–5.8	5.6–6.8	4.3–5.3	3.2–4.1	13.7–13.9
Male	%	67.5	4.8	2.0	2.6	4.0	4.8	6.1	4.2	4.0	13.76
(n=3,108)	CI	65.8–69.1	4.1–5.6	1.6–2.6	2.1–3.3	3.4–4.7	4.1–5.6	5.3–7.0	3.5–5.0	3.4–4.8	13.6–13.9
Female	%	66.8	3.4	1.7	2.7	4.9	5.4	6.5	5.3	3.4	13.95
(n=3,156)	CI	65.1–68.4	2.8–4.0	1.3–2.2	2.2–3.3	4.2–5.7	4.7–6.3	5.6–7.4	4.6–6.2	2.8–4.1	13.8–14.1
White	%	67.8	3.8	1.7	2.7	4.3	5.2	6.0	4.8	3.6	13.89
(n=5,364)	CI	66.6–69.1	3.3–4.3	1.4–2.1	2.3–3.2	3.8–4.9	4.6–5.8	5.4–6.7	4.3–5.5	3.1–4.1	13.8–14.0
Black	%	60.6	6.3	1.3	1.4	6.8	2.1	11.5	6.8	3.1	13.89
(n=142)	CI	52.2–68.5	3.3–11.8	0.3–5.2	0.3–5.3	3.7–12.3	0.7–6.3	7.1–18.1	3.6–12.7	1.2–8.2	13.3–14.5
Hispanic	%	62.6	4.5	1.3	2.6	4.2	6.9	10.7	2.4	4.8	13.96
(n=230)	CI	56.1–68.7	2.5–8.0	0.4–4.0	1.2–5.8	2.3–7.6	4.2–11.0	7.3–15.5	1.0–5.6	2.6–8.8	13.5–14.4
8th	%	80.5	4.0	2.4	3.6	6.2	3.1	0.0	0.0	0.0	12.11
(n=2,383)	CI	78.9–82.1	3.3–4.9	1.9–3.1	3.0–4.5	5.3–7.3	2.5–3.9	0.0–0.3	–	–	12.0–12.2
10th	%	63.4	5.2	2.0	2.5	4.7	8.6	10.3	3.1	0.2	13.47
(n=2,222)	CI	61.3–65.3	4.4–6.2	1.5–2.7	1.9–3.2	3.9–5.6	7.5–9.9	9.1–11.7	2.5–3.9	0.1–0.5	13.3–13.6
12th	%	56.2	3.1	1.0	1.9	2.4	3.7	8.5	11.8	11.3	15.01
(n=1,815)	CI	53.9–58.5	2.4–4.0	0.7–1.6	1.4–2.7	1.8–3.2	3.0–4.7	7.3–9.8	10.4–13.4	9.9–12.8	14.9–15.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.12. Age of First Cigarette Use

		How old were you when you first <u>smoked a cigarette, even just a puff?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	81.1	3.9	1.7	2.2	2.5	2.4	2.4	2.0	1.9	13.18
(n=6,414)	CI	80.1–82.1	3.4–4.4	1.4–2.0	1.8–2.6	2.2–3.0	2.1–2.8	2.0–2.8	1.6–2.3	1.6–2.3	13.0–13.3
Male	%	80.9	4.6	1.7	1.9	2.2	2.4	2.3	1.8	2.1	13.09
(n=3,107)	CI	79.5–82.3	3.9–5.4	1.3–2.2	1.5–2.5	1.8–2.8	1.9–3.0	1.8–2.9	1.4–2.3	1.7–2.7	12.9–13.3
Female	%	81.8	2.9	1.6	2.4	2.6	2.4	2.4	2.2	1.7	13.35
(n=3,152)	CI	80.4–83.1	2.4–3.5	1.2–2.0	1.9–3.0	2.1–3.3	1.9–3.0	1.9–3.0	1.7–2.8	1.3–2.2	13.2–13.5
White	%	81.2	3.8	1.6	2.2	2.5	2.5	2.5	1.9	1.8	13.21
(n=5,360)	CI	80.2–82.3	3.3–4.3	1.3–1.9	1.9–2.6	2.1–2.9	2.1–3.0	2.1–2.9	1.6–2.4	1.5–2.2	13.1–13.4
Black	%	81.9	2.7	3.5	2.2	1.3	0.8	0.7	3.1	3.8	13.53
(n=143)	CI	74.5–87.5	1.0–7.0	1.4–8.2	0.7–6.7	0.3–5.1	0.1–5.4	0.1–4.9	1.2–8.1	1.6–8.9	12.4–14.7
Hispanic	%	86.2	2.7	0.4	1.2	2.2	2.1	1.3	1.5	2.4	13.64
(n=229)	CI	80.9–90.1	1.2–5.8	0.1–2.9	0.4–3.8	0.9–5.2	0.9–5.0	0.4–4.1	0.5–4.4	1.0–5.7	12.7–14.5
8th	%	88.3	4.0	1.8	2.5	2.3	1.0	0.1	0.0	0.0	11.58
(n=2,386)	CI	86.9–89.5	3.3–4.8	1.3–2.4	2.0–3.2	1.8–3.0	0.7–1.5	0.0–0.3	0.0–0.3	–	11.4–11.7
10th	%	79.4	4.8	1.8	2.3	3.0	4.1	3.7	0.8	0.2	12.73
(n=2,213)	CI	77.6–81.0	4.0–5.8	1.3–2.5	1.7–3.0	2.3–3.8	3.4–5.0	3.0–4.6	0.5–1.2	0.1–0.5	12.5–12.9
12th	%	75.2	2.8	1.4	1.7	2.3	2.0	3.4	5.3	5.8	14.41
(n=1,815)	CI	73.2–77.1	2.1–3.7	1.0–2.1	1.2–2.4	1.7–3.1	1.5–2.8	2.7–4.3	4.4–6.5	4.8–6.9	14.2–14.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	73.5	1.3	1.2	2.0	3.7	4.6	5.6	3.6	4.5	14.36
(n=6,384)	CI	72.4–74.5	1.1–1.6	0.9–1.5	1.7–2.4	3.3–4.2	4.2–5.2	5.1–6.2	3.1–4.1	4.0–5.0	14.3–14.5
Male	%	71.9	1.5	1.0	2.3	3.6	5.2	6.1	3.9	4.6	14.36
(n=3,088)	CI	70.2–73.4	1.1–2.0	0.7–1.5	1.8–2.9	3.0–4.3	4.4–6.0	5.3–7.0	3.2–4.6	3.9–5.4	14.2–14.5
Female	%	75.2	1.0	1.2	1.8	3.7	4.1	5.1	3.4	4.5	14.43
(n=3,145)	CI	73.6–76.7	0.7–1.4	0.9–1.7	1.4–2.3	3.1–4.4	3.4–4.8	4.4–6.0	2.8–4.1	3.8–5.3	14.3–14.6
White	%	73.5	1.2	1.1	2.0	3.7	4.7	5.8	3.7	4.5	14.40
(n=5,337)	CI	72.2–74.6	0.9–1.5	0.9–1.5	1.7–2.4	3.2–4.2	4.1–5.3	5.2–6.4	3.2–4.2	3.9–5.1	14.3–14.5
Black	%	68.1	2.1	3.3	1.3	4.7	2.6	7.0	4.6	6.2	14.28
(n=143)	CI	59.9–75.4	0.7–6.4	1.4–7.8	0.3–5.3	2.2–9.7	1.0–6.9	3.8–12.6	2.1–9.9	3.1–11.9	13.6–14.9
Hispanic	%	82.7	0.4	0.0	2.1	4.2	3.5	2.7	0.5	3.9	14.30
(n=226)	CI	77.1–87.1	0.1–2.9	–	0.9–5.1	2.3–7.7	1.7–6.9	1.2–6.0	0.1–3.4	1.9–7.6	13.7–14.9
8th	%	82.2	1.9	1.7	3.4	6.7	3.9	0.2	0.0	0.0	12.54
(n=2,379)	CI	80.6–83.7	1.4–2.5	1.3–2.3	2.7–4.2	5.7–7.8	3.2–4.8	0.1–0.5	–	–	12.4–12.7
10th	%	70.5	1.4	1.4	1.5	2.8	7.1	12.2	3.0	0.2	14.13
(n=2,196)	CI	68.6–72.4	1.0–1.9	1.0–1.9	1.0–2.1	2.2–3.6	6.1–8.2	10.9–13.6	2.4–3.9	0.1–0.5	14.0–14.2
12th	%	67.1	0.6	0.4	1.1	1.5	2.9	4.5	8.0	14.0	15.67
(n=1,809)	CI	64.9–69.2	0.3–1.1	0.2–0.8	0.7–1.7	1.0–2.2	2.2–3.7	3.7–5.6	6.8–9.3	12.4–15.6	15.5–15.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	83.3	1.2	1.0	1.5	2.4	2.8	3.3	2.3	2.2	14.09
(n=6,421)	CI	82.3–84.2	0.9–1.5	0.8–1.2	1.3–1.9	2.1–2.8	2.4–3.2	2.9–3.8	2.0–2.7	1.8–2.6	14.0–14.2
Male	%	83.7	1.4	1.1	1.6	2.0	2.7	3.5	2.0	2.0	13.95
(n=3,107)	CI	82.3–84.9	1.1–1.9	0.8–1.6	1.2–2.1	1.6–2.6	2.2–3.3	2.9–4.2	1.5–2.5	1.6–2.6	13.8–14.1
Female	%	83.4	0.7	0.8	1.4	2.6	2.9	3.1	2.6	2.4	14.30
(n=3,158)	CI	82.1–84.7	0.5–1.1	0.5–1.1	1.1–1.9	2.1–3.2	2.4–3.6	2.6–3.8	2.1–3.3	1.9–3.0	14.1–14.5
White	%	84.5	0.9	0.9	1.4	2.2	2.7	3.1	2.2	2.1	14.16
(n=5,366)	CI	83.5–85.4	0.7–1.2	0.7–1.2	1.1–1.7	1.8–2.6	2.3–3.1	2.7–3.7	1.8–2.7	1.7–2.5	14.0–14.3
Black	%	66.6	4.3	2.6	2.7	6.9	4.4	6.4	3.7	2.3	13.49
(n=143)	CI	58.4–74.0	1.9–9.3	1.0–6.9	1.0–7.2	3.7–12.4	2.0–9.6	3.3–12.0	1.5–8.7	0.7–6.8	12.9–14.1
Hispanic	%	83.9	0.9	0.9	1.9	2.2	2.6	4.4	1.9	1.5	14.04
(n=229)	CI	78.4–88.2	0.2–3.4	0.2–3.4	0.7–4.9	0.9–5.2	1.2–5.7	2.3–7.9	0.7–5.1	0.5–4.4	13.4–14.7
8th	%	92.3	0.9	1.1	1.7	2.7	1.2	0.1	0.0	0.0	12.31
(n=2,381)	CI	91.2–93.3	0.6–1.4	0.7–1.6	1.2–2.3	2.1–3.5	0.8–1.7	0.0–0.3	–	–	12.1–12.5
10th	%	80.7	1.5	1.3	1.5	3.1	4.4	6.1	1.4	0.0	13.65
(n=2,221)	CI	79.0–82.3	1.1–2.1	0.9–1.8	1.1–2.1	2.4–3.9	3.6–5.3	5.2–7.2	1.0–2.0	0.0–0.3	13.5–13.8
12th	%	76.1	1.1	0.5	1.4	1.3	2.8	3.9	5.9	6.9	15.10
(n=1,819)	CI	74.1–78.0	0.7–1.7	0.3–1.0	1.0–2.1	0.9–2.0	2.1–3.7	3.1–4.9	4.9–7.0	5.8–8.2	14.9–15.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.15. Past Month Prescription Drug Misuse

During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>									
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	97.4	1.5	0.5	0.2	0.2	0.1	0.2	2.6
(n=6,435)	CI	97.0–97.7	1.2–1.8	0.3–0.7	0.1–0.3	0.1–0.4	0.0–0.2	0.1–0.3	2.3–3.0
Male	%	98.0	1.1	0.4	0.2	0.1	0.1	0.0	2.0
(n=3,117)	CI	97.5–98.5	0.8–1.6	0.2–0.7	0.1–0.5	0.1–0.4	0.0–0.3	0.0–0.2	1.5–2.5
Female	%	97.2	1.9	0.5	0.1	0.2	0.0	0.1	2.8
(n=3,162)	CI	96.5–97.7	1.4–2.4	0.3–0.8	0.0–0.3	0.1–0.5	0.0–0.2	0.0–0.3	2.3–3.5
White	%	97.7	1.4	0.4	0.2	0.2	0.1	0.1	2.3
(n=5,377)	CI	97.2–98.1	1.1–1.7	0.3–0.7	0.1–0.3	0.1–0.3	0.0–0.2	0.0–0.2	1.9–2.8
Black	%	95.1	3.6	0.0	0.0	0.7	0.0	0.7	4.9
(n=143)	CI	89.9–97.7	1.5–8.4	–	–	0.1–4.9	–	0.1–4.9	2.3–10.1
Hispanic	%	97.4	2.1	0.0	0.5	0.0	0.0	0.0	2.6
(n=228)	CI	94.2–98.8	0.9–5.1	–	0.1–3.4	–	–	–	1.2–5.8
8th	%	97.6	1.5	0.4	0.2	0.2	0.1	0.1	2.4
(n=2,388)	CI	96.9–98.2	1.1–2.0	0.2–0.7	0.1–0.4	0.1–0.4	0.0–0.3	0.0–0.4	1.8–3.1
10th	%	96.8	1.8	0.9	0.2	0.2	0.0	0.2	3.2
(n=2,224)	CI	95.9–97.4	1.3–2.4	0.5–1.3	0.1–0.5	0.1–0.5	0.0–0.3	0.1–0.5	2.6–4.1
12th	%	97.8	1.3	0.2	0.2	0.3	0.1	0.2	2.2
(n=1,823)	CI	97.0–98.4	0.9–2.0	0.1–0.5	0.1–0.5	0.1–0.7	0.0–0.4	0.1–0.6	1.6–3.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.16. Past Month Alcohol Use

During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>									
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	85.0	8.2	3.8	0.9	1.2	0.5	0.4	15.0
(n=6,411)	CI	84.1–85.9	7.6–8.9	3.3–4.3	0.7–1.2	0.9–1.5	0.3–0.7	0.3–0.6	14.1–15.9
Male	%	85.1	7.6	3.9	1.1	1.2	0.6	0.4	14.9
(n=3,104)	CI	83.8–86.3	6.7–8.6	3.3–4.7	0.8–1.6	0.9–1.7	0.4–1.0	0.2–0.7	13.7–16.2
Female	%	85.1	8.8	3.8	0.7	1.1	0.3	0.3	14.9
(n=3,151)	CI	83.8–86.4	7.8–9.8	3.1–4.5	0.5–1.1	0.8–1.5	0.1–0.5	0.1–0.5	13.6–16.2
White	%	85.4	7.9	3.9	1.0	1.1	0.4	0.4	14.6
(n=5,360)	CI	84.4–86.4	7.2–8.7	3.4–4.4	0.7–1.3	0.8–1.4	0.2–0.6	0.3–0.6	13.6–15.6
Black	%	83.0	8.8	5.2	0.0	1.5	0.7	0.8	17.0
(n=139)	CI	75.6–88.5	5.0–14.9	2.5–10.6	–	0.4–6.0	0.1–4.7	0.1–5.6	11.5–24.4
Hispanic	%	82.2	10.4	3.7	0.4	1.4	1.4	0.4	17.8
(n=227)	CI	76.6–86.8	7.0–15.2	1.8–7.2	0.1–3.1	0.5–4.3	0.5–4.3	0.1–3.1	13.2–23.4
8th	%	92.1	5.2	1.5	0.3	0.4	0.3	0.2	7.9
(n=2,383)	CI	91.0–93.1	4.4–6.2	1.1–2.0	0.2–0.7	0.2–0.7	0.1–0.6	0.1–0.5	6.9–9.0
10th	%	82.6	9.6	4.7	1.0	1.2	0.4	0.4	17.4
(n=2,217)	CI	81.0–84.2	8.5–10.9	3.9–5.7	0.7–1.5	0.8–1.8	0.2–0.7	0.2–0.8	15.8–19.0
12th	%	79.8	10.0	5.3	1.5	2.0	0.8	0.7	20.2
(n=1,811)	CI	77.9–81.6	8.7–11.5	4.3–6.4	1.0–2.2	1.4–2.7	0.5–1.3	0.4–1.2	18.4–22.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.17. Past Month Binge Drinking

		During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	92.5	3.9	1.9	0.7	0.5	0.1	0.3	7.5
(n=6,419)	CI	91.8–93.1	3.5–4.5	1.6–2.3	0.5–0.9	0.4–0.8	0.1–0.3	0.2–0.4	6.9–8.2
Male	%	91.3	4.3	2.4	0.8	0.7	0.2	0.3	8.7
(n=3,107)	CI	90.2–92.3	3.6–5.1	1.9–3.0	0.6–1.2	0.4–1.0	0.1–0.5	0.1–0.5	7.7–9.8
Female	%	93.9	3.6	1.5	0.6	0.3	0.0	0.1	6.1
(n=3,156)	CI	92.9–94.7	3.0–4.3	1.1–2.0	0.4–0.9	0.2–0.6	0.0–0.2	0.0–0.3	5.3–7.1
White	%	92.8	3.8	1.9	0.7	0.5	0.1	0.2	7.2
(n=5,367)	CI	92.0–93.5	3.3–4.4	1.5–2.3	0.5–1.0	0.3–0.7	0.1–0.3	0.1–0.4	6.5–8.0
Black	%	89.5	6.7	1.5	0.8	0.7	0.0	0.8	10.5
(n=141)	CI	83.0–93.7	3.5–12.4	0.4–5.9	0.1–5.5	0.1–4.9	–	0.1–5.5	6.3–17.0
Hispanic	%	90.7	4.2	2.8	1.4	0.4	0.5	0.0	9.3
(n=227)	CI	86.0–94.0	2.2–7.9	1.2–6.1	0.5–4.3	0.1–3.1	0.1–3.4	–	6.0–14.0
8th	%	97.8	1.5	0.3	0.1	0.1	0.1	0.0	2.2
(n=2,385)	CI	97.1–98.3	1.1–2.1	0.1–0.6	0.0–0.4	0.0–0.4	0.0–0.4	0.0–0.3	1.7–2.9
10th	%	91.5	4.3	2.4	0.8	0.7	0.0	0.2	8.5
(n=2,221)	CI	90.3–92.6	3.5–5.2	1.8–3.1	0.5–1.3	0.4–1.2	0.0–0.3	0.1–0.5	7.4–9.7
12th	%	87.7	6.2	3.2	1.2	0.8	0.3	0.6	12.3
(n=1,813)	CI	86.1–89.2	5.2–7.4	2.5–4.2	0.8–1.8	0.5–1.4	0.1–0.7	0.3–1.0	10.8–13.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.18. Past Month Cigarette Use

		During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	91.2	3.6	1.9	0.6	0.8	0.6	1.4	8.8
(n=6,418)	CI	90.5–91.9	3.2–4.1	1.6–2.2	0.4–0.8	0.6–1.0	0.4–0.8	1.1–1.7	8.1–9.5
Male	%	91.1	3.9	2.0	0.5	0.7	0.6	1.2	8.9
(n=3,107)	CI	90.0–92.0	3.2–4.6	1.6–2.6	0.3–0.9	0.4–1.1	0.4–1.0	0.9–1.7	8.0–10.0
Female	%	91.6	3.3	1.7	0.7	0.8	0.5	1.4	8.4
(n=3,154)	CI	90.6–92.5	2.7–4.0	1.3–2.3	0.4–1.0	0.6–1.2	0.3–0.8	1.1–1.9	7.5–9.4
White	%	91.4	3.5	1.8	0.6	0.8	0.5	1.4	8.6
(n=5,360)	CI	90.6–92.1	3.1–4.1	1.5–2.2	0.4–0.8	0.6–1.1	0.3–0.7	1.1–1.8	7.9–9.4
Black	%	92.1	2.9	3.0	0.0	0.0	1.4	0.7	7.9
(n=143)	CI	86.1–95.6	1.1–7.6	1.1–7.8	–	–	0.3–5.5	0.1–4.6	4.4–13.9
Hispanic	%	92.9	2.7	0.5	1.3	0.9	0.5	1.3	7.1
(n=229)	CI	88.7–95.6	1.2–5.8	0.1–3.4	0.4–3.9	0.2–3.6	0.1–3.4	0.4–4.0	4.4–11.3
8th	%	94.5	2.8	1.0	0.4	0.3	0.3	0.7	5.5
(n=2,386)	CI	93.5–95.3	2.3–3.6	0.7–1.5	0.2–0.7	0.1–0.6	0.1–0.6	0.4–1.1	4.7–6.5
10th	%	90.0	3.7	2.1	0.8	0.7	0.8	1.9	10.0
(n=2,217)	CI	88.7–91.2	2.9–4.5	1.6–2.8	0.5–1.3	0.4–1.1	0.5–1.3	1.4–2.6	8.8–11.3
12th	%	88.9	4.3	2.5	0.7	1.5	0.5	1.6	11.1
(n=1,815)	CI	87.4–90.3	3.5–5.4	1.9–3.3	0.4–1.2	1.0–2.2	0.3–1.0	1.1–2.3	9.7–12.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	82.7	6.0	3.0	1.1	2.0	1.5	3.7	17.3
(n=6,402)	CI	81.7–83.6	5.4–6.6	2.6–3.5	0.9–1.4	1.6–2.3	1.3–1.9	3.3–4.2	16.4–18.3
Male	%	81.1	6.2	3.1	1.3	2.2	1.7	4.3	18.9
(n=3,094)	CI	79.7–82.5	5.4–7.1	2.5–3.8	1.0–1.8	1.8–2.8	1.3–2.3	3.7–5.1	17.5–20.3
Female	%	84.3	5.7	3.1	0.9	1.7	1.4	3.0	15.7
(n=3,153)	CI	83.0–85.6	4.9–6.5	2.5–3.7	0.6–1.3	1.3–2.2	1.0–1.8	2.5–3.7	14.4–17.0
White	%	82.8	5.9	3.0	1.1	2.0	1.6	3.5	17.2
(n=5,349)	CI	81.8–83.8	5.3–6.5	2.6–3.5	0.9–1.4	1.7–2.4	1.3–2.0	3.1–4.1	16.2–18.2
Black	%	80.7	3.5	4.2	2.9	2.1	0.7	5.9	19.3
(n=142)	CI	73.3–86.5	1.4–8.2	1.9–9.2	1.1–7.5	0.7–6.4	0.1–4.6	2.9–11.4	13.5–26.7
Hispanic	%	87.6	5.1	2.7	0.0	1.9	1.5	1.3	12.4
(n=228)	CI	82.5–91.3	2.9–8.9	1.2–5.9	–	0.7–5.0	0.5–4.5	0.4–3.9	8.7–17.5
8th	%	89.5	4.6	2.3	0.7	1.1	0.5	1.3	10.5
(n=2,376)	CI	88.2–90.7	3.8–5.5	1.7–3.0	0.4–1.1	0.8–1.7	0.3–0.9	1.0–1.9	9.3–11.8
10th	%	79.7	6.7	3.8	1.4	2.7	1.6	4.1	20.3
(n=2,213)	CI	78.0–81.3	5.8–7.8	3.1–4.7	1.0–2.0	2.1–3.5	1.1–2.2	3.4–5.0	18.7–22.0
12th	%	78.4	6.6	3.1	1.4	2.0	2.6	5.8	21.6
(n=1,813)	CI	76.5–80.3	5.5–7.8	2.4–4.0	0.9–2.0	1.5–2.8	2.0–3.5	4.8–7.0	19.7–23.5

Note: The confidence interval (CI) is 95 percent.

Table BH2.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	94.5	1.8	0.7	0.3	0.4	0.4	2.0	5.5
(n=6,424)	CI	93.9–95.1	1.5–2.1	0.5–0.9	0.2–0.5	0.3–0.6	0.3–0.6	1.6–2.3	4.9–6.1
Male	%	91.4	2.6	0.9	0.5	0.7	0.6	3.4	8.6
(n=3,107)	CI	90.3–92.3	2.1–3.2	0.6–1.3	0.3–0.8	0.4–1.0	0.4–1.0	2.8–4.1	7.7–9.7
Female	%	97.8	0.8	0.4	0.2	0.1	0.2	0.5	2.2
(n=3,161)	CI	97.3–98.3	0.5–1.2	0.2–0.7	0.1–0.4	0.0–0.3	0.1–0.4	0.3–0.9	1.7–2.7
White	%	94.3	1.7	0.7	0.3	0.5	0.4	2.1	5.7
(n=5,367)	CI	93.7–94.9	1.4–2.1	0.5–1.0	0.2–0.5	0.3–0.7	0.3–0.6	1.7–2.5	5.1–6.3
Black	%	94.4	2.1	1.3	0.0	0.0	0.7	1.5	5.6
(n=143)	CI	89.0–97.2	0.7–6.5	0.3–5.3	–	–	0.1–4.9	0.4–5.8	2.8–11.0
Hispanic	%	98.6	0.9	0.0	0.0	0.0	0.0	0.5	1.4
(n=227)	CI	95.8–99.6	0.2–3.4	–	–	–	–	0.1–3.4	0.4–4.2
8th	%	96.7	1.1	0.7	0.3	0.2	0.2	0.8	3.3
(n=2,387)	CI	95.9–97.3	0.7–1.6	0.4–1.1	0.2–0.7	0.1–0.5	0.1–0.5	0.5–1.2	2.7–4.1
10th	%	93.6	2.2	0.7	0.4	0.6	0.6	1.9	6.4
(n=2,219)	CI	92.5–94.5	1.7–2.9	0.4–1.2	0.2–0.7	0.3–1.0	0.4–1.1	1.4–2.6	5.5–7.5
12th	%	93.2	2.0	0.5	0.2	0.4	0.3	3.3	6.8
(n=1,818)	CI	91.9–94.3	1.5–2.8	0.3–1.0	0.1–0.6	0.2–0.8	0.1–0.7	2.6–4.2	5.7–8.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	91.1	3.1	1.6	0.8	1.1	0.8	1.6	8.9
(n=6,396)	CI	90.4–91.8	2.7–3.6	1.3–2.0	0.6–1.0	0.9–1.4	0.6–1.0	1.3–1.9	8.2–9.6
Male	%	91.2	3.2	1.6	0.5	1.2	0.6	1.7	8.8
(n=3,096)	CI	90.2–92.2	2.6–3.8	1.2–2.1	0.3–0.9	0.9–1.6	0.4–1.0	1.3–2.2	7.8–9.8
Female	%	91.2	3.1	1.5	1.0	1.0	0.9	1.3	8.8
(n=3,147)	CI	90.2–92.2	2.6–3.8	1.1–2.0	0.7–1.4	0.7–1.4	0.6–1.2	0.9–1.7	7.8–9.8
White	%	91.8	2.9	1.4	0.7	1.2	0.7	1.3	8.2
(n=5,351)	CI	91.0–92.5	2.5–3.4	1.1–1.7	0.5–0.9	0.9–1.5	0.5–1.0	1.1–1.7	7.5–9.0
Black	%	79.7	5.8	5.7	0.7	1.5	0.7	6.0	20.3
(n=141)	CI	72.1–85.7	2.9–11.3	2.8–11.0	0.1–4.6	0.4–5.9	0.1–4.6	3.0–11.6	14.3–27.9
Hispanic	%	91.2	5.6	1.4	0.9	0.4	0.4	0.0	8.8
(n=226)	CI	86.6–94.3	3.2–9.7	0.4–4.2	0.2–3.7	0.1–3.1	0.1–3.1	–	5.7–13.4
8th	%	95.6	1.5	1.1	0.4	0.5	0.3	0.5	4.4
(n=2,383)	CI	94.7–96.4	1.1–2.0	0.8–1.6	0.2–0.8	0.3–0.9	0.1–0.6	0.3–0.9	3.6–5.3
10th	%	89.4	3.7	1.8	0.8	1.5	0.9	1.9	10.6
(n=2,204)	CI	88.1–90.7	3.0–4.5	1.3–2.5	0.5–1.3	1.1–2.1	0.6–1.4	1.4–2.5	9.3–11.9
12th	%	87.9	4.3	1.9	1.0	1.3	1.2	2.4	12.1
(n=1,809)	CI	86.3–89.3	3.5–5.3	1.4–2.7	0.7–1.6	0.8–1.9	0.8–1.8	1.8–3.2	10.7–13.7

Note: The confidence interval (CI) is 95 percent.

Table BH2.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.5	0.2	0.1	0.0	0.1	0.0	0.1	0.5
(n=6,433)	CI	99.3–99.6	0.1–0.3	0.1–0.2	0.0–0.1	0.0–0.2	–	0.0–0.2	0.4–0.7
Male	%	99.5	0.1	0.1	0.0	0.1	0.0	0.1	0.5
(n=3,115)	CI	99.2–99.7	0.1–0.4	0.0–0.4	–	0.1–0.4	–	0.0–0.3	0.3–0.8
Female	%	99.6	0.2	0.1	0.0	0.1	0.0	0.0	0.4
(n=3,162)	CI	99.4–99.8	0.1–0.5	0.0–0.3	–	0.0–0.3	–	–	0.2–0.6
White	%	99.6	0.2	0.1	0.0	0.1	0.0	0.0	0.4
(n=5,376)	CI	99.4–99.8	0.1–0.3	0.1–0.3	0.0–0.1	0.0–0.2	–	0.0–0.1	0.2–0.6
Black	%	98.5	0.0	0.0	0.0	0.7	0.0	0.8	1.5
(n=143)	CI	94.2–99.6	–	–	–	0.1–4.9	–	0.1–5.4	0.4–5.8
Hispanic	%	99.6	0.0	0.0	0.0	0.4	0.0	0.0	0.4
(n=227)	CI	96.9–99.9	–	–	–	0.1–3.1	–	–	0.1–3.1
8th	%	99.9	0.0	0.0	0.0	0.0	0.0	0.0	0.1
(n=2,389)	CI	99.7–100.0	0.0–0.3	0.0–0.3	–	–	–	–	0.0–0.3
10th	%	99.4	0.3	0.1	0.0	0.1	0.0	0.1	0.6
(n=2,222)	CI	98.9–99.6	0.2–0.7	0.0–0.4	0.0–0.3	0.0–0.4	–	0.0–0.4	0.4–1.1
12th	%	99.1	0.2	0.2	0.0	0.2	0.0	0.2	0.9
(n=1,822)	CI	98.6–99.5	0.1–0.6	0.1–0.6	–	0.1–0.6	–	0.1–0.6	0.5–1.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.5	0.2	0.1	0.0	0.0	0.0	0.1	0.5
(n=6,433)	CI	99.3–99.7	0.1–0.3	0.0–0.2	0.0–0.1	0.0–0.1	–	0.1–0.3	0.3–0.7
Male	%	99.4	0.2	0.2	0.1	0.0	0.0	0.1	0.6
(n=3,115)	CI	99.1–99.6	0.1–0.5	0.1–0.4	0.0–0.3	–	–	0.0–0.4	0.4–0.9
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=3,162)	CI	99.5–99.9	0.0–0.3	0.0–0.2	0.0–0.2	0.0–0.2	–	–	0.1–0.5
White	%	99.6	0.2	0.1	0.1	0.0	0.0	0.1	0.4
(n=5,377)	CI	99.4–99.8	0.1–0.3	0.0–0.2	0.0–0.2	–	–	0.0–0.2	0.2–0.6
Black	%	98.5	0.0	0.7	0.0	0.0	0.0	0.8	1.5
(n=143)	CI	94.2–99.6	–	0.1–4.9	–	–	–	0.1–5.4	0.4–5.8
Hispanic	%	99.6	0.0	0.4	0.0	0.0	0.0	0.0	0.4
(n=227)	CI	96.9–99.9	–	0.1–3.1	–	–	–	–	0.1–3.1
8th	%	99.9	0.1	0.0	0.0	0.0	0.0	0.0	0.1
(n=2,389)	CI	99.6–100.0	0.0–0.3	–	–	–	–	0.0–0.3	0.0–0.4
10th	%	99.3	0.2	0.2	0.1	0.0	0.0	0.1	0.7
(n=2,223)	CI	98.9–99.6	0.1–0.5	0.1–0.5	0.0–0.4	–	–	0.0–0.4	0.4–1.1
12th	%	99.3	0.3	0.1	0.0	0.1	0.0	0.3	0.7
(n=1,821)	CI	98.8–99.6	0.1–0.7	0.0–0.4	–	0.0–0.4	–	0.1–0.7	0.4–1.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.4	0.1	0.0	0.1	0.0	0.1	0.8
(n=6,431)	CI	99.0–99.4	0.3–0.6	0.1–0.3	0.0–0.1	0.0–0.2	0.0–0.1	0.0–0.2	0.6–1.0
Male	%	99.1	0.6	0.1	0.1	0.1	0.0	0.1	0.9
(n=3,115)	CI	98.7–99.4	0.4–0.9	0.0–0.3	0.0–0.3	0.0–0.3	–	0.0–0.3	0.6–1.3
Female	%	99.6	0.2	0.2	0.0	0.0	0.0	0.0	0.4
(n=3,162)	CI	99.3–99.8	0.1–0.4	0.1–0.4	–	0.0–0.2	0.0–0.2	–	0.2–0.7
White	%	99.4	0.3	0.1	0.0	0.1	0.0	0.0	0.6
(n=5,376)	CI	99.2–99.6	0.2–0.5	0.1–0.3	0.0–0.2	0.0–0.2	–	0.0–0.1	0.4–0.8
Black	%	97.8	0.7	0.7	0.0	0.0	0.0	0.8	2.2
(n=143)	CI	93.4–99.3	0.1–4.9	0.1–4.9	–	–	–	0.1–5.4	0.7–6.6
Hispanic	%	99.1	0.4	0.0	0.0	0.4	0.0	0.0	0.9
(n=227)	CI	96.6–99.8	0.1–3.1	–	–	0.1–3.1	–	–	0.2–3.4
8th	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=2,389)	CI	99.5–99.9	0.0–0.4	0.0–0.3	0.0–0.3	–	–	–	0.1–0.5
10th	%	99.2	0.4	0.1	0.0	0.1	0.0	0.1	0.8
(n=2,222)	CI	98.7–99.5	0.2–0.8	0.0–0.4	–	0.0–0.4	–	0.0–0.4	0.5–1.3
12th	%	98.7	0.7	0.2	0.1	0.1	0.1	0.2	1.3
(n=1,820)	CI	98.0–99.1	0.4–1.2	0.1–0.6	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.5	0.9–2.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.6	0.1	0.1	0.0	0.1	0.0	0.1	0.4
(n=6,432)	CI	99.5–99.8	0.1–0.3	0.0–0.2	–	0.0–0.2	–	0.0–0.2	0.2–0.5
Male	%	99.7	0.1	0.1	0.0	0.1	0.0	0.0	0.3
(n=3,114)	CI	99.5–99.9	0.0–0.3	0.0–0.3	–	0.0–0.3	–	0.0–0.3	0.1–0.5
Female	%	99.8	0.1	0.1	0.0	0.1	0.0	0.0	0.2
(n=3,162)	CI	99.5–99.9	0.0–0.3	0.0–0.2	–	0.0–0.3	–	–	0.1–0.5
White	%	99.8	0.1	0.1	0.0	0.0	0.0	0.0	0.2
(n=5,376)	CI	99.7–99.9	0.0–0.2	0.0–0.2	–	0.0–0.1	–	–	0.1–0.3
Black	%	97.8	0.7	0.0	0.0	0.7	0.0	0.8	2.2
(n=143)	CI	93.4–99.3	0.1–4.9	–	–	0.1–4.9	–	0.1–5.4	0.7–6.6
Hispanic	%	99.6	0.0	0.0	0.0	0.4	0.0	0.0	0.4
(n=227)	CI	96.9–99.9	–	–	–	0.1–3.1	–	–	0.1–3.1
8th	%	99.9	0.0	0.1	0.0	0.0	0.0	0.0	0.1
(n=2,389)	CI	99.6–100.0	0.0–0.3	0.0–0.3	–	–	–	–	0.0–0.4
10th	%	99.6	0.2	0.1	0.0	0.1	0.0	0.0	0.4
(n=2,223)	CI	99.2–99.8	0.1–0.5	0.0–0.4	–	0.0–0.4	–	0.0–0.3	0.2–0.8
12th	%	99.5	0.2	0.1	0.0	0.1	0.0	0.2	0.5
(n=1,820)	CI	99.1–99.7	0.1–0.5	0.0–0.4	–	0.0–0.4	–	0.1–0.6	0.3–0.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.7	0.1	0.0	0.0	0.0	0.0	0.1	0.3
(n=6,431)	CI	99.5–99.8	0.0–0.2	0.0–0.2	0.0–0.1	0.0–0.1	0.0–0.1	0.0–0.2	0.2–0.5
Male	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=3,114)	CI	99.5–99.9	0.0–0.3	0.0–0.3	0.0–0.3	–	0.0–0.2	0.0–0.3	0.1–0.5
Female	%	99.8	0.0	0.0	0.0	0.1	0.0	0.0	0.2
(n=3,162)	CI	99.6–99.9	0.0–0.2	–	0.0–0.2	0.0–0.3	0.0–0.2	–	0.1–0.4
White	%	99.8	0.1	0.0	0.0	0.1	0.0	0.0	0.2
(n=5,375)	CI	99.6–99.9	0.0–0.2	0.0–0.2	0.0–0.2	0.0–0.2	–	–	0.1–0.4
Black	%	99.2	0.0	0.0	0.0	0.0	0.0	0.8	0.8
(n=143)	CI	94.6–99.9	–	–	–	–	–	0.1–5.4	0.1–5.4
Hispanic	%	99.6	0.0	0.0	0.0	0.0	0.4	0.0	0.4
(n=227)	CI	96.9–99.9	–	–	–	–	0.1–3.1	–	0.1–3.1
8th	%	99.9	0.0	0.0	0.0	0.1	0.0	0.0	0.1
(n=2,389)	CI	99.6–100.0	0.0–0.3	–	–	0.0–0.3	–	–	0.0–0.4
10th	%	99.7	0.1	0.0	0.0	0.0	0.0	0.0	0.3
(n=2,222)	CI	99.3–99.8	0.0–0.4	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.3	0.2–0.7
12th	%	99.5	0.1	0.1	0.1	0.0	0.1	0.2	0.5
(n=1,820)	CI	99.0–99.7	0.0–0.4	0.0–0.4	0.0–0.4	–	0.0–0.4	0.1–0.6	0.3–1.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you use ecstasy (MDMA)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.6	0.2	0.1	0.1	0.0	0.0	0.1	0.4
(n=6,433)	CI	99.4-99.7	0.1-0.3	0.0-0.2	0.0-0.2	-	0.0-0.1	0.0-0.2	0.3-0.6
Male	%	99.6	0.3	0.0	0.1	0.0	0.0	0.0	0.4
(n=3,115)	CI	99.3-99.8	0.1-0.5	0.0-0.2	0.0-0.3	-	-	0.0-0.3	0.2-0.7
Female	%	99.8	0.1	0.1	0.0	0.0	0.0	0.0	0.2
(n=3,163)	CI	99.6-99.9	0.0-0.3	0.0-0.2	0.0-0.2	-	0.0-0.2	-	0.1-0.4
White	%	99.7	0.2	0.1	0.1	0.0	0.0	0.0	0.3
(n=5,376)	CI	99.5-99.8	0.1-0.3	0.0-0.2	0.0-0.2	-	-	-	0.2-0.5
Black	%	99.2	0.0	0.0	0.0	0.0	0.0	0.8	0.8
(n=143)	CI	94.6-99.9	-	-	-	-	-	0.1-5.4	0.1-5.4
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=227)	CI	-	-	-	-	-	-	-	-
8th	%	99.8	0.0	0.1	0.0	0.0	0.0	0.0	0.2
(n=2,390)	CI	99.6-99.9	0.0-0.3	0.0-0.3	-	-	0.0-0.3	-	0.1-0.4
10th	%	99.7	0.1	0.0	0.1	0.0	0.0	0.0	0.3
(n=2,222)	CI	99.3-99.8	0.0-0.4	0.0-0.3	0.0-0.4	-	-	0.0-0.3	0.2-0.7
12th	%	99.3	0.3	0.1	0.1	0.0	0.0	0.2	0.7
(n=1,821)	CI	98.8-99.6	0.1-0.7	0.0-0.4	0.0-0.4	-	-	0.1-0.6	0.4-1.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you use methamphetamines (crank, etc.)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.6	0.1	0.1	0.0	0.0	0.0	0.1	0.4
(n=6,436)	CI	99.4-99.8	0.1-0.3	0.0-0.2	0.0-0.1	0.0-0.1	0.0-0.1	0.0-0.2	0.2-0.6
Male	%	99.6	0.2	0.1	0.0	0.1	0.0	0.0	0.4
(n=3,117)	CI	99.3-99.8	0.1-0.5	0.0-0.3	-	0.0-0.3	-	0.0-0.2	0.2-0.7
Female	%	99.9	0.0	0.0	0.0	0.0	0.0	0.0	0.1
(n=3,162)	CI	99.7-100.0	0.0-0.2	0.0-0.2	-	-	0.0-0.2	-	0.0-0.3
White	%	99.8	0.1	0.1	0.0	0.0	0.0	0.0	0.2
(n=5,377)	CI	99.6-99.9	0.1-0.3	0.0-0.2	-	0.0-0.1	0.0-0.1	0.0-0.1	0.1-0.4
Black	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=143)	CI	-	-	-	-	-	-	-	-
Hispanic	%	99.6	0.0	0.0	0.0	0.4	0.0	0.0	0.4
(n=229)	CI	97.0-99.9	-	-	-	0.1-3.0	-	-	0.1-3.0
8th	%	99.9	0.0	0.0	0.0	0.0	0.0	0.0	0.1
(n=2,388)	CI	99.6-100.0	0.0-0.3	0.0-0.3	-	-	-	0.0-0.3	0.0-0.4
10th	%	99.5	0.1	0.1	0.0	0.1	0.0	0.1	0.5
(n=2,222)	CI	99.1-99.7	0.0-0.4	0.0-0.4	-	0.0-0.4	0.0-0.3	0.0-0.4	0.3-0.9
12th	%	99.5	0.3	0.1	0.1	0.0	0.0	0.1	0.5
(n=1,826)	CI	99.1-99.7	0.1-0.7	0.0-0.4	0.0-0.4	-	-	0.0-0.4	0.3-0.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	19.7	20.4	15.8	44.2	59.9
(n=2,823)	CI	18.2–21.2	18.9–21.9	14.5–17.2	42.4–46.0	58.1–61.7
Male	%	17.5	18.7	16.9	46.9	63.8
(n=1,323)	CI	15.5–19.7	16.6–20.9	15.0–19.0	44.2–49.6	61.2–66.4
Female	%	20.2	22.3	15.1	42.3	57.5
(n=1,416)	CI	18.2–22.4	20.2–24.5	13.4–17.1	39.8–44.9	54.9–60.0
White	%	19.1	21.0	16.3	43.5	59.8
(n=2,345)	CI	17.6–20.8	19.4–22.7	14.9–17.9	41.5–45.5	57.8–61.8
Black	%	20.3	13.5	11.8	54.4	66.2
(n=69)	CI	12.2–31.7	7.1–24.3	5.9–22.2	42.3–66.0	54.0–76.6
Hispanic	%	14.1	13.0	18.6	54.2	72.9
(n=93)	CI	8.3–23.0	7.5–21.8	11.8–28.1	43.8–64.3	62.7–81.1
8th	%	17.8	16.8	11.4	54.0	65.4
(n=1,058)	CI	15.6–20.2	14.7–19.2	9.7–13.5	51.0–57.0	62.5–68.2
10th	%	19.2	21.3	17.7	41.8	59.5
(n=982)	CI	16.9–21.8	18.8–24.0	15.5–20.2	38.7–44.9	56.4–62.5
12th	%	22.3	23.4	18.4	35.9	54.3
(n=783)	CI	19.5–25.3	20.6–26.5	15.9–21.3	32.6–39.3	50.8–57.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	23.5	34.2	16.3	25.9	42.2
(n=3,795)	CI	22.2–24.9	32.7–35.8	15.2–17.6	24.5–27.3	40.7–43.8
Male	%	22.5	33.7	16.9	26.9	43.8
(n=1,798)	CI	20.6–24.5	31.5–35.9	15.3–18.8	24.9–29.0	41.6–46.1
Female	%	23.8	35.5	16.0	24.7	40.7
(n=1,888)	CI	21.9–25.8	33.3–37.7	14.4–17.7	22.8–26.7	38.5–42.9
White	%	23.1	34.8	16.4	25.7	42.1
(n=3,152)	CI	21.7–24.7	33.1–36.5	15.1–17.7	24.2–27.2	40.3–43.8
Black	%	24.2	28.5	5.4	41.9	47.3
(n=88)	CI	16.2–34.5	19.9–39.0	2.2–12.6	31.9–52.6	36.9–57.9
Hispanic	%	19.1	34.5	19.7	26.7	46.5
(n=134)	CI	13.2–26.8	26.8–43.0	13.7–27.5	19.9–34.9	38.1–55.1
8th	%	16.7	27.5	16.8	39.0	55.8
(n=1,239)	CI	14.7–18.9	25.1–30.1	14.8–19.0	36.3–41.7	53.0–58.5
10th	%	22.7	37.2	18.0	22.1	40.1
(n=1,357)	CI	20.5–25.0	34.7–39.8	16.0–20.1	20.0–24.4	37.5–42.7
12th	%	30.3	37.0	14.3	18.4	32.7
(n=1,199)	CI	27.7–32.9	34.3–39.8	12.4–16.4	16.3–20.7	30.1–35.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	38.1	24.5	12.0	25.4	37.4
(n=3,872)	CI	36.5–39.6	23.2–25.9	11.1–13.1	24.1–26.8	35.9–39.0
Male	%	40.4	22.6	13.1	23.9	37.0
(n=1,891)	CI	38.2–42.7	20.8–24.5	11.7–14.7	22.0–25.8	34.8–39.2
Female	%	35.0	26.8	11.1	27.1	38.2
(n=1,877)	CI	32.9–37.3	24.8–28.8	9.8–12.6	25.1–29.1	36.0–40.4
White	%	37.9	25.3	12.1	24.7	36.8
(n=3,254)	CI	36.2–39.6	23.8–26.8	11.0–13.3	23.3–26.3	35.2–38.5
Black	%	37.9	17.9	10.1	34.0	44.1
(n=87)	CI	28.1–48.8	11.2–27.6	5.3–18.5	24.7–44.8	33.9–54.9
Hispanic	%	28.2	23.4	10.7	37.8	48.4
(n=120)	CI	20.7–37.1	16.6–31.9	6.2–17.6	29.4–46.9	39.5–57.5
8th	%	23.7	21.2	15.0	40.0	55.0
(n=1,276)	CI	21.5–26.2	19.1–23.6	13.1–17.0	37.4–42.8	52.3–57.7
10th	%	34.4	28.8	13.0	23.9	36.9
(n=1,359)	CI	31.9–36.9	26.4–31.2	11.3–14.8	21.7–26.3	34.3–39.5
12th	%	54.0	23.1	8.6	14.3	22.9
(n=1,237)	CI	51.2–56.7	20.9–25.6	7.2–10.3	12.4–16.3	20.6–25.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	23.1	22.0	13.2	41.6	54.8
(n=3,191)	CI	21.7–24.6	20.6–23.5	12.0–14.4	39.9–43.4	53.1–56.6
Male	%	23.1	21.6	14.1	41.2	55.3
(n=1,498)	CI	21.0–25.3	19.5–23.8	12.4–16.0	38.7–43.7	52.8–57.8
Female	%	22.5	22.3	12.5	42.7	55.2
(n=1,601)	CI	20.5–24.6	20.3–24.4	11.0–14.2	40.3–45.1	52.7–57.6
White	%	22.5	21.9	14.3	41.4	55.7
(n=2,627)	CI	20.9–24.1	20.3–23.5	13.0–15.7	39.5–43.3	53.8–57.6
Black	%	31.6	23.9	3.3	41.1	44.5
(n=84)	CI	22.3–42.6	15.8–34.5	1.0–10.0	31.0–52.1	34.0–55.4
Hispanic	%	17.0	23.3	9.5	50.2	59.7
(n=116)	CI	11.0–25.2	16.3–32.2	5.3–16.5	41.0–59.3	50.3–68.4
8th	%	10.8	11.2	11.8	66.1	77.9
(n=1,051)	CI	9.1–12.9	9.5–13.3	10.0–13.9	63.2–68.9	75.3–80.3
10th	%	25.0	24.7	14.3	36.0	50.3
(n=1,161)	CI	22.6–27.6	22.3–27.3	12.4–16.4	33.3–38.8	47.4–53.2
12th	%	32.2	28.9	13.2	25.7	38.9
(n=979)	CI	29.3–35.2	26.2–31.8	11.3–15.5	23.0–28.5	35.9–42.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	1.1	98.9
(n=6,419)	CI	0.9–1.4	98.6–99.1
Male	%	1.0	99.0
(n=3,113)	CI	0.7–1.4	98.6–99.3
Female	%	1.2	98.8
(n=3,153)	CI	0.8–1.6	98.4–99.2
White	%	1.0	99.0
(n=5,365)	CI	0.8–1.3	98.7–99.2
Black	%	2.2	97.8
(n=141)	CI	0.7–6.7	93.3–99.3
Hispanic	%	1.7	98.3
(n=229)	CI	0.6–4.5	95.5–99.4
8th	%	0.8	99.2
(n=2,387)	CI	0.5–1.3	98.7–99.5
10th	%	1.5	98.5
(n=2,212)	CI	1.1–2.1	97.9–98.9
12th	%	1.1	98.9
(n=1,820)	CI	0.7–1.7	98.3–99.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	12.5	87.5
(n=6,373)	CI	11.7–13.3	86.7–88.3
Male	%	12.2	87.8
(n=3,076)	CI	11.1–13.4	86.6–88.9
Female	%	12.7	87.3
(n=3,146)	CI	11.6–13.9	86.1–88.4
White	%	12.5	87.5
(n=5,338)	CI	11.6–13.4	86.6–88.4
Black	%	11.4	88.6
(n=138)	CI	7.0–18.2	81.8–93.0
Hispanic	%	12.2	87.8
(n=223)	CI	8.4–17.3	82.7–91.6
8th	%	4.9	95.1
(n=2,366)	CI	4.1–5.8	94.2–95.9
10th	%	14.7	85.3
(n=2,206)	CI	13.3–16.3	83.7–86.7
12th	%	18.3	81.7
(n=1,801)	CI	16.6–20.1	79.9–83.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=73)	Male (n=31)	Female (n=37)	White (n=53)	Black (n=3)	Hispanic (n=4)	8th (n=20)	10th (n=33)	12th (n=20)
Got a prescription from one doctor	%	8.3	6.5	7.8	7.2	0.0	0.0	10.0	6.1	10.0
	CI	3.7–17.5	1.5–24.2	2.4–22.7	2.6–18.1	–	–	2.2–35.5	1.4–22.6	2.2–35.5
Got a prescription from more than one doctor	%	9.2	12.6	7.5	9.1	0.0	0.0	25.0	3.0	5.0
	CI	4.4–18.4	4.5–30.6	2.3–21.9	3.7–20.7	–	–	9.9–50.3	0.4–20.3	0.6–32.3
Bought on the Internet	%	8.5	16.7	0.0	9.8	0.0	25.4	5.0	6.1	15.0
	CI	3.8–18.0	6.8–35.7	–	4.0–22.0	–	0.5–95.9	0.6–32.3	1.4–22.6	4.4–40.4
Bought from a friend, relative, or someone you know	%	32.7	31.7	32.7	35.6	64.0	0.0	30.0	39.4	25.0
	CI	22.7–44.5	17.3–50.8	19.1–50.1	23.6–49.7	0.3–99.9	–	13.1–55.0	23.7–57.6	9.9–50.3
Got from a friend, relative, or someone you know for free	%	49.0	58.1	45.2	50.8	32.0	49.2	50.0	54.5	40.0
	CI	37.5–60.6	39.4–74.7	29.5–62.0	37.2–64.3	0.1–99.7	2.4–97.5	27.7–72.3	36.8–71.2	20.0–64.0
Took from a friend, relative, or someone you know without asking	%	29.0	16.9	43.1	30.1	36.0	25.4	30.0	24.2	35.0
	CI	19.5–40.7	6.9–36.0	27.6–60.0	19.0–44.2	0.1–99.7	0.5–95.9	13.1–55.0	12.1–42.6	16.4–59.6
Took from someone's house or workplace without asking	%	9.4	3.5	13.0	11.1	0.0	25.4	10.0	12.1	5.0
	CI	4.5–18.8	0.4–23.3	5.3–28.7	4.9–23.1	–	0.5–95.9	2.2–35.5	4.4–29.4	0.6–32.3
Got it at a party	%	22.3	26.8	18.9	22.7	64.0	0.0	5.0	30.3	25.0
	CI	14.0–33.6	13.5–46.2	8.9–35.5	13.1–36.4	0.3–99.9	–	0.6–32.3	16.6–48.8	9.9–50.3
Got it some other way	%	32.1	37.0	29.2	34.4	32.0	25.4	15.0	36.4	40.0
	CI	22.1–43.9	21.2–56.2	16.4–46.4	22.5–48.6	0.1–99.7	0.5–95.9	4.4–40.4	21.3–54.7	20.0–64.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=770)	Male (n=363)	Female (n=388)	White (n=645)	Black (n=15)	Hispanic (n=26)	8th (n=116)	10th (n=325)	12th (n=329)
I bought it in a store	%	3.8	5.6	2.4	3.9	0.0	11.8	2.6	3.1	4.8
	CI	2.7–5.5	3.6–8.5	1.2–4.5	2.7–5.7	–	3.6–32.8	0.8–7.8	1.7–5.6	3.0–7.8
I bought it in a gas station	%	5.1	7.0	3.0	5.0	0.0	8.2	2.6	3.4	7.3
	CI	3.7–6.9	4.8–10.2	1.6–5.3	3.6–7.1	–	1.9–29.5	0.8–7.8	1.9–6.0	4.9–10.6
I bought it in a restaurant	%	0.5	0.8	0.3	0.3	0.0	0.0	1.7	0.3	0.3
	CI	0.2–1.3	0.2–2.3	0.0–2.0	0.1–1.1	–	–	0.4–6.8	0.0–2.2	0.0–2.1
I bought it in a bar or club	%	2.4	2.5	2.4	2.0	7.1	11.8	2.6	1.5	3.0
	CI	1.5–3.7	1.3–4.7	1.2–4.6	1.2–3.5	0.8–43.2	3.6–32.8	0.8–7.8	0.6–3.7	1.6–5.6
I bought it at a public event	%	2.7	3.3	2.1	2.2	0.0	15.5	3.4	1.8	3.3
	CI	1.8–4.2	1.9–5.8	1.0–4.1	1.3–3.7	–	5.5–36.5	1.3–8.9	0.8–4.1	1.8–5.9
I gave someone money to buy it for me	%	27.2	30.8	24.2	28.5	14.2	31.4	13.8	27.1	31.2
	CI	24.1–30.5	26.3–35.8	20.2–28.8	25.1–32.1	3.0–47.1	15.7–52.8	8.6–21.5	22.5–32.2	26.4–36.4
I got it from a sibling (under 21)	%	4.8	5.2	4.8	4.5	0.0	8.2	11.2	3.4	4.2
	CI	3.5–6.6	3.3–8.0	3.0–7.4	3.2–6.4	–	1.9–29.5	6.6–18.5	1.9–6.0	2.5–7.1
I got it from a friend (under 21)	%	21.3	18.6	24.1	22.0	13.4	19.1	16.4	23.4	20.9
	CI	18.5–24.3	14.9–22.9	20.1–28.6	18.9–25.3	2.8–45.6	7.6–40.2	10.6–24.4	19.1–28.3	16.8–25.7
I got it at a party	%	45.0	49.3	41.7	43.4	60.2	73.6	29.3	44.3	50.3
	CI	41.5–48.6	44.1–54.5	36.8–46.7	39.6–47.3	31.9–83.0	52.4–87.6	21.7–38.4	39.0–49.8	44.9–55.7
I took it from a store or family member	%	14.3	13.6	14.9	13.6	13.4	19.1	19.8	15.7	11.5
	CI	12.0–17.0	10.5–17.6	11.7–18.8	11.2–16.5	2.8–45.6	7.6–40.2	13.5–28.2	12.1–20.1	8.5–15.5
I took it home without permission	%	19.6	16.6	22.0	18.9	6.3	26.8	32.8	23.7	12.1
	CI	16.9–22.5	13.2–20.8	18.2–26.4	16.1–22.1	0.7–40.1	12.6–48.2	24.7–41.9	19.4–28.6	9.0–16.1
I got it from my parents with permission	%	21.5	16.9	26.1	22.6	6.3	7.7	25.9	22.2	19.7
	CI	18.8–24.6	13.4–21.1	22.0–30.8	19.6–26.0	0.7–40.1	1.7–28.3	18.6–34.7	17.9–27.0	15.7–24.4
I got it from an older sibling (over 21)	%	12.0	9.8	14.0	11.8	20.5	8.2	12.9	13.2	10.6
	CI	9.9–14.5	7.2–13.4	10.9–17.9	9.6–14.6	5.8–51.9	1.9–29.5	7.9–20.5	9.9–17.4	7.7–14.4
I got it from an older friend (over 21)	%	26.3	25.1	27.3	26.5	13.4	22.7	13.8	25.5	30.6
	CI	23.3–29.5	20.8–29.8	23.1–32.0	23.2–30.1	2.8–45.6	9.9–43.9	8.6–21.5	21.1–30.6	25.8–35.8
I got it from some other adult	%	19.7	17.6	21.6	19.7	33.8	15.0	14.7	22.8	18.5
	CI	17.0–22.7	14.0–21.9	17.7–25.9	16.8–23.0	13.1–63.4	5.3–35.6	9.2–22.5	18.5–27.7	14.6–23.1
I got it some other way	%	22.0	25.4	17.5	21.1	26.4	23.6	26.7	20.9	21.5
	CI	19.2–25.1	21.2–30.2	14.1–21.6	18.1–24.5	8.9–56.8	10.4–45.2	19.4–35.6	16.8–25.7	17.4–26.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?				
		I bought or tried to buy alcohol	Yes, asked for proof of age	No, not asked for proof of age
Total	%	3.0	29.8	70.2
(n=153)	CI	2.5–3.4	23.0–37.7	62.3–77.0
Male	%	3.6	27.7	72.3
(n=88)	CI	2.9–4.4	19.2–38.2	61.8–80.8
Female	%	1.8	25.1	74.9
(n=46)	CI	1.3–2.3	14.3–40.4	59.6–85.7
White	%	2.5	26.0	74.0
(n=111)	CI	2.1–3.0	18.5–35.2	64.8–81.5
Black	%	6.3	50.5	49.5
(n=6)	CI	3.1–12.2	9.3–91.1	8.9–90.7
Hispanic	%	5.2	36.4	63.6
(n=8)	CI	2.8–9.5	8.2–78.5	21.5–91.8
8th	%	1.7	15.2	84.8
(n=33)	CI	1.2–2.3	6.1–32.8	67.2–93.9
10th	%	3.2	22.8	77.2
(n=57)	CI	2.5–4.0	13.5–35.9	64.1–86.5
12th	%	4.1	41.5	58.5
(n=63)	CI	3.3–5.2	30.0–54.1	45.9–70.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?					
		I bought or tried to buy alcohol	No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	3.0	38.2	32.6	29.2
(n=153)	CI	2.5–3.4	31.0–46.0	25.7–40.3	22.7–36.7
Male	%	3.6	35.8	40.7	23.5
(n=88)	CI	2.9–4.4	26.6–46.1	31.0–51.2	15.9–33.4
Female	%	1.8	53.3	16.4	30.3
(n=46)	CI	1.3–2.3	38.9–67.1	8.2–30.1	18.8–45.0
White	%	2.5	37.4	32.5	30.1
(n=111)	CI	2.1–3.0	29.0–46.7	24.4–41.7	22.3–39.2
Black	%	6.3	51.2	12.6	36.3
(n=6)	CI	3.1–12.2	14.9–86.3	1.0–68.2	8.2–78.3
Hispanic	%	5.2	33.1	23.0	43.9
(n=8)	CI	2.8–9.5	8.0–73.7	4.0–68.0	13.1–80.2
8th	%	1.7	50.0	13.9	36.1
(n=33)	CI	1.2–2.3	33.5–66.5	5.6–30.3	21.7–53.6
10th	%	3.2	37.7	32.8	29.5
(n=57)	CI	2.5–4.0	26.2–50.8	22.0–45.8	19.2–42.4
12th	%	4.1	33.3	40.9	25.8
(n=63)	CI	3.3–5.2	22.8–45.8	29.5–53.4	16.5–37.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.39. Ever Ridden with a Prescription Drugs Impaired Driver

		Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		
		No	Yes	Not sure
Total	%	83.5	7.4	9.1
(n=6,168)	CI	82.6–84.4	6.8–8.1	8.4–9.8
Male	%	85.7	5.9	8.5
(n=2,975)	CI	84.4–86.9	5.1–6.8	7.5–9.5
Female	%	82.4	8.2	9.4
(n=3,042)	CI	81.0–83.7	7.3–9.3	8.4–10.5
White	%	84.0	7.0	9.0
(n=5,172)	CI	83.0–85.0	6.3–7.7	8.2–9.8
Black	%	80.1	9.5	10.4
(n=136)	CI	72.4–86.1	5.6–15.8	6.2–16.9
Hispanic	%	88.0	4.6	7.4
(n=209)	CI	82.9–91.7	2.5–8.4	4.6–11.8
8th	%	81.3	7.1	11.6
(n=2,282)	CI	79.6–82.8	6.2–8.3	10.3–12.9
10th	%	83.4	7.8	8.8
(n=2,136)	CI	81.8–84.9	6.7–9.0	7.7–10.1
12th	%	86.1	7.2	6.6
(n=1,750)	CI	84.4–87.7	6.1–8.6	5.5–7.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.40. Frequency Riding with a Prescription Drugs Impaired Driver

		During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?					
		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	96.6	1.1	1.0	0.3	1.1	3.4
(n=6,295)	CI	96.1–97.0	0.9–1.4	0.8–1.3	0.2–0.4	0.8–1.3	3.0–3.9
Male	%	97.3	0.7	0.8	0.2	1.0	2.7
(n=3,044)	CI	96.7–97.9	0.5–1.1	0.5–1.2	0.1–0.4	0.7–1.4	2.1–3.3
Female	%	96.5	1.4	1.1	0.4	0.7	3.5
(n=3,096)	CI	95.7–97.0	1.0–1.9	0.8–1.5	0.2–0.7	0.5–1.1	3.0–4.3
White	%	96.9	1.1	0.9	0.3	0.8	3.1
(n=5,268)	CI	96.4–97.4	0.9–1.5	0.6–1.1	0.2–0.4	0.6–1.1	2.6–3.6
Black	%	92.9	0.7	4.3	0.7	1.4	7.1
(n=137)	CI	87.1–96.2	0.1–4.8	1.9–9.4	0.1–5.1	0.3–5.5	3.8–12.9
Hispanic	%	96.9	0.9	0.9	0.0	1.3	3.1
(n=220)	CI	93.7–98.5	0.2–3.4	0.2–3.5	–	0.4–4.0	1.5–6.3
8th	%	96.9	1.0	1.0	0.3	0.8	3.1
(n=2,327)	CI	96.1–97.5	0.7–1.5	0.7–1.5	0.1–0.6	0.5–1.2	2.5–3.9
10th	%	96.2	1.1	1.1	0.4	1.1	3.8
(n=2,179)	CI	95.3–96.9	0.7–1.6	0.8–1.7	0.2–0.8	0.8–1.7	3.1–4.7
12th	%	96.7	1.2	0.8	0.1	1.3	3.3
(n=1,789)	CI	95.7–97.4	0.8–1.8	0.5–1.3	0.0–0.4	0.9–1.9	2.6–4.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	72.6	19.0	8.3
(n=6,283)	CI	71.5–73.7	18.1–20.0	7.7–9.0
Male	%	74.3	17.5	8.2
(n=3,036)	CI	72.7–75.8	16.2–18.9	7.3–9.3
Female	%	71.8	20.0	8.2
(n=3,092)	CI	70.2–73.3	18.6–21.5	7.3–9.2
White	%	73.5	18.3	8.2
(n=5,266)	CI	72.3–74.6	17.3–19.4	7.5–9.0
Black	%	65.1	24.8	10.1
(n=137)	CI	56.6–72.8	18.1–32.9	6.0–16.4
Hispanic	%	75.4	17.7	6.9
(n=217)	CI	69.2–80.7	13.2–23.4	4.2–11.1
8th	%	72.4	16.2	11.4
(n=2,322)	CI	70.6–74.2	14.7–17.7	10.1–12.7
10th	%	72.3	20.1	7.6
(n=2,177)	CI	70.4–74.2	18.5–21.8	6.5–8.8
12th	%	73.2	20.9	5.9
(n=1,784)	CI	71.1–75.2	19.1–22.9	4.9–7.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	91.0	3.6	3.0	0.6	1.7	9.0
(n=6,292)	CI	90.3–91.7	3.2–4.1	2.6–3.5	0.4–0.8	1.4–2.1	8.3–9.7
Male	%	92.1	3.3	2.5	0.7	1.5	7.9
(n=3,045)	CI	91.0–93.0	2.7–4.0	2.0–3.1	0.4–1.0	1.1–2.0	7.0–9.0
Female	%	90.4	4.1	3.5	0.5	1.5	9.6
(n=3,093)	CI	89.3–91.4	3.4–4.8	2.9–4.2	0.3–0.9	1.1–2.0	8.6–10.7
White	%	91.5	3.5	3.0	0.6	1.4	8.5
(n=5,272)	CI	90.7–92.3	3.1–4.1	2.5–3.5	0.4–0.8	1.1–1.8	7.7–9.3
Black	%	88.9	3.2	5.0	0.7	2.2	11.1
(n=137)	CI	82.3–93.3	1.2–8.2	2.4–10.3	0.1–5.1	0.7–6.6	6.7–17.7
Hispanic	%	88.9	5.0	4.0	0.5	1.7	11.1
(n=218)	CI	84.0–92.4	2.8–8.8	2.1–7.5	0.1–3.2	0.6–4.6	7.6–16.0
8th	%	92.8	2.9	2.5	0.4	1.4	7.2
(n=2,321)	CI	91.7–93.8	2.3–3.7	1.9–3.2	0.2–0.7	1.0–2.0	6.2–8.3
10th	%	89.8	3.8	3.8	0.7	1.9	10.2
(n=2,184)	CI	88.4–91.0	3.1–4.7	3.0–4.6	0.4–1.2	1.4–2.6	9.0–11.6
12th	%	90.5	4.1	2.8	0.7	1.8	9.5
(n=1,787)	CI	89.1–91.8	3.3–5.2	2.2–3.7	0.4–1.2	1.3–2.5	8.2–10.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.4	1.6	4.9	92.1	97.0
(n=6,277)	CI	1.1–1.7	1.3–1.9	4.4–5.5	91.4–92.7	96.6–97.4
Male	%	1.3	1.2	5.0	92.5	97.5
(n=3,029)	CI	0.9–1.8	0.9–1.7	4.3–5.8	91.5–93.4	96.8–98.0
Female	%	1.1	1.7	4.8	92.5	97.2
(n=3,094)	CI	0.8–1.5	1.3–2.2	4.1–5.6	91.5–93.3	96.6–97.7
White	%	1.1	1.5	5.1	92.2	97.3
(n=5,255)	CI	0.9–1.5	1.2–1.9	4.5–5.7	91.5–92.9	96.9–97.7
Black	%	5.0	0.7	4.7	89.7	94.3
(n=137)	CI	2.4–10.2	0.1–4.8	2.1–10.1	83.2–93.9	89.0–97.2
Hispanic	%	1.4	1.8	5.1	91.8	96.8
(n=217)	CI	0.4–4.3	0.7–4.6	2.8–8.9	87.3–94.8	93.5–98.5
8th	%	1.4	1.6	4.2	92.8	97.1
(n=2,317)	CI	1.0–1.9	1.1–2.1	3.5–5.1	91.7–93.8	96.3–97.7
10th	%	1.3	1.5	5.0	92.2	97.2
(n=2,173)	CI	0.9–1.9	1.0–2.1	4.2–6.0	91.0–93.3	96.5–97.9
12th	%	1.5	1.7	5.6	91.2	96.8
(n=1,787)	CI	1.0–2.2	1.2–2.5	4.6–6.7	89.8–92.4	95.8–97.5

Note: The confidence interval (CI) is 95 percent.

Table BH2.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.0	4.5	10.5	83.0	93.5
(n=6,302)	CI	1.7–2.4	4.0–5.0	9.8–11.3	82.1–83.9	92.9–94.1
Male	%	2.0	4.7	11.2	82.1	93.3
(n=3,042)	CI	1.5–2.6	4.0–5.6	10.1–12.4	80.7–83.4	92.3–94.1
Female	%	1.6	4.2	9.9	84.4	94.3
(n=3,104)	CI	1.2–2.1	3.5–4.9	8.9–11.0	83.1–85.6	93.4–95.0
White	%	1.7	4.4	10.4	83.5	93.9
(n=5,274)	CI	1.4–2.1	3.8–5.0	9.6–11.3	82.5–84.5	93.2–94.5
Black	%	4.9	1.4	10.4	83.2	93.7
(n=137)	CI	2.3–10.1	0.3–5.5	6.2–17.0	75.9–88.7	88.2–96.7
Hispanic	%	3.7	3.8	11.0	81.6	92.6
(n=219)	CI	1.8–7.2	1.9–7.4	7.5–16.0	75.8–86.2	88.2–95.4
8th	%	1.2	3.1	7.7	87.9	95.7
(n=2,328)	CI	0.8–1.7	2.5–3.9	6.7–8.9	86.5–89.2	94.8–96.4
10th	%	1.9	3.9	10.0	84.1	94.1
(n=2,183)	CI	1.4–2.6	3.2–4.8	8.8–11.4	82.5–85.6	93.1–95.1
12th	%	3.0	6.5	14.1	76.5	90.6
(n=1,791)	CI	2.3–3.8	5.4–7.7	12.6–15.8	74.5–78.4	89.1–91.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.45. Parents Feel Regularly Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.0	7.8	12.2	75.0	87.1
(n=6,295)	CI	4.5–5.6	7.2–8.5	11.4–13.0	73.9–76.0	86.3–88.0
Male	%	4.8	7.3	12.9	75.1	87.9
(n=3,041)	CI	4.0–5.6	6.4–8.3	11.7–14.1	73.5–76.6	86.7–89.1
Female	%	4.9	8.4	11.4	75.3	86.7
(n=3,100)	CI	4.2–5.7	7.5–9.4	10.3–12.6	73.7–76.8	85.5–87.9
White	%	4.7	7.8	12.2	75.4	87.6
(n=5,271)	CI	4.1–5.3	7.1–8.5	11.3–13.1	74.2–76.5	86.6–88.4
Black	%	8.0	7.0	11.7	73.3	85.0
(n=137)	CI	4.5–14.1	3.6–12.9	7.2–18.4	65.1–80.2	77.8–90.2
Hispanic	%	5.1	6.0	12.3	76.6	88.9
(n=216)	CI	2.8–9.1	3.5–10.0	8.6–17.5	70.4–81.8	83.9–92.5
8th	%	3.1	5.2	10.0	81.7	91.7
(n=2,326)	CI	2.5–3.9	4.4–6.2	8.9–11.3	80.0–83.2	90.5–92.7
10th	%	4.4	7.8	11.7	76.0	87.8
(n=2,181)	CI	3.6–5.3	6.8–9.0	10.4–13.2	74.2–77.8	86.3–89.1
12th	%	7.8	10.7	15.1	66.5	81.6
(n=1,788)	CI	6.6–9.1	9.3–12.2	13.5–16.8	64.3–68.7	79.7–83.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.46. Parents Feel Smoking Tobacco is Wrong

		How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.5	4.5	10.2	82.9	93.1
(n=6,287)	CI	2.1–2.9	4.0–5.0	9.4–10.9	81.9–83.8	92.4–93.7
Male	%	2.7	5.3	10.3	81.7	92.0
(n=3,039)	CI	2.2–3.4	4.5–6.1	9.3–11.5	80.3–83.1	91.0–93.0
Female	%	1.8	3.7	9.8	84.7	94.5
(n=3,094)	CI	1.4–2.3	3.1–4.4	8.8–10.9	83.4–86.0	93.7–95.3
White	%	2.3	4.6	10.5	82.6	93.1
(n=5,264)	CI	1.9–2.8	4.0–5.2	9.7–11.4	81.6–83.6	92.4–93.8
Black	%	3.6	5.0	11.1	80.3	91.4
(n=137)	CI	1.5–8.4	2.4–10.3	6.7–17.7	72.7–86.2	85.4–95.1
Hispanic	%	1.8	2.6	6.9	88.6	95.6
(n=218)	CI	0.7–4.8	1.2–5.8	4.2–11.3	83.6–92.2	91.9–97.6
8th	%	1.4	2.0	7.3	89.2	96.6
(n=2,323)	CI	1.0–2.0	1.5–2.7	6.3–8.4	87.9–90.4	95.7–97.2
10th	%	1.9	3.8	8.8	85.5	94.3
(n=2,177)	CI	1.4–2.6	3.0–4.7	7.7–10.1	83.9–86.9	93.3–95.2
12th	%	4.2	7.9	14.7	73.3	87.9
(n=1,787)	CI	3.4–5.2	6.7–9.2	13.1–16.4	71.2–75.3	86.3–89.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.7	1.8	6.7	89.8	96.5
(n=6,291)	CI	1.4–2.1	1.5–2.2	6.1–7.3	89.0–90.6	96.0–96.9
Male	%	1.8	1.8	7.3	89.0	96.4
(n=3,036)	CI	1.4–2.4	1.4–2.3	6.4–8.3	87.9–90.1	95.6–97.0
Female	%	1.2	1.7	5.9	91.2	97.1
(n=3,100)	CI	0.9–1.7	1.3–2.3	5.1–6.7	90.2–92.2	96.4–97.6
White	%	1.6	1.8	6.8	89.8	96.6
(n=5,268)	CI	1.2–1.9	1.5–2.2	6.2–7.6	88.9–90.6	96.1–97.1
Black	%	5.1	0.7	5.7	88.4	94.2
(n=137)	CI	2.4–10.4	0.1–5.1	2.9–11.2	81.8–92.8	88.7–97.1
Hispanic	%	1.8	0.9	4.6	92.7	97.3
(n=216)	CI	0.7–4.8	0.2–3.5	2.5–8.3	88.4–95.5	94.1–98.8
8th	%	1.0	0.8	4.3	93.8	98.2
(n=2,323)	CI	0.7–1.5	0.5–1.2	3.6–5.3	92.8–94.8	97.6–98.7
10th	%	1.3	1.7	5.6	91.4	97.0
(n=2,178)	CI	0.9–1.9	1.2–2.3	4.7–6.6	90.2–92.5	96.2–97.7
12th	%	2.8	3.1	10.4	83.7	94.1
(n=1,790)	CI	2.2–3.7	2.4–4.0	9.0–11.9	81.9–85.4	92.9–95.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.48. Parents Feel Using Electronic Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.5	7.9	12.2	75.3	87.5
(n=6,286)	CI	4.0–5.1	7.3–8.7	11.4–13.0	74.2–76.4	86.7–88.3
Male	%	4.8	8.8	12.5	74.0	86.5
(n=3,038)	CI	4.1–5.6	7.8–9.8	11.3–13.7	72.4–75.5	85.2–87.6
Female	%	3.8	7.2	11.7	77.3	89.0
(n=3,094)	CI	3.2–4.6	6.3–8.1	10.6–12.9	75.8–78.7	87.8–90.1
White	%	4.3	7.9	12.6	75.2	87.8
(n=5,265)	CI	3.7–4.9	7.2–8.7	11.7–13.5	74.1–76.4	86.9–88.7
Black	%	7.9	10.0	10.5	71.6	82.1
(n=136)	CI	4.4–13.8	5.9–16.6	6.2–17.0	63.3–78.7	74.6–87.8
Hispanic	%	4.6	3.7	7.4	84.3	91.7
(n=216)	CI	2.5–8.4	1.9–7.3	4.5–11.7	78.8–88.6	87.1–94.7
8th	%	3.0	4.6	11.2	81.2	92.4
(n=2,323)	CI	2.4–3.7	3.8–5.5	10.0–12.5	79.6–82.8	91.3–93.4
10th	%	3.8	7.0	12.2	77.1	89.2
(n=2,178)	CI	3.0–4.6	6.0–8.2	10.9–13.6	75.2–78.8	87.8–90.5
12th	%	7.0	12.6	13.3	67.1	80.4
(n=1,785)	CI	5.9–8.3	11.1–14.2	11.8–15.0	64.9–69.2	78.5–82.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.1	4.8	8.2	82.9	91.1
(n=6,284)	CI	3.6–4.6	4.3–5.4	7.5–8.9	82.0–83.9	90.4–91.8
Male	%	3.8	5.1	8.9	82.2	91.1
(n=3,031)	CI	3.2–4.6	4.3–5.9	8.0–10.0	80.7–83.5	90.0–92.1
Female	%	3.7	4.6	7.1	84.6	91.7
(n=3,098)	CI	3.1–4.5	3.9–5.4	6.3–8.1	83.2–85.8	90.7–92.6
White	%	3.5	4.9	8.0	83.6	91.6
(n=5,262)	CI	3.1–4.1	4.3–5.5	7.3–8.8	82.6–84.6	90.8–92.3
Black	%	8.2	4.4	13.7	73.6	87.3
(n=135)	CI	4.6–14.4	2.0–9.6	8.8–20.8	65.4–80.5	80.4–92.0
Hispanic	%	2.9	3.2	6.9	87.1	94.0
(n=218)	CI	1.3–6.2	1.5–6.5	4.2–11.2	81.8–91.0	89.9–96.5
8th	%	2.5	2.3	5.5	89.7	95.2
(n=2,321)	CI	1.9–3.2	1.8–3.0	4.7–6.5	88.4–90.9	94.3–96.0
10th	%	3.8	5.6	8.5	82.1	90.6
(n=2,178)	CI	3.1–4.7	4.7–6.6	7.4–9.8	80.4–83.6	89.3–91.7
12th	%	6.1	6.7	10.6	76.6	87.2
(n=1,785)	CI	5.1–7.3	5.6–8.0	9.3–12.1	74.6–78.5	85.6–88.7

Note: The confidence interval (CI) is 95 percent.

Table BH2.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.7	3.5	6.7	86.0	92.7
(n=6,277)	CI	3.3–4.3	3.1–4.0	6.1–7.3	85.1–86.9	92.0–93.3
Male	%	3.5	3.5	7.8	85.2	93.0
(n=3,027)	CI	2.9–4.2	2.9–4.2	6.9–8.9	83.9–86.4	92.1–93.9
Female	%	3.5	3.6	5.3	87.6	92.9
(n=3,096)	CI	2.9–4.2	3.0–4.3	4.6–6.2	86.4–88.7	92.0–93.8
White	%	3.3	3.5	6.6	86.6	93.2
(n=5,260)	CI	2.9–3.9	3.0–4.0	5.9–7.3	85.6–87.5	92.4–93.8
Black	%	7.5	5.9	9.1	77.5	86.6
(n=134)	CI	4.1–13.6	2.9–11.4	5.2–15.4	69.5–83.9	79.6–91.4
Hispanic	%	2.9	1.8	4.6	90.7	95.4
(n=216)	CI	1.3–6.3	0.7–4.7	2.5–8.5	86.0–94.0	91.5–97.5
8th	%	2.2	1.7	4.3	91.8	96.1
(n=2,317)	CI	1.7–2.9	1.2–2.3	3.5–5.2	90.6–92.9	95.2–96.8
10th	%	3.5	4.2	6.9	85.4	92.3
(n=2,177)	CI	2.8–4.3	3.4–5.1	5.9–8.1	83.9–86.8	91.1–93.4
12th	%	5.7	4.9	9.0	80.5	89.5
(n=1,783)	CI	4.7–6.8	4.0–6.0	7.8–10.4	78.6–82.2	88.0–90.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.51. Friends Feel Misusing Prescription Drugs is Wrong

How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.6	6.6	14.3	72.5	86.8
(n=6,285)	CI	6.0–7.3	6.0–7.2	13.4–15.2	71.4–73.6	85.9–87.6
Male	%	7.2	7.0	15.2	70.6	85.8
(n=3,033)	CI	6.3–8.2	6.1–8.0	14.0–16.5	68.9–72.2	84.5–87.0
Female	%	5.6	6.2	13.4	74.7	88.2
(n=3,097)	CI	4.8–6.5	5.4–7.2	12.3–14.7	73.2–76.2	87.0–89.3
White	%	6.2	6.5	14.2	73.2	87.3
(n=5,259)	CI	5.6–6.9	5.8–7.2	13.3–15.2	71.9–74.3	86.4–88.2
Black	%	12.2	6.8	13.3	67.7	81.1
(n=137)	CI	7.6–18.8	3.5–12.6	8.5–20.3	59.3–75.1	73.5–86.8
Hispanic	%	6.8	5.9	16.8	70.5	87.2
(n=218)	CI	4.1–11.1	3.5–10.0	12.3–22.3	64.0–76.2	82.1–91.1
8th	%	5.3	4.8	12.8	77.1	89.9
(n=2,321)	CI	4.5–6.3	4.0–5.7	11.5–14.2	75.4–78.8	88.6–91.1
10th	%	7.7	7.4	15.4	69.5	84.9
(n=2,176)	CI	6.7–8.9	6.4–8.6	13.9–17.0	67.5–71.4	83.3–86.3
12th	%	6.9	7.7	14.7	70.6	85.4
(n=1,788)	CI	5.8–8.2	6.6–9.0	13.2–16.5	68.5–72.7	83.7–86.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.52. Friends Feel Drinking Alcohol is Wrong

How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.4	14.9	17.2	53.6	70.8
(n=6,286)	CI	13.5–15.3	14.0–15.8	16.3–18.2	52.3–54.8	69.6–71.9
Male	%	15.9	14.5	17.6	52.0	69.6
(n=3,033)	CI	14.7–17.3	13.3–15.8	16.3–19.0	50.2–53.7	67.9–71.2
Female	%	12.5	15.2	16.8	55.4	72.2
(n=3,098)	CI	11.4–13.8	14.0–16.6	15.6–18.2	53.6–57.1	70.6–73.8
White	%	13.8	14.8	17.4	54.0	71.4
(n=5,263)	CI	12.9–14.8	13.8–15.7	16.4–18.4	52.7–55.4	70.2–72.6
Black	%	21.4	8.9	18.1	51.6	69.7
(n=137)	CI	15.2–29.2	5.1–15.1	12.5–25.6	43.1–60.0	61.4–77.0
Hispanic	%	15.3	16.5	17.5	50.7	68.2
(n=216)	CI	11.0–20.8	12.1–22.2	12.9–23.2	44.0–57.4	61.6–74.1
8th	%	7.6	11.1	16.4	64.9	81.3
(n=2,322)	CI	6.6–8.7	9.9–12.5	15.0–18.0	62.9–66.8	79.7–82.9
10th	%	16.8	16.6	18.1	48.5	66.6
(n=2,178)	CI	15.3–18.5	15.1–18.2	16.5–19.8	46.4–50.6	64.6–68.5
12th	%	19.1	17.1	17.1	46.8	63.9
(n=1,786)	CI	17.3–20.9	15.4–18.9	15.4–18.9	44.5–49.1	61.6–66.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.5	12.5	17.4	55.6	73.0
(n=6,279)	CI	13.6–15.4	11.7–13.4	16.5–18.3	54.4–56.8	71.8–74.1
Male	%	16.2	12.2	18.5	53.1	71.6
(n=3,028)	CI	14.9–17.6	11.1–13.5	17.2–19.9	51.3–54.9	69.9–73.2
Female	%	12.4	12.8	16.6	58.1	74.7
(n=3,096)	CI	11.3–13.6	11.7–14.1	15.3–18.0	56.4–59.9	73.2–76.2
White	%	14.2	12.7	17.5	55.7	73.1
(n=5,257)	CI	13.3–15.2	11.8–13.6	16.5–18.5	54.3–57.0	71.9–74.3
Black	%	20.8	6.5	13.1	59.6	72.7
(n=137)	CI	14.7–28.6	3.4–12.1	8.4–20.0	51.1–67.6	64.5–79.7
Hispanic	%	10.7	11.2	22.4	55.7	78.1
(n=215)	CI	7.2–15.6	7.6–16.3	17.3–28.5	48.9–62.3	72.0–83.2
8th	%	8.3	9.1	16.2	66.4	82.7
(n=2,317)	CI	7.2–9.5	8.0–10.3	14.8–17.8	64.5–68.3	81.1–84.1
10th	%	15.8	13.3	18.4	52.5	70.9
(n=2,177)	CI	14.3–17.4	11.9–14.8	16.8–20.1	50.4–54.6	69.0–72.8
12th	%	19.8	15.5	17.5	47.1	64.7
(n=1,785)	CI	18.0–21.7	13.9–17.3	15.8–19.3	44.8–49.5	62.4–66.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	24.4	14.4	14.3	46.9	61.2
(n=6,264)	CI	23.4–25.5	13.5–15.3	13.5–15.2	45.6–48.1	60.0–62.4
Male	%	26.1	13.3	15.3	45.4	60.7
(n=3,022)	CI	24.5–27.7	12.1–14.5	14.1–16.6	43.6–47.1	58.9–62.4
Female	%	22.5	15.6	13.6	48.3	61.9
(n=3,088)	CI	21.0–24.0	14.3–16.9	12.5–14.9	46.5–50.1	60.2–63.6
White	%	24.3	14.3	14.6	46.9	61.5
(n=5,247)	CI	23.1–25.4	13.3–15.2	13.7–15.6	45.5–48.2	60.2–62.8
Black	%	27.5	9.2	14.1	49.2	63.3
(n=135)	CI	20.5–35.7	5.3–15.6	9.1–21.2	40.7–57.7	54.7–71.1
Hispanic	%	19.8	14.1	14.1	52.0	66.1
(n=212)	CI	14.9–25.8	10.0–19.5	10.0–19.5	45.2–58.7	59.4–72.2
8th	%	15.2	12.6	14.8	57.3	72.2
(n=2,317)	CI	13.8–16.7	11.3–14.1	13.4–16.3	55.3–59.3	70.3–74.0
10th	%	26.8	15.3	14.4	43.5	57.9
(n=2,167)	CI	25.0–28.7	13.8–16.8	13.0–15.9	41.5–45.6	55.8–60.0
12th	%	31.9	15.4	13.6	39.1	52.7
(n=1,780)	CI	29.8–34.1	13.8–17.1	12.1–15.3	36.9–41.4	50.4–55.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	19.1	11.4	14.4	55.1	69.5
(n=6,282)	CI	18.2–20.1	10.6–12.2	13.5–15.3	53.8–56.3	68.3–70.6
Male	%	20.3	10.6	15.5	53.6	69.1
(n=3,028)	CI	18.9–21.8	9.6–11.8	14.3–16.9	51.8–55.4	67.4–70.7
Female	%	17.5	12.2	13.4	57.0	70.4
(n=3,099)	CI	16.1–18.8	11.1–13.4	12.2–14.6	55.2–58.7	68.7–72.0
White	%	18.2	11.2	14.7	56.0	70.6
(n=5,257)	CI	17.1–19.2	10.4–12.1	13.7–15.7	54.6–57.3	69.4–71.9
Black	%	31.9	10.2	11.7	46.2	57.9
(n=137)	CI	24.5–40.3	6.1–16.6	7.2–18.4	38.0–54.7	49.4–66.0
Hispanic	%	17.7	11.1	16.4	54.8	71.2
(n=217)	CI	13.1–23.5	7.5–16.1	12.0–21.9	48.0–61.3	64.7–76.9
8th	%	9.9	8.1	13.4	68.6	82.0
(n=2,321)	CI	8.8–11.2	7.0–9.2	12.1–14.8	66.7–70.5	80.4–83.5
10th	%	23.2	12.4	15.9	48.6	64.4
(n=2,173)	CI	21.5–25.0	11.1–13.8	14.4–17.5	46.5–50.7	62.4–66.4
12th	%	24.7	14.0	13.8	47.4	61.3
(n=1,788)	CI	22.8–26.8	12.5–15.7	12.3–15.5	45.1–49.8	59.0–63.5

Note: The confidence interval (CI) is 95 percent.

Table BH2.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	88.7	5.4	2.3	0.7	2.9	11.3
(n=6,270)	CI	87.9–89.5	4.8–6.0	2.0–2.7	0.5–0.9	2.5–3.3	10.5–12.1
Male	%	89.9	4.3	2.2	0.5	3.1	10.1
(n=3,032)	CI	88.7–90.9	3.6–5.1	1.7–2.8	0.3–0.9	2.5–3.8	9.1–11.3
Female	%	88.1	6.5	2.5	0.8	2.1	11.9
(n=3,084)	CI	86.9–89.2	5.7–7.4	2.0–3.1	0.5–1.2	1.6–2.7	10.8–13.1
White	%	89.1	5.4	2.4	0.7	2.5	10.9
(n=5,255)	CI	88.2–89.9	4.8–6.0	2.0–2.8	0.5–1.0	2.1–2.9	10.1–11.8
Black	%	88.6	6.2	2.3	0.7	2.2	11.4
(n=136)	CI	81.9–93.1	3.1–12.0	0.7–6.9	0.1–4.8	0.7–6.8	6.9–18.1
Hispanic	%	90.2	5.2	0.9	0.5	3.2	9.8
(n=212)	CI	85.4–93.6	2.9–9.2	0.2–3.6	0.1–3.7	1.5–6.6	6.4–14.6
8th	%	91.2	4.7	1.9	0.4	1.8	8.8
(n=2,319)	CI	89.9–92.3	3.9–5.6	1.5–2.6	0.2–0.8	1.3–2.4	7.7–10.1
10th	%	86.5	6.3	2.7	1.0	3.5	13.5
(n=2,171)	CI	85.0–87.8	5.4–7.4	2.1–3.5	0.7–1.5	2.8–4.4	12.2–15.0
12th	%	88.6	5.2	2.4	0.6	3.4	11.4
(n=1,780)	CI	87.0–90.0	4.2–6.3	1.7–3.2	0.3–1.0	2.6–4.3	10.0–13.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	55.6	13.7	10.1	4.7	15.9	44.4
(n=6,318)	CI	54.3–56.8	12.9–14.6	9.4–10.9	4.2–5.2	15.0–16.9	43.2–45.7
Male	%	58.6	12.4	8.7	4.0	16.4	41.4
(n=3,057)	CI	56.8–60.3	11.3–13.6	7.7–9.8	3.3–4.7	15.1–17.7	39.7–43.2
Female	%	52.7	15.2	11.6	5.4	15.1	47.3
(n=3,105)	CI	50.9–54.4	14.0–16.5	10.5–12.8	4.7–6.3	13.9–16.4	45.6–49.1
White	%	56.1	13.7	10.0	4.8	15.5	43.9
(n=5,291)	CI	54.7–57.4	12.8–14.6	9.2–10.9	4.2–5.4	14.5–16.5	42.6–45.3
Black	%	58.0	10.0	10.2	4.6	17.2	42.0
(n=139)	CI	49.5–66.0	6.0–16.3	6.1–16.6	2.1–9.9	11.7–24.5	34.0–50.5
Hispanic	%	55.2	16.8	7.9	2.8	17.3	44.8
(n=216)	CI	48.4–61.7	12.3–22.5	5.0–12.4	1.2–6.1	12.8–23.1	38.3–51.6
8th	%	71.2	12.2	6.8	2.7	7.1	28.8
(n=2,336)	CI	69.3–73.0	11.0–13.6	5.8–7.9	2.1–3.5	6.1–8.2	27.0–30.7
10th	%	48.3	15.5	12.4	5.4	18.4	51.7
(n=2,182)	CI	46.2–50.4	14.1–17.1	11.1–13.8	4.5–6.4	16.8–20.1	49.6–53.8
12th	%	46.5	13.4	11.4	6.0	22.8	53.5
(n=1,800)	CI	44.2–48.8	11.9–15.0	10.0–12.9	5.0–7.2	21.0–24.8	51.2–55.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	69.6	10.0	6.1	3.6	10.7	30.4
(n=6,298)	CI	68.5–70.8	9.3–10.8	5.5–6.7	3.1–4.1	10.0–11.5	29.2–31.5
Male	%	70.5	9.4	5.9	3.0	11.1	29.5
(n=3,045)	CI	68.8–72.1	8.4–10.5	5.1–6.8	2.5–3.7	10.0–12.3	27.9–31.2
Female	%	69.1	10.7	6.3	4.2	9.8	30.9
(n=3,098)	CI	67.4–70.7	9.6–11.8	5.4–7.2	3.5–5.0	8.8–10.9	29.3–32.6
White	%	69.7	10.1	6.1	3.6	10.5	30.3
(n=5,277)	CI	68.5–71.0	9.3–11.0	5.4–6.7	3.1–4.2	9.7–11.4	29.0–31.5
Black	%	71.8	7.5	5.9	3.8	11.0	28.2
(n=137)	CI	63.5–78.8	4.0–13.4	3.0–11.5	1.6–9.0	6.7–17.5	21.2–36.5
Hispanic	%	74.5	9.1	4.3	3.3	8.9	25.5
(n=214)	CI	68.2–80.0	5.8–13.8	2.2–8.1	1.6–6.8	5.7–13.5	20.0–31.8
8th	%	86.4	6.0	2.7	1.4	3.4	13.6
(n=2,331)	CI	84.9–87.7	5.1–7.1	2.2–3.5	1.0–2.0	2.8–4.3	12.3–15.1
10th	%	63.5	12.1	8.1	4.1	12.3	36.5
(n=2,171)	CI	61.4–65.5	10.8–13.5	7.0–9.3	3.3–5.0	11.0–13.8	34.5–38.6
12th	%	58.1	12.1	7.6	5.4	16.8	41.9
(n=1,796)	CI	55.8–60.4	10.6–13.6	6.4–8.9	4.5–6.6	15.2–18.6	39.6–44.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have smoked part, or all of a cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	70.6	11.0	6.1	3.3	9.0	29.4
(n=6,283)	CI	69.4–71.7	10.3–11.9	5.5–6.7	2.9–3.8	8.3–9.7	28.3–30.6
Male	%	71.9	10.8	5.1	2.7	9.5	28.1
(n=3,037)	CI	70.3–73.5	9.7–11.9	4.4–6.0	2.1–3.3	8.5–10.6	26.5–29.7
Female	%	69.6	11.5	7.1	3.9	8.0	30.4
(n=3,092)	CI	67.9–71.2	10.4–12.6	6.2–8.0	3.3–4.6	7.1–9.0	28.8–32.1
White	%	70.4	11.2	6.2	3.5	8.7	29.6
(n=5,264)	CI	69.2–71.7	10.4–12.1	5.6–6.9	3.0–4.0	7.9–9.5	28.3–30.8
Black	%	73.4	12.4	3.2	3.7	7.4	26.6
(n=137)	CI	65.3–80.3	7.8–19.1	1.2–8.2	1.5–8.6	4.0–13.3	19.7–34.7
Hispanic	%	78.2	8.0	3.3	1.0	9.6	21.8
(n=214)	CI	72.1–83.2	5.0–12.5	1.5–6.7	0.2–3.9	6.3–14.4	16.8–27.9
8th	%	79.8	9.0	4.3	2.1	4.8	20.2
(n=2,328)	CI	78.1–81.4	7.9–10.3	3.6–5.2	1.6–2.7	4.0–5.7	18.6–21.9
10th	%	67.4	12.7	6.7	3.3	9.9	32.6
(n=2,165)	CI	65.4–69.3	11.3–14.1	5.8–7.9	2.6–4.2	8.7–11.2	30.7–34.6
12th	%	63.9	11.5	7.4	4.6	12.7	36.1
(n=1,790)	CI	61.6–66.1	10.1–13.1	6.2–8.7	3.7–5.6	11.2–14.3	33.9–38.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used an e-cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	59.4	10.1	7.1	4.7	18.7	40.6
(n=6,271)	CI	58.2–60.6	9.3–10.8	6.5–7.8	4.2–5.3	17.8–19.7	39.4–41.8
Male	%	59.9	9.2	6.6	4.4	19.9	40.1
(n=3,032)	CI	58.1–61.6	8.2–10.3	5.8–7.6	3.7–5.2	18.5–21.3	38.4–41.9
Female	%	59.1	11.0	7.7	5.0	17.2	40.9
(n=3,083)	CI	57.3–60.8	9.9–12.1	6.8–8.7	4.3–5.8	15.9–18.6	39.2–42.7
White	%	59.0	10.5	7.3	4.7	18.5	41.0
(n=5,253)	CI	57.6–60.3	9.7–11.4	6.6–8.0	4.2–5.4	17.4–19.5	39.7–42.4
Black	%	68.2	4.4	6.1	2.3	19.0	31.8
(n=137)	CI	59.7–75.5	2.0–9.6	3.0–11.7	0.7–7.0	13.2–26.6	24.5–40.3
Hispanic	%	68.5	6.1	7.0	2.4	16.1	31.5
(n=211)	CI	61.8–74.4	3.5–10.2	4.2–11.4	1.0–5.7	11.7–21.7	25.6–38.2
8th	%	70.2	8.9	5.9	3.1	11.8	29.8
(n=2,320)	CI	68.3–72.1	7.8–10.1	5.1–7.0	2.4–3.8	10.6–13.2	27.9–31.7
10th	%	54.1	11.3	8.5	5.3	20.8	45.9
(n=2,166)	CI	52.0–56.2	10.0–12.7	7.4–9.7	4.4–6.3	19.2–22.6	43.8–48.0
12th	%	53.3	10.0	6.9	5.9	23.9	46.7
(n=1,785)	CI	51.0–55.6	8.7–11.5	5.8–8.2	4.9–7.1	22.0–26.0	44.4–49.0

Note: The confidence interval (CI) is 95 percent.

Table BH2.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	68.8	9.8	5.8	3.9	11.7	31.2
(n=6,303)	CI	67.7–70.0	9.1–10.6	5.2–6.4	3.4–4.4	10.9–12.5	30.0–32.3
Male	%	70.6	8.7	5.5	3.8	11.4	29.4
(n=3,045)	CI	68.9–72.2	7.8–9.8	4.7–6.4	3.2–4.6	10.3–12.6	27.8–31.1
Female	%	67.7	10.8	6.1	4.0	11.4	32.3
(n=3,102)	CI	66.0–69.3	9.7–11.9	5.3–7.0	3.4–4.8	10.3–12.6	30.7–34.0
White	%	69.6	10.0	5.7	3.9	10.7	30.4
(n=5,280)	CI	68.3–70.8	9.3–10.9	5.1–6.4	3.5–4.5	9.9–11.6	29.2–31.7
Black	%	58.0	6.6	9.8	5.8	19.8	42.0
(n=137)	CI	49.4–66.1	3.4–12.3	5.7–16.3	2.9–11.2	13.9–27.5	33.9–50.6
Hispanic	%	73.6	6.4	4.2	3.8	12.0	26.4
(n=216)	CI	67.3–79.1	3.8–10.5	2.2–7.9	1.9–7.5	8.3–17.2	20.9–32.7
8th	%	82.2	7.3	3.3	2.0	5.1	17.8
(n=2,334)	CI	80.6–83.7	6.3–8.5	2.6–4.1	1.5–2.7	4.3–6.1	16.3–19.4
10th	%	62.5	11.4	6.9	5.3	13.9	37.5
(n=2,179)	CI	60.5–64.5	10.2–12.8	5.9–8.1	4.4–6.3	12.5–15.4	35.5–39.5
12th	%	61.0	10.7	7.2	4.5	16.6	39.0
(n=1,790)	CI	58.8–63.3	9.4–12.2	6.1–8.5	3.6–5.5	14.9–18.3	36.7–41.2

Note: The confidence interval (CI) is 95 percent.

Table BH2.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	9.3	5.2	12.0	73.5	85.5
(n=6,214)	CI	8.6–10.1	4.6–5.7	11.2–12.8	72.4–74.6	84.6–86.4
Male	%	11.0	4.8	12.3	71.9	84.2
(n=3,005)	CI	9.9–12.2	4.1–5.6	11.1–13.5	70.3–73.5	82.9–85.5
Female	%	7.0	5.5	11.8	75.7	87.5
(n=3,055)	CI	6.2–8.0	4.7–6.3	10.7–13.0	74.1–77.2	86.3–88.6
White	%	8.7	5.2	12.0	74.0	86.0
(n=5,205)	CI	8.0–9.5	4.7–5.9	11.2–13.0	72.8–75.1	85.0–86.9
Black	%	16.7	5.1	5.8	72.4	78.2
(n=137)	CI	11.3–24.0	2.4–10.3	2.9–11.2	64.2–79.3	70.4–84.4
Hispanic	%	12.9	6.9	10.5	69.6	80.2
(n=210)	CI	8.9–18.3	4.2–11.3	7.1–15.4	63.0–75.5	74.1–85.1
8th	%	8.6	6.0	11.8	73.6	85.4
(n=2,301)	CI	7.5–9.8	5.1–7.1	10.6–13.2	71.7–75.4	83.9–86.8
10th	%	9.5	5.3	12.7	72.5	85.2
(n=2,150)	CI	8.3–10.8	4.5–6.4	11.3–14.2	70.5–74.3	83.6–86.6
12th	%	10.0	4.0	11.4	74.6	86.0
(n=1,763)	CI	8.6–11.4	3.2–5.0	10.0–12.9	72.6–76.6	84.3–87.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.63. Perceived Risk of Regularly Drinking Alcohol

		How much do you think people risk harming themselves physically and in other ways when they <u>take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	12.6	15.0	23.0	49.4	72.4
(n=6,256)	CI	11.8–13.5	14.1–15.9	22.0–24.1	48.1–50.6	71.3–73.5
Male	%	14.8	15.9	23.3	46.0	69.3
(n=3,021)	CI	13.5–16.1	14.6–17.3	21.8–24.8	44.3–47.8	67.7–71.0
Female	%	10.0	13.9	23.0	53.1	76.1
(n=3,081)	CI	9.0–11.1	12.7–15.2	21.5–24.5	51.3–54.9	74.5–77.6
White	%	12.1	15.1	23.5	49.4	72.9
(n=5,242)	CI	11.2–13.0	14.1–16.0	22.3–24.6	48.0–50.8	71.6–74.1
Black	%	17.5	17.5	16.0	49.0	65.0
(n=137)	CI	11.9–24.9	11.9–24.9	10.7–23.2	40.6–57.5	56.5–72.6
Hispanic	%	15.5	11.3	18.7	54.5	73.2
(n=212)	CI	11.1–21.1	7.7–16.4	14.0–24.6	47.7–61.1	66.7–78.8
8th	%	11.6	13.8	20.4	54.2	74.6
(n=2,314)	CI	10.4–13.0	12.5–15.3	18.8–22.1	52.1–56.2	72.7–76.3
10th	%	13.0	15.6	24.2	47.2	71.4
(n=2,167)	CI	11.7–14.5	14.2–17.2	22.4–26.0	45.1–49.3	69.4–73.2
12th	%	13.3	15.5	24.7	46.5	71.2
(n=1,775)	CI	11.8–14.9	13.9–17.3	22.7–26.7	44.2–48.9	69.1–73.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.64. Perceived Risk of Binge Drinking

		How much do you think people risk harming themselves physically and in other ways when they <u>take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	11.0	10.1	21.3	57.5	78.8
(n=6,253)	CI	10.3–11.8	9.4–10.9	20.3–22.4	56.3–58.7	77.8–79.8
Male	%	12.9	10.9	21.7	54.5	76.2
(n=3,026)	CI	11.7–14.1	9.8–12.1	20.3–23.3	52.7–56.3	74.7–77.7
Female	%	8.6	9.5	20.9	61.0	81.8
(n=3,073)	CI	7.7–9.7	8.5–10.6	19.4–22.3	59.2–62.7	80.4–83.2
White	%	10.4	10.1	21.4	58.1	79.5
(n=5,240)	CI	9.6–11.2	9.3–11.0	20.3–22.5	56.7–59.4	78.4–80.6
Black	%	18.4	11.7	18.9	51.0	69.9
(n=136)	CI	12.7–25.9	7.3–18.4	13.1–26.5	42.5–59.4	61.5–77.1
Hispanic	%	13.6	10.4	17.6	58.3	76.0
(n=213)	CI	9.5–19.0	6.9–15.4	13.1–23.4	51.5–64.8	69.6–81.3
8th	%	10.6	8.8	18.2	62.3	80.6
(n=2,313)	CI	9.4–12.0	7.7–10.0	16.7–19.9	60.3–64.3	78.9–82.1
10th	%	11.1	10.5	22.3	56.0	78.4
(n=2,161)	CI	9.9–12.5	9.3–11.9	20.6–24.1	53.9–58.1	76.6–80.0
12th	%	11.3	11.2	23.6	53.9	77.5
(n=1,779)	CI	9.9–12.9	9.8–12.8	21.6–25.6	51.6–56.2	75.5–79.3

Note: The confidence interval (CI) is 95 percent.

Table BH2.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.3	5.9	16.0	67.8	83.8
(n=6,266)	CI	9.6–11.1	5.4–6.6	15.1–16.9	66.6–68.9	82.8–84.7
Male	%	12.0	5.9	17.1	65.0	82.1
(n=3,026)	CI	10.9–13.2	5.2–6.9	15.8–18.5	63.3–66.7	80.6–83.4
Female	%	8.0	6.0	14.9	71.2	86.0
(n=3,086)	CI	7.1–9.0	5.2–6.9	13.7–16.2	69.5–72.7	84.8–87.2
White	%	9.7	5.9	15.8	68.6	84.4
(n=5,250)	CI	8.9–10.5	5.3–6.6	14.8–16.8	67.3–69.8	83.4–85.3
Black	%	19.2	6.6	13.1	61.1	74.2
(n=136)	CI	13.4–26.8	3.4–12.3	8.3–19.9	52.5–69.0	66.0–80.9
Hispanic	%	14.4	5.5	15.1	65.0	80.1
(n=213)	CI	10.2–19.9	3.1–9.5	10.9–20.5	58.3–71.2	74.1–85.0
8th	%	9.8	6.1	14.9	69.2	84.1
(n=2,318)	CI	8.6–11.1	5.2–7.2	13.5–16.4	67.3–71.0	82.5–85.5
10th	%	10.0	6.0	16.6	67.3	83.9
(n=2,166)	CI	8.8–11.4	5.1–7.1	15.1–18.2	65.3–69.3	82.3–85.4
12th	%	11.1	5.7	16.4	66.8	83.2
(n=1,782)	CI	9.8–12.7	4.7–6.8	14.8–18.2	64.6–68.9	81.4–84.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	13.6	19.1	25.5	41.7	67.3
(n=6,251)	CI	12.8–14.5	18.1–20.1	24.5–26.6	40.5–43.0	66.1–68.4
Male	%	16.2	21.3	23.6	38.9	62.5
(n=3,022)	CI	14.9–17.5	19.9–22.8	22.1–25.2	37.2–40.7	60.8–64.2
Female	%	10.6	16.8	27.7	44.9	72.6
(n=3,076)	CI	9.6–11.8	15.5–18.1	26.2–29.3	43.1–46.6	71.0–74.1
White	%	12.7	19.2	26.3	41.8	68.1
(n=5,241)	CI	11.9–13.7	18.1–20.3	25.1–27.5	40.5–43.1	66.8–69.3
Black	%	26.6	15.7	21.9	35.8	57.7
(n=137)	CI	19.8–34.8	10.5–22.8	15.7–29.7	28.1–44.3	49.1–65.8
Hispanic	%	17.9	15.5	19.5	47.1	66.6
(n=212)	CI	13.2–23.8	11.2–21.1	14.7–25.4	40.4–53.9	59.9–72.7
8th	%	12.8	17.3	22.3	47.5	69.8
(n=2,313)	CI	11.5–14.3	15.8–18.9	20.7–24.1	45.5–49.5	67.9–71.7
10th	%	13.5	20.1	26.1	40.3	66.4
(n=2,158)	CI	12.1–15.0	18.5–21.8	24.3–28.0	38.2–42.4	64.4–68.4
12th	%	14.7	19.9	28.4	37.1	65.4
(n=1,780)	CI	13.1–16.4	18.1–21.8	26.3–30.5	34.8–39.3	63.2–67.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	28.5	20.0	15.1	36.4	51.5
(n=6,209)	CI	27.4–29.6	19.1–21.1	14.2–16.0	35.2–37.6	50.2–52.7
Male	%	31.3	19.5	14.4	34.9	49.3
(n=2,999)	CI	29.6–33.0	18.1–20.9	13.2–15.7	33.2–36.6	47.5–51.1
Female	%	25.0	20.8	15.9	38.4	54.2
(n=3,057)	CI	23.5–26.6	19.4–22.2	14.6–17.2	36.7–40.1	52.5–56.0
White	%	27.6	20.0	15.2	37.2	52.4
(n=5,207)	CI	26.4–28.8	19.0–21.1	14.2–16.2	35.9–38.5	51.0–53.7
Black	%	41.4	22.2	11.5	24.9	36.4
(n=136)	CI	33.3–50.0	15.9–30.1	7.1–18.0	18.3–33.0	28.7–44.9
Hispanic	%	25.6	17.1	15.8	41.5	57.3
(n=207)	CI	20.0–32.1	12.5–22.9	11.4–21.5	34.9–48.4	50.4–64.0
8th	%	19.9	17.5	15.6	47.1	62.7
(n=2,285)	CI	18.3–21.5	16.0–19.1	14.2–17.2	45.0–49.1	60.7–64.6
10th	%	31.0	20.0	15.7	33.3	49.0
(n=2,152)	CI	29.1–33.0	18.4–21.8	14.2–17.3	31.3–35.3	46.8–51.1
12th	%	35.0	22.9	13.8	28.3	42.1
(n=1,772)	CI	32.8–37.3	21.0–24.9	12.3–15.5	26.3–30.5	39.8–44.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	20.8	15.5	18.9	44.8	63.7
(n=6,227)	CI	19.8–21.9	14.6–16.4	17.9–19.9	43.6–46.0	62.5–64.9
Male	%	23.5	16.0	18.4	42.1	60.6
(n=3,005)	CI	22.0–25.0	14.7–17.3	17.1–19.9	40.4–43.9	58.8–62.3
Female	%	17.7	14.9	19.5	48.0	67.4
(n=3,070)	CI	16.3–19.1	13.7–16.2	18.1–20.9	46.2–49.7	65.7–69.1
White	%	19.6	15.5	19.2	45.7	64.9
(n=5,277)	CI	18.5–20.7	14.5–16.5	18.2–20.3	44.4–47.1	63.6–66.2
Black	%	39.9	12.2	13.3	34.6	47.9
(n=134)	CI	31.8–48.5	7.6–19.2	8.5–20.2	27.0–43.2	39.4–56.5
Hispanic	%	21.1	12.3	20.1	46.5	66.6
(n=210)	CI	16.0–27.2	8.5–17.6	15.1–26.1	39.8–53.4	59.8–72.7
8th	%	15.2	11.1	17.4	56.2	73.7
(n=2,305)	CI	13.8–16.7	9.9–12.5	15.9–19.0	54.2–58.3	71.8–75.4
10th	%	22.8	16.4	19.4	41.3	60.7
(n=2,153)	CI	21.1–24.7	14.9–18.1	17.8–21.1	39.3–43.4	58.6–62.8
12th	%	24.8	19.2	20.0	36.1	56.1
(n=1,769)	CI	22.8–26.8	17.4–21.1	18.2–21.9	33.9–38.4	53.7–58.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.7	3.7	10.8	82.7	93.5
(n=6,261)	CI	2.4–3.2	3.3–4.2	10.1–11.6	81.8–83.6	92.9–94.1
Male	%	3.1	3.8	10.5	82.6	93.1
(n=3,024)	CI	2.5–3.8	3.2–4.6	9.4–11.6	81.2–83.9	92.1–93.9
Female	%	1.9	3.5	11.1	83.5	94.6
(n=3,084)	CI	1.4–2.4	2.9–4.2	10.1–12.3	82.1–84.8	93.7–95.4
White	%	2.4	3.7	10.9	83.0	93.9
(n=5,248)	CI	2.0–2.8	3.2–4.3	10.1–11.8	82.0–84.0	93.3–94.6
Black	%	7.4	6.0	9.6	77.1	86.6
(n=136)	CI	4.0–13.3	3.0–11.7	5.6–15.9	69.1–83.5	79.6–91.5
Hispanic	%	3.8	4.2	11.1	80.9	92.0
(n=213)	CI	1.9–7.5	2.2–7.9	7.5–16.1	75.0–85.6	87.4–95.0
8th	%	2.0	2.9	9.0	86.0	95.1
(n=2,313)	CI	1.5–2.6	2.3–3.7	7.9–10.3	84.6–87.4	94.1–95.9
10th	%	2.8	4.5	11.6	81.2	92.7
(n=2,169)	CI	2.2–3.5	3.7–5.5	10.3–13.0	79.4–82.7	91.5–93.7
12th	%	3.5	3.8	11.9	80.8	92.7
(n=1,779)	CI	2.8–4.5	3.0–4.7	10.5–13.5	78.9–82.6	91.4–93.8

Note: The confidence interval (CI) is 95 percent.

Table BH2.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.4	9.9	18.3	66.4	84.7
(n=6,280)	CI	4.9–6.0	9.2–10.7	17.4–19.3	65.2–67.5	83.7–85.5
Male	%	5.9	9.9	17.7	66.4	84.2
(n=3,035)	CI	5.1–6.8	8.9–11.0	16.4–19.1	64.7–68.1	82.8–85.4
Female	%	4.4	9.9	19.2	66.5	85.7
(n=3,091)	CI	3.7–5.2	8.9–11.0	17.8–20.6	64.8–68.1	84.4–86.9
White	%	4.9	10.1	18.0	67.0	85.0
(n=5,262)	CI	4.4–5.6	9.3–10.9	16.9–19.0	65.7–68.3	84.0–85.9
Black	%	10.5	8.8	17.0	63.7	80.7
(n=137)	CI	6.3–17.1	5.0–14.9	11.5–24.5	55.1–71.4	73.1–86.6
Hispanic	%	6.2	9.7	20.0	64.1	84.1
(n=216)	CI	3.6–10.4	6.4–14.5	15.2–26.0	57.4–70.3	78.5–88.5
8th	%	3.0	5.8	12.9	78.3	91.2
(n=2,321)	CI	2.4–3.8	4.9–6.8	11.6–14.3	76.6–79.9	90.0–92.3
10th	%	5.6	11.4	19.8	63.3	83.1
(n=2,173)	CI	4.7–6.6	10.1–12.8	18.2–21.5	61.2–65.3	81.4–84.6
12th	%	7.8	12.9	22.6	56.7	79.3
(n=1,786)	CI	6.7–9.2	11.4–14.5	20.7–24.6	54.4–59.0	77.3–81.1

Note: The confidence interval (CI) is 95 percent.

Table BH2.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.4	10.3	17.8	64.5	82.3
(n=6,278)	CI	6.7–8.1	9.6–11.1	16.9–18.8	63.3–65.7	81.3–83.2
Male	%	8.4	9.7	17.6	64.3	81.9
(n=3,031)	CI	7.5–9.5	8.7–10.8	16.2–19.0	62.6–66.0	80.5–83.2
Female	%	5.9	10.8	18.2	65.1	83.3
(n=3,095)	CI	5.1–6.8	9.8–12.0	16.8–19.6	63.4–66.8	81.9–84.6
White	%	7.1	10.8	17.8	64.3	82.1
(n=5,262)	CI	6.4–7.8	10.0–11.7	16.8–18.9	63.0–65.6	81.0–83.2
Black	%	10.1	5.8	17.2	66.9	84.0
(n=136)	CI	5.9–16.8	2.9–11.3	11.6–24.6	58.3–74.4	76.6–89.4
Hispanic	%	5.8	5.9	15.0	73.3	88.4
(n=215)	CI	3.3–9.9	3.4–9.9	10.8–20.5	66.9–78.9	83.3–92.0
8th	%	3.0	5.7	13.1	78.2	91.4
(n=2,323)	CI	2.4–3.7	4.8–6.7	11.8–14.6	76.5–79.9	90.1–92.4
10th	%	6.8	10.3	19.6	63.3	82.9
(n=2,171)	CI	5.8–8.0	9.1–11.6	18.0–21.3	61.3–65.3	81.3–84.4
12th	%	12.8	15.4	21.0	50.8	71.8
(n=1,784)	CI	11.3–14.4	13.8–17.2	19.1–22.9	48.5–53.2	69.7–73.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	12.7	15.7	17.0	54.6	71.6
(n=6,265)	CI	11.8–13.5	14.9–16.7	16.1–18.0	53.4–55.8	70.5–72.7
Male	%	13.7	14.8	16.5	55.0	71.5
(n=3,029)	CI	12.5–15.0	13.6–16.2	15.2–17.8	53.2–56.8	69.8–73.1
Female	%	11.1	16.7	17.9	54.3	72.2
(n=3,085)	CI	10.0–12.3	15.4–18.1	16.5–19.3	52.6–56.1	70.6–73.8
White	%	12.2	16.1	17.3	54.5	71.7
(n=5,253)	CI	11.3–13.1	15.1–17.1	16.3–18.3	53.1–55.8	70.5–72.9
Black	%	18.9	13.0	11.9	56.2	68.1
(n=135)	CI	13.1–26.6	8.2–20.0	7.3–18.6	47.6–64.5	59.6–75.6
Hispanic	%	11.0	12.2	13.5	63.4	76.9
(n=214)	CI	7.4–16.0	8.4–17.3	9.5–18.8	56.7–69.7	70.7–82.1
8th	%	7.0	10.4	14.6	68.0	82.5
(n=2,318)	CI	6.1–8.1	9.3–11.7	13.2–16.1	66.0–69.8	80.9–84.0
10th	%	12.2	16.5	18.2	53.1	71.2
(n=2,164)	CI	10.9–13.7	15.0–18.2	16.6–19.8	51.0–55.2	69.3–73.1
12th	%	19.2	20.6	18.4	41.7	60.2
(n=1,783)	CI	17.4–21.1	18.8–22.6	16.7–20.3	39.5–44.1	57.9–62.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	11.2	11.1	14.2	63.6	77.7
(n=6,271)	CI	10.5–12.0	10.3–11.9	13.3–15.0	62.3–64.7	76.7–78.7
Male	%	11.8	10.0	14.3	63.9	78.2
(n=3,029)	CI	10.7–13.0	9.0–11.1	13.1–15.6	62.1–65.6	76.7–79.7
Female	%	10.2	11.9	14.1	63.8	77.9
(n=3,089)	CI	9.2–11.3	10.8–13.1	12.9–15.3	62.1–65.5	76.4–79.3
White	%	10.3	11.0	14.3	64.3	78.6
(n=5,257)	CI	9.5–11.2	10.2–11.9	13.4–15.3	63.0–65.6	77.5–79.7
Black	%	25.0	9.6	12.5	52.8	65.3
(n=136)	CI	18.3–33.2	5.6–16.0	7.9–19.3	44.3–61.2	56.8–73.0
Hispanic	%	10.1	8.4	13.2	68.2	81.4
(n=214)	CI	6.7–15.1	5.3–13.0	9.2–18.5	61.6–74.2	75.5–86.2
8th	%	5.0	5.5	10.3	79.2	89.4
(n=2,317)	CI	4.2–6.0	4.7–6.5	9.1–11.6	77.5–80.8	88.1–90.6
10th	%	12.8	13.4	14.6	59.1	73.8
(n=2,172)	CI	11.5–14.3	12.1–14.9	13.2–16.2	57.1–61.2	71.9–75.6
12th	%	16.2	14.5	17.9	51.4	69.3
(n=1,782)	CI	14.6–18.0	12.9–16.2	16.2–19.7	49.1–53.7	67.1–71.4

Note: The confidence interval (CI) is 95 percent.

Table BH2.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	61.9	27.3	10.8	38.1
(n=6,255)	CI	60.7–63.1	26.3–28.5	10.0–11.6	36.9–39.3
Male	%	63.8	25.6	10.6	36.2
(n=3,022)	CI	62.0–65.5	24.1–27.2	9.6–11.8	34.5–38.0
Female	%	60.2	29.5	10.3	39.8
(n=3,081)	CI	58.5–61.9	27.9–31.1	9.3–11.5	38.1–41.5
White	%	62.0	27.7	10.3	38.0
(n=5,242)	CI	60.7–63.3	26.5–28.9	9.5–11.2	36.7–39.3
Black	%	57.0	29.4	13.6	43.0
(n=137)	CI	48.5–65.1	22.3–37.7	8.7–20.6	34.9–51.5
Hispanic	%	64.3	22.9	12.8	35.7
(n=215)	CI	57.6–70.5	17.7–29.0	9.0–18.0	29.5–42.4
8th	%	58.2	30.4	11.5	41.8
(n=2,310)	CI	56.1–60.2	28.5–32.3	10.2–12.8	39.8–43.9
10th	%	61.3	27.5	11.3	38.7
(n=2,165)	CI	59.2–63.3	25.6–29.4	10.0–12.7	36.7–40.8
12th	%	66.6	23.9	9.5	33.4
(n=1,780)	CI	64.4–68.7	22.0–26.0	8.2–10.9	31.3–35.6

Note: The confidence interval (CI) is 95 percent.

Table BH2.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	47.7	33.6	18.7	52.3
(n=6,259)	CI	46.5–49.0	32.4–34.8	17.8–19.7	51.0–53.5
Male	%	50.9	32.1	17.1	49.1
(n=3,030)	CI	49.1–52.6	30.4–33.8	15.8–18.5	47.4–50.9
Female	%	44.4	35.4	20.2	55.6
(n=3,077)	CI	42.7–46.2	33.7–37.1	18.8–21.6	53.8–57.3
White	%	47.9	34.1	18.0	52.1
(n=5,246)	CI	46.6–49.3	32.8–35.4	17.0–19.0	50.7–53.4
Black	%	43.8	31.4	24.8	56.2
(n=136)	CI	35.6–52.4	24.1–39.8	18.1–32.9	47.6–64.4
Hispanic	%	49.1	29.7	21.3	50.9
(n=215)	CI	42.4–55.8	23.9–36.2	16.3–27.3	44.2–57.6
8th	%	45.1	34.9	20.0	54.9
(n=2,311)	CI	43.0–47.1	33.0–36.9	18.4–21.7	52.9–57.0
10th	%	46.0	34.1	19.9	54.0
(n=2,170)	CI	43.9–48.1	32.1–36.1	18.3–21.6	51.9–56.1
12th	%	52.5	31.5	16.0	47.5
(n=1,778)	CI	50.1–54.8	29.4–33.7	14.4–17.8	45.2–49.9

Note: The confidence interval (CI) is 95 percent.

Table BH2.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	36.8	34.1	29.2	63.2
(n=6,257)	CI	35.6–38.0	32.9–35.2	28.0–30.3	62.0–64.4
Male	%	40.2	31.5	28.2	59.8
(n=3,026)	CI	38.5–42.0	29.9–33.2	26.7–29.9	58.0–61.5
Female	%	32.9	37.2	29.9	67.1
(n=3,080)	CI	31.3–34.6	35.5–39.0	28.3–31.5	65.4–68.7
White	%	35.8	34.9	29.3	64.2
(n=5,248)	CI	34.5–37.1	33.6–36.2	28.1–30.5	62.9–65.5
Black	%	41.1	29.8	29.0	58.9
(n=137)	CI	33.1–49.7	22.7–38.1	21.9–37.4	50.3–66.9
Hispanic	%	46.4	27.6	26.0	53.6
(n=213)	CI	39.8–53.2	21.9–34.0	20.5–32.4	46.8–60.2
8th	%	33.8	33.2	33.0	66.2
(n=2,312)	CI	31.9–35.8	31.3–35.2	31.1–34.9	64.2–68.1
10th	%	36.3	34.7	28.9	63.7
(n=2,166)	CI	34.3–38.4	32.8–36.8	27.0–30.9	61.6–65.7
12th	%	40.5	34.3	25.2	59.5
(n=1,779)	CI	38.2–42.8	32.1–36.5	23.3–27.3	57.2–61.8

Note: The confidence interval (CI) is 95 percent.

Behavioral Health Planning Region 3 Results Summary

Table BH3.1. Lifetime Prescription Drug Misuse

		On how many occasions (if any) have you ever <u>used prescription drugs not prescribed to you?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	91.7	4.2	1.4	0.8	0.6	0.1	1.0	8.3
(n=5,150)	CI	90.9-92.5	3.7-4.9	1.1-1.8	0.6-1.1	0.4-0.9	0.1-0.3	0.7-1.4	7.5-9.1
Male	%	93.1	2.8	1.2	0.7	0.7	0.1	1.4	6.9
(n=2,572)	CI	91.9-94.1	2.2-3.6	0.8-1.8	0.4-1.1	0.4-1.2	0.0-0.4	1.0-2.0	5.9-8.1
Female	%	90.9	5.5	1.5	0.9	0.5	0.2	0.4	9.1
(n=2,456)	CI	89.6-92.0	4.6-6.6	1.1-2.1	0.6-1.4	0.3-1.0	0.1-0.5	0.2-0.8	8.0-10.4
White	%	92.4	4.0	1.3	0.8	0.6	0.1	0.9	7.6
(n=3,577)	CI	91.4-93.2	3.3-4.7	0.9-1.7	0.6-1.2	0.3-0.9	0.0-0.4	0.6-1.3	6.8-8.6
Black	%	90.9	4.8	1.3	0.4	0.5	0.1	1.8	9.1
(n=491)	CI	87.7-93.4	3.1-7.5	0.6-3.0	0.1-1.9	0.1-2.1	0.0-1.0	0.9-3.7	6.6-12.3
Hispanic	%	92.3	4.2	2.0	0.7	0.7	0.0	0.2	7.7
(n=466)	CI	89.1-94.5	2.6-6.7	1.0-4.1	0.3-1.9	0.2-2.6	-	0.0-1.1	5.5-10.9
8th	%	94.0	3.1	1.3	0.9	0.4	0.1	0.4	6.0
(n=2,545)	CI	93.0-94.9	2.5-3.8	0.9-1.8	0.6-1.3	0.2-0.7	0.0-0.3	0.2-0.7	5.1-7.0
10th	%	90.9	4.8	1.7	0.7	0.5	0.1	1.3	9.1
(n=1,552)	CI	89.3-92.2	3.9-6.0	1.2-2.5	0.4-1.3	0.2-0.9	0.0-0.5	0.8-2.0	7.8-10.7
12th	%	90.2	4.9	1.2	0.9	1.1	0.2	1.4	9.8
(n=1,053)	CI	88.2-91.8	3.8-6.4	0.7-2.1	0.5-1.8	0.6-2.0	0.0-0.8	0.9-2.3	8.2-11.8

Note: The confidence interval (CI) is 95 percent.

Table BH3.2. Lifetime Alcohol Use

		On how many occasions (if any) have you ever <u>had one or more drinks of an alcoholic beverage?</u>							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	59.5	18.4	8.6	4.2	3.3	2.2	3.9	40.5
(n=5,162)	CI	58.0-60.9	17.3-19.5	7.8-9.4	3.6-4.8	2.8-3.9	1.8-2.7	3.4-4.6	39.1-42.0
Male	%	61.6	15.3	7.9	4.7	3.8	2.0	4.5	38.4
(n=2,580)	CI	59.6-63.6	13.9-16.8	6.9-9.2	3.9-5.7	3.0-4.7	1.5-2.8	3.7-5.5	36.4-40.4
Female	%	58.0	21.1	9.1	3.7	2.9	2.3	2.9	42.0
(n=2,456)	CI	55.9-60.0	19.4-22.9	7.9-10.4	2.9-4.6	2.3-3.8	1.7-3.1	2.3-3.8	40.0-44.1
White	%	58.0	18.3	9.0	4.5	3.4	2.4	4.5	42.0
(n=3,584)	CI	56.3-59.7	17.0-19.7	8.0-10.1	3.8-5.3	2.8-4.1	1.9-3.0	3.7-5.3	40.3-43.7
Black	%	67.3	17.6	6.6	3.7	2.7	0.8	1.3	32.7
(n=495)	CI	62.8-71.6	14.3-21.4	4.6-9.4	2.2-6.0	1.5-4.9	0.3-2.5	0.5-3.1	28.4-37.2
Hispanic	%	67.9	19.6	4.6	1.4	1.9	2.0	2.6	32.1
(n=468)	CI	63.1-72.3	16.0-23.7	2.9-7.2	0.6-3.1	0.9-4.0	0.9-4.4	1.4-5.0	27.7-36.9
8th	%	72.7	16.7	5.6	2.2	1.3	0.7	0.9	27.3
(n=2,537)	CI	70.9-74.4	15.3-18.2	4.7-6.5	1.7-2.9	0.9-1.8	0.5-1.2	0.6-1.3	25.6-29.1
10th	%	55.9	20.0	10.1	4.9	3.2	2.0	4.0	44.1
(n=1,562)	CI	53.4-58.3	18.1-22.1	8.7-11.7	3.9-6.0	2.4-4.2	1.4-2.8	3.1-5.1	41.7-46.6
12th	%	49.1	18.5	10.1	5.5	5.6	3.9	7.2	50.9
(n=1,063)	CI	46.1-52.1	16.2-20.9	8.4-12.1	4.3-7.1	4.4-7.2	2.9-5.3	5.8-8.9	47.9-53.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.3. Lifetime Binge Drinking

		On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	82.2	6.6	3.9	2.4	1.6	1.3	2.0	17.8
(n=5,166)	CI	81.0–83.3	5.9–7.4	3.4–4.6	1.9–2.9	1.3–2.1	1.0–1.7	1.6–2.5	16.7–19.0
Male	%	81.5	6.3	4.0	2.3	1.9	1.4	2.4	18.5
(n=2,582)	CI	79.8–83.1	5.4–7.4	3.3–5.0	1.7–3.1	1.4–2.7	1.0–2.1	1.8–3.2	16.9–20.2
Female	%	83.1	6.9	3.8	2.3	1.3	1.2	1.3	16.9
(n=2,460)	CI	81.4–84.7	5.9–8.1	3.1–4.8	1.7–3.1	0.9–2.0	0.8–1.8	0.9–1.9	15.3–18.6
White	%	80.6	7.1	4.3	2.5	1.7	1.5	2.3	19.4
(n=3,593)	CI	79.2–82.0	6.2–8.0	3.6–5.2	2.0–3.1	1.3–2.3	1.1–2.0	1.8–3.0	18.0–20.8
Black	%	89.2	4.9	2.3	1.6	0.7	0.8	0.4	10.8
(n=493)	CI	85.7–91.8	3.2–7.5	1.2–4.4	0.7–3.5	0.2–2.3	0.3–2.5	0.1–1.9	8.2–14.3
Hispanic	%	88.5	4.3	2.0	2.0	1.0	0.7	1.6	11.5
(n=466)	CI	84.7–91.5	2.6–6.8	0.9–4.1	0.9–4.4	0.3–3.1	0.2–2.6	0.7–3.8	8.5–15.3
8th	%	93.4	3.6	1.3	0.7	0.3	0.2	0.5	6.6
(n=2,539)	CI	92.4–94.3	3.0–4.4	0.9–1.8	0.5–1.2	0.1–0.6	0.1–0.4	0.3–0.8	5.7–7.6
10th	%	80.9	8.5	3.7	2.3	1.1	1.5	2.0	19.1
(n=1,564)	CI	78.9–82.8	7.2–10.0	2.9–4.8	1.7–3.2	0.7–1.7	1.0–2.3	1.4–2.8	17.2–21.1
12th	%	71.5	7.8	7.0	4.1	3.7	2.2	3.7	28.5
(n=1,063)	CI	68.7–74.1	6.3–9.5	5.6–8.7	3.1–5.5	2.7–5.0	1.4–3.2	2.8–5.1	25.9–31.3

Note: The confidence interval (CI) is 95 percent.

Table BH3.4. Lifetime Cigarette Use

		On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.7	9.7	2.5	1.8	1.4	1.0	3.8	20.3
(n=5,171)	CI	78.5–80.9	8.9–10.6	2.0–3.0	1.5–2.3	1.1–1.8	0.7–1.4	3.3–4.5	19.1–21.5
Male	%	80.6	8.7	2.2	1.8	1.5	0.9	4.2	19.4
(n=2,586)	CI	78.9–82.2	7.6–9.9	1.6–2.9	1.3–2.5	1.1–2.2	0.6–1.4	3.4–5.2	17.8–21.1
Female	%	79.2	10.6	2.8	1.9	1.2	1.0	3.3	20.8
(n=2,460)	CI	77.4–80.9	9.4–12.0	2.2–3.6	1.3–2.6	0.8–1.8	0.6–1.6	2.6–4.2	19.1–22.6
White	%	77.4	10.4	2.7	2.1	1.5	1.3	4.6	22.6
(n=3,593)	CI	75.8–78.8	9.4–11.5	2.2–3.3	1.6–2.7	1.1–2.0	0.9–1.8	3.9–5.5	21.2–24.2
Black	%	88.1	6.7	1.4	1.0	1.3	0.2	1.3	11.9
(n=495)	CI	84.6–90.9	4.7–9.5	0.6–3.1	0.4–2.5	0.5–3.0	0.0–1.5	0.5–3.3	9.1–15.4
Hispanic	%	88.1	7.8	1.6	1.3	0.2	0.1	0.8	11.9
(n=469)	CI	84.4–91.0	5.4–11.0	0.7–3.5	0.5–3.2	0.0–1.7	0.0–1.1	0.3–2.5	9.0–15.6
8th	%	88.3	7.1	1.6	0.5	0.8	0.5	1.1	11.7
(n=2,548)	CI	87.0–89.5	6.2–8.2	1.2–2.2	0.3–0.9	0.5–1.3	0.3–0.8	0.8–1.6	10.5–13.0
10th	%	79.4	9.8	2.4	2.1	1.7	1.2	3.5	20.6
(n=1,561)	CI	77.3–81.3	8.4–11.4	1.7–3.3	1.5–3.0	1.1–2.4	0.7–1.8	2.7–4.6	18.7–22.7
12th	%	70.7	12.5	3.5	2.9	1.9	1.4	7.1	29.3
(n=1,062)	CI	67.9–73.4	10.6–14.6	2.5–4.8	2.1–4.1	1.2–2.9	0.8–2.3	5.7–8.8	26.6–32.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	70.7	9.7	3.9	2.6	2.3	1.7	9.2	29.3
(n=5,163)	CI	69.4–72.0	8.8–10.5	3.4–4.5	2.1–3.1	1.9–2.7	1.3–2.1	8.4–10.1	28.0–30.6
Male	%	71.1	8.8	3.6	2.3	2.2	1.6	10.5	28.9
(n=2,582)	CI	69.2–73.0	7.7–10.0	2.9–4.4	1.7–3.0	1.6–2.8	1.1–2.2	9.2–11.9	27.0–30.8
Female	%	70.8	10.5	4.3	2.6	2.4	1.8	7.6	29.2
(n=2,457)	CI	68.8–72.7	9.2–11.8	3.5–5.2	2.0–3.4	1.8–3.2	1.3–2.5	6.5–8.9	27.3–31.2
White	%	67.4	10.1	4.1	2.7	2.7	1.9	11.0	32.6
(n=3,590)	CI	65.8–69.1	9.1–11.2	3.5–4.9	2.2–3.4	2.2–3.3	1.5–2.5	9.9–12.2	30.9–34.2
Black	%	86.8	5.6	2.2	1.5	1.2	0.1	2.5	13.2
(n=492)	CI	83.3–89.7	3.8–8.2	1.2–4.0	0.7–3.3	0.5–2.9	0.0–1.0	1.4–4.5	10.3–16.7
Hispanic	%	81.3	7.7	3.5	1.5	1.1	2.1	2.9	18.7
(n=468)	CI	77.1–84.8	5.5–10.6	2.0–5.9	0.7–3.5	0.5–2.5	1.0–4.3	1.6–5.1	15.2–22.9
8th	%	80.7	8.1	2.9	2.0	1.8	1.1	3.3	19.3
(n=2,541)	CI	79.1–82.2	7.1–9.3	2.3–3.6	1.5–2.6	1.3–2.4	0.8–1.6	2.7–4.1	17.8–20.9
10th	%	65.9	11.3	4.4	2.4	2.9	1.9	11.2	34.1
(n=1,559)	CI	63.5–68.2	9.8–12.9	3.4–5.5	1.8–3.3	2.2–3.8	1.3–2.7	9.7–12.9	31.8–36.5
12th	%	65.0	9.6	4.5	3.3	2.2	2.1	13.4	35.0
(n=1,063)	CI	62.1–67.9	7.9–11.5	3.4–5.9	2.4–4.5	1.4–3.2	1.4–3.1	11.5–15.6	32.1–37.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	88.7	4.2	1.5	0.9	0.8	0.5	3.5	11.3
(n=5,149)	CI	87.7–89.6	3.7–4.9	1.1–1.9	0.6–1.2	0.6–1.1	0.3–0.7	2.9–4.1	10.4–12.3
Male	%	84.2	5.0	1.9	1.3	1.4	0.7	5.5	15.8
(n=2,579)	CI	82.6–85.7	4.2–6.0	1.4–2.6	0.9–1.9	1.0–2.0	0.4–1.1	4.6–6.7	14.3–17.4
Female	%	93.6	3.5	1.0	0.4	0.3	0.2	0.9	6.4
(n=2,446)	CI	92.5–94.6	2.7–4.4	0.7–1.6	0.2–0.8	0.1–0.7	0.1–0.5	0.6–1.5	5.4–7.5
White	%	86.5	4.8	1.8	0.9	1.0	0.6	4.5	13.5
(n=3,575)	CI	85.2–87.7	4.0–5.6	1.3–2.3	0.6–1.3	0.7–1.4	0.4–0.9	3.8–5.4	12.3–14.8
Black	%	96.1	2.4	0.5	0.0	0.2	0.0	0.7	3.9
(n=492)	CI	93.7–97.6	1.3–4.4	0.1–2.1	–	0.0–1.6	–	0.2–2.3	2.4–6.3
Hispanic	%	96.1	2.1	0.0	0.8	0.4	0.3	0.2	3.9
(n=468)	CI	93.7–97.7	1.1–4.1	–	0.3–2.5	0.1–1.6	0.0–2.3	0.0–1.7	2.3–6.3
8th	%	94.5	2.8	0.8	0.5	0.2	0.4	0.8	5.5
(n=2,542)	CI	93.5–95.3	2.2–3.5	0.5–1.2	0.3–0.9	0.1–0.5	0.2–0.7	0.5–1.3	4.7–6.5
10th	%	87.5	4.9	1.5	0.8	1.3	0.5	3.5	12.5
(n=1,552)	CI	85.8–89.1	3.9–6.1	1.0–2.2	0.5–1.4	0.8–2.0	0.2–0.9	2.7–4.6	10.9–14.2
12th	%	83.7	5.1	2.2	1.3	0.9	0.6	6.2	16.3
(n=1,055)	CI	81.3–85.8	3.9–6.6	1.4–3.2	0.8–2.2	0.5–1.7	0.3–1.3	4.9–7.9	14.2–18.7

Note: The confidence interval (CI) is 95 percent.

Table BH3.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.0	6.3	3.2	1.8	1.9	1.4	6.3	21.0
(n=5,158)	CI	77.8–80.2	5.7–7.1	2.7–3.8	1.4–2.2	1.6–2.4	1.1–1.8	5.6–7.1	19.8–22.2
Male	%	79.3	6.0	2.4	1.7	2.0	1.4	7.1	20.7
(n=2,577)	CI	77.6–81.0	5.1–7.0	1.9–3.2	1.3–2.4	1.5–2.7	1.0–2.0	6.1–8.3	19.0–22.4
Female	%	79.1	6.7	3.7	1.9	1.8	1.4	5.4	20.9
(n=2,456)	CI	77.2–80.8	5.7–7.9	3.0–4.6	1.3–2.6	1.3–2.6	1.0–2.0	4.5–6.5	19.2–22.8
White	%	80.3	5.7	3.0	1.6	1.8	1.4	6.2	19.7
(n=3,588)	CI	78.8–81.7	5.0–6.6	2.4–3.7	1.2–2.1	1.4–2.4	1.0–1.9	5.3–7.1	18.3–21.2
Black	%	72.1	9.5	4.3	3.0	2.3	1.2	7.5	27.9
(n=488)	CI	67.5–76.3	7.1–12.8	2.6–7.0	1.7–5.2	1.2–4.3	0.5–2.9	5.2–10.7	23.7–32.5
Hispanic	%	84.4	5.8	2.6	1.2	2.5	1.0	2.6	15.6
(n=467)	CI	80.3–87.8	3.9–8.8	1.4–4.7	0.4–3.1	1.3–4.7	0.3–2.7	1.4–5.1	12.2–19.7
8th	%	89.7	4.3	2.2	0.8	0.8	0.7	1.5	10.3
(n=2,540)	CI	88.5–90.9	3.6–5.2	1.7–2.8	0.5–1.2	0.5–1.3	0.5–1.2	1.1–2.0	9.1–11.5
10th	%	76.8	7.3	2.9	2.2	2.2	1.6	7.0	23.2
(n=1,554)	CI	74.6–78.8	6.1–8.7	2.2–3.9	1.6–3.0	1.6–3.0	1.1–2.4	5.8–8.4	21.2–25.4
12th	%	69.9	7.5	4.7	2.3	2.9	2.0	10.7	30.1
(n=1,064)	CI	67.1–72.6	6.1–9.2	3.6–6.1	1.6–3.4	2.0–4.1	1.3–3.0	9.0–12.7	27.4–32.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	98.6	0.6	0.2	0.1	0.1	0.1	0.3	1.4
(n=5,157)	CI	98.2–99.0	0.4–0.8	0.1–0.4	0.0–0.3	0.0–0.3	0.0–0.2	0.2–0.6	1.0–1.8
Male	%	98.8	0.5	0.3	0.1	0.1	0.0	0.2	1.2
(n=2,578)	CI	98.2–99.2	0.3–0.9	0.1–0.7	0.0–0.4	0.0–0.3	–	0.1–0.5	0.8–1.8
Female	%	98.9	0.5	0.0	0.1	0.2	0.1	0.2	1.1
(n=2,455)	CI	98.3–99.2	0.3–1.0	0.0–0.2	0.0–0.4	0.1–0.6	0.0–0.2	0.1–0.6	0.8–1.7
White	%	98.9	0.5	0.1	0.1	0.1	0.0	0.2	1.1
(n=3,583)	CI	98.5–99.3	0.3–0.8	0.0–0.3	0.0–0.3	0.0–0.4	0.0–0.1	0.1–0.5	0.7–1.5
Black	%	98.4	1.2	0.0	0.0	0.0	0.2	0.2	1.6
(n=490)	CI	96.6–99.2	0.5–2.9	–	–	–	0.0–1.6	0.0–1.6	0.8–3.4
Hispanic	%	99.2	0.0	0.3	0.0	0.1	0.0	0.3	0.8
(n=469)	CI	97.4–99.8	–	0.0–2.3	–	0.0–1.1	–	0.0–2.3	0.2–2.6
8th	%	99.3	0.4	0.0	0.0	0.1	0.1	0.0	0.7
(n=2,539)	CI	98.9–99.6	0.2–0.7	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.3	0.4–1.1
10 th	%	98.5	0.6	0.1	0.1	0.1	0.1	0.4	1.5
(n=1,559)	CI	97.8–99.0	0.3–1.2	0.0–0.5	0.0–0.5	0.0–0.5	0.0–0.5	0.2–0.9	1.0–2.2
12th	%	98.0	0.7	0.4	0.2	0.2	0.0	0.6	2.0
(n=1,059)	CI	97.0–98.7	0.3–1.4	0.1–1.0	0.0–0.8	0.0–0.8	–	0.3–1.3	1.3–3.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	97.0	1.2	0.6	0.4	0.3	0.1	0.4	3.0
(n=5,167)	CI	96.4–97.5	0.9–1.5	0.4–0.9	0.3–0.7	0.2–0.5	0.0–0.3	0.3–0.7	2.5–3.6
Male	%	97.0	1.2	0.7	0.5	0.4	0.0	0.3	3.0
(n=2,583)	CI	96.2–97.7	0.8–1.7	0.4–1.1	0.3–0.9	0.2–0.8	–	0.1–0.6	2.3–3.8
Female	%	97.2	1.2	0.5	0.3	0.2	0.2	0.5	2.8
(n=2,460)	CI	96.3–97.8	0.8–1.8	0.3–1.0	0.1–0.7	0.0–0.5	0.1–0.5	0.2–0.9	2.2–3.7
White	%	97.2	1.1	0.7	0.3	0.3	0.1	0.3	2.8
(n=3,591)	CI	96.5–97.7	0.8–1.5	0.4–1.1	0.2–0.6	0.2–0.6	0.0–0.3	0.2–0.6	2.3–3.5
Black	%	98.3	0.6	0.5	0.3	0.0	0.0	0.2	1.7
(n=490)	CI	96.6–99.2	0.2–1.7	0.1–2.1	0.0–2.1	–	–	0.0–1.6	0.8–3.4
Hispanic	%	97.6	1.3	0.0	1.0	0.1	0.0	0.0	2.4
(n=470)	CI	95.4–98.8	0.5–3.0	–	0.3–3.0	0.0–1.1	–	–	1.2–4.6
8th	%	98.5	0.8	0.2	0.2	0.1	0.0	0.1	1.5
(n=2,546)	CI	98.0–98.9	0.5–1.3	0.1–0.5	0.1–0.5	0.0–0.3	0.0–0.3	0.0–0.3	1.1–2.0
10 th	%	97.1	1.0	0.5	0.2	0.4	0.2	0.6	2.9
(n=1,561)	CI	96.1–97.8	0.6–1.7	0.3–1.0	0.1–0.6	0.2–0.9	0.1–0.6	0.3–1.2	2.2–3.9
12th	%	95.2	1.7	1.0	0.9	0.5	0.1	0.6	4.8
(n=1,060)	CI	93.7–96.3	1.1–2.7	0.6–1.9	0.5–1.7	0.2–1.1	0.0–0.7	0.3–1.3	3.7–6.3

Note: The confidence interval (CI) is 95 percent.

Table BH3.10. Age of First Prescription Drug Misuse

How old were you when you first <u>used prescription drugs not prescribed to you</u> ?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	92.8	1.1	0.5	0.7	1.4	1.0	1.1	0.6	0.8	13.42
(n=5,180)	CI	92.0–93.5	0.8–1.4	0.4–0.8	0.5–1.0	1.1–1.8	0.7–1.3	0.8–1.5	0.4–0.9	0.5–1.1	13.2–13.7
Male	%	94.3	1.0	0.3	0.5	1.0	0.6	0.9	0.6	0.8	13.61
(n=2,592)	CI	93.2–95.2	0.7–1.4	0.1–0.6	0.3–0.9	0.7–1.6	0.3–1.0	0.6–1.5	0.4–1.1	0.5–1.4	13.2–14.0
Female	%	91.6	1.0	0.8	1.0	1.7	1.2	1.3	0.7	0.8	13.37
(n=2,463)	CI	90.4–92.7	0.7–1.5	0.5–1.2	0.7–1.4	1.3–2.4	0.8–1.8	0.8–1.9	0.4–1.2	0.4–1.3	13.1–13.7
White	%	93.2	0.9	0.5	0.6	1.4	0.9	1.2	0.7	0.6	13.51
(n=3,603)	CI	92.2–94.0	0.7–1.3	0.3–0.8	0.4–0.9	1.0–1.8	0.6–1.2	0.8–1.7	0.5–1.1	0.4–1.0	13.2–13.8
Black	%	93.3	0.8	0.6	1.2	1.3	0.2	0.5	0.8	1.2	13.57
(n=492)	CI	90.4–95.4	0.3–2.2	0.2–1.9	0.5–2.8	0.6–3.1	0.0–1.5	0.1–2.1	0.3–2.6	0.5–3.2	12.6–14.5
Hispanic	%	93.3	0.5	0.1	0.7	0.8	2.0	0.9	0.0	1.6	0.00
(n=469)	CI	90.3–95.4	0.2–1.7	0.0–1.1	0.3–1.9	0.3–2.1	1.0–4.1	0.3–2.8	–	0.7–3.7	–
8th	%	94.8	1.3	1.0	1.0	1.3	0.6	0.0	0.0	0.0	11.82
(n=2,551)	CI	93.9–95.6	0.9–1.8	0.7–1.5	0.7–1.4	0.9–1.8	0.4–1.0	0.0–0.3	–	–	11.6–12.1
10th	%	92.1	1.4	0.3	1.0	1.3	1.3	1.8	0.6	0.1	13.21
(n=1,567)	CI	90.7–93.3	0.9–2.1	0.1–0.8	0.6–1.6	0.8–2.0	0.9–2.0	1.3–2.6	0.3–1.2	0.0–0.5	12.9–13.6
12th	%	91.4	0.6	0.3	0.2	1.6	0.9	1.5	1.3	2.3	14.67
(n=1,062)	CI	89.5–92.9	0.3–1.2	0.1–0.9	0.0–0.7	1.0–2.6	0.5–1.7	0.9–2.4	0.8–2.2	1.5–3.3	14.2–15.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.11. Age of First Alcohol Use

How old were you when you first <u>had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)</u> ?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	63.1	5.1	2.2	3.5	5.4	5.6	6.5	5.7	2.9	13.65
(n=5,166)	CI	61.6–64.5	4.5–5.8	1.8–2.7	3.0–4.0	4.8–6.1	5.0–6.3	5.7–7.3	5.0–6.5	2.4–3.6	13.5–13.8
Male	%	65.3	6.0	2.0	3.4	5.0	5.1	5.8	4.6	2.8	13.47
(n=2,584)	CI	63.3–67.3	5.1–7.0	1.5–2.6	2.7–4.2	4.3–6.0	4.2–6.1	4.8–6.9	3.7–5.6	2.1–3.7	13.3–13.6
Female	%	61.2	4.0	2.3	3.4	5.9	5.8	7.2	7.1	3.1	13.88
(n=2,457)	CI	59.2–63.3	3.2–4.8	1.8–3.0	2.8–4.2	5.0–6.9	4.9–6.9	6.1–8.5	6.0–8.4	2.4–4.1	13.7–14.0
White	%	61.0	5.3	2.0	3.7	5.3	6.3	7.4	6.0	3.0	13.70
(n=3,596)	CI	59.3–62.7	4.6–6.1	1.6–2.6	3.1–4.4	4.7–6.1	5.5–7.2	6.4–8.4	5.1–7.0	2.4–3.8	13.6–13.8
Black	%	73.6	3.1	2.4	3.3	2.9	3.1	3.6	4.8	3.3	13.78
(n=493)	CI	69.2–77.5	1.9–5.0	1.4–4.1	2.1–5.2	1.7–4.8	1.8–5.3	2.1–6.0	3.0–7.6	1.8–5.9	13.3–14.2
Hispanic	%	71.7	3.4	2.3	3.2	5.5	2.9	2.7	4.9	3.3	13.65
(n=466)	CI	67.0–75.9	2.0–5.7	1.3–4.1	1.9–5.4	3.7–8.1	1.6–5.1	1.4–5.0	3.0–7.9	1.8–6.1	13.2–14.1
8th	%	76.4	6.0	3.3	4.6	7.4	2.2	0.1	0.0	0.0	11.86
(n=2,543)	CI	74.7–78.0	5.2–7.0	2.7–4.1	3.9–5.5	6.4–8.4	1.7–2.8	0.0–0.4	–	–	11.8–12.0
10th	%	59.6	5.2	1.6	3.4	5.8	9.3	11.0	4.0	0.1	13.53
(n=1,563)	CI	57.1–62.0	4.2–6.5	1.1–2.4	2.6–4.4	4.8–7.1	8.0–10.9	9.5–12.6	3.1–5.1	0.0–0.5	13.4–13.7
12th	%	52.4	4.0	1.7	2.3	2.9	5.4	8.5	13.6	9.1	14.72
(n=1,060)	CI	49.4–55.4	3.0–5.4	1.1–2.7	1.6–3.5	2.1–4.1	4.2–6.9	6.9–10.3	11.7–15.8	7.5–11.0	14.5–14.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.12. Age of First Cigarette Use

		How old were you when you first <u>smoked a cigarette, even just a puff?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	80.2	3.8	1.7	2.5	2.7	2.3	2.6	2.7	1.4	13.21
(n=5,166)	CI	79.0–81.4	3.3–4.3	1.4–2.1	2.1–3.0	2.3–3.2	1.9–2.8	2.2–3.2	2.2–3.2	1.1–1.9	13.0–13.4
Male	%	81.1	4.0	1.6	2.4	2.9	2.0	2.1	2.6	1.1	13.05
(n=2,584)	CI	79.4–82.7	3.3–4.9	1.2–2.2	1.9–3.2	2.3–3.7	1.5–2.8	1.6–2.9	2.0–3.5	0.7–1.8	12.8–13.3
Female	%	79.7	3.2	1.8	2.4	2.6	2.6	3.2	2.8	1.8	13.43
(n=2,457)	CI	77.9–81.4	2.6–4.0	1.3–2.4	1.9–3.1	2.0–3.3	2.0–3.4	2.5–4.1	2.1–3.7	1.2–2.5	13.2–13.7
White	%	77.8	4.1	1.7	2.7	3.0	2.7	3.1	3.3	1.5	13.28
(n=3,590)	CI	76.3–79.3	3.5–4.9	1.3–2.2	2.2–3.4	2.5–3.6	2.2–3.4	2.5–3.8	2.7–4.1	1.1–2.1	13.1–13.5
Black	%	89.4	1.7	0.8	1.8	0.9	1.0	2.0	1.2	1.1	13.42
(n=491)	CI	86.1–92.1	0.9–3.4	0.3–2.1	0.9–3.6	0.3–2.4	0.4–2.6	1.0–4.0	0.5–3.2	0.4–3.0	12.7–14.1
Hispanic	%	88.5	0.6	0.4	2.7	2.4	1.6	0.6	0.9	2.3	13.81
(n=469)	CI	84.8–91.3	0.2–2.0	0.1–1.6	1.5–4.9	1.3–4.4	0.7–3.6	0.1–2.3	0.3–2.8	1.1–4.8	13.1–14.5
8th	%	88.9	3.4	2.0	2.2	3.1	0.5	0.0	0.0	0.0	11.59
(n=2,545)	CI	87.6–90.1	2.7–4.2	1.5–2.6	1.7–2.8	2.5–3.8	0.3–0.8	0.0–0.3	–	–	11.4–11.7
10th	%	79.9	4.9	1.5	2.7	3.1	2.9	4.0	1.0	0.1	12.69
(n=1,562)	CI	77.9–81.8	3.9–6.0	1.0–2.2	2.0–3.7	2.3–4.1	2.2–3.9	3.1–5.1	0.6–1.6	0.0–0.5	12.5–12.9
12th	%	71.1	3.0	1.6	2.7	2.1	3.7	4.0	7.3	4.4	14.26
(n=1,059)	CI	68.3–73.8	2.1–4.2	1.0–2.6	1.9–3.9	1.4–3.1	2.7–5.0	3.0–5.4	5.9–9.1	3.3–5.8	14.0–14.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	72.2	1.1	1.1	2.5	4.7	4.4	6.2	4.7	3.2	14.28
(n=5,146)	CI	70.9–73.5	0.8–1.4	0.8–1.4	2.1–2.9	4.2–5.3	3.9–5.1	5.5–6.9	4.0–5.4	2.6–3.8	14.2–14.4
Male	%	72.6	1.2	1.0	3.0	4.4	4.5	6.2	4.5	2.5	14.16
(n=2,569)	CI	70.7–74.4	0.8–1.7	0.7–1.5	2.4–3.8	3.7–5.3	3.7–5.4	5.2–7.3	3.7–5.6	1.9–3.4	14.0–14.3
Female	%	72.3	0.8	1.1	1.9	5.0	4.3	6.0	4.8	3.8	14.42
(n=2,452)	CI	70.3–74.1	0.6–1.3	0.7–1.5	1.4–2.5	4.3–5.9	3.5–5.2	5.1–7.2	3.9–5.9	3.0–4.9	14.3–14.6
White	%	68.7	1.1	1.2	2.7	5.0	5.1	7.0	5.6	3.6	14.34
(n=3,576)	CI	67.1–70.3	0.8–1.5	0.9–1.6	2.3–3.3	4.3–5.7	4.4–5.9	6.1–8.0	4.8–6.6	3.0–4.5	14.2–14.5
Black	%	88.5	0.2	0.4	1.6	2.6	1.2	2.5	1.9	1.1	14.19
(n=493)	CI	85.2–91.2	0.0–1.5	0.1–1.5	0.8–3.1	1.6–4.2	0.5–2.8	1.3–4.7	0.9–4.1	0.4–3.0	13.7–14.7
Hispanic	%	83.1	0.5	0.2	2.1	2.4	2.8	3.3	3.0	2.7	14.49
(n=467)	CI	79.1–86.5	0.1–2.1	0.0–1.1	1.1–3.9	1.4–4.1	1.6–4.8	1.9–5.7	1.6–5.5	1.3–5.2	14.0–14.9
8th	%	82.1	1.4	1.9	4.0	8.0	2.5	0.2	0.0	0.0	12.51
(n=2,535)	CI	80.5–83.5	1.0–1.9	1.4–2.5	3.3–4.8	7.0–9.1	2.0–3.2	0.1–0.4	–	0.0–0.3	12.4–12.6
10th	%	67.6	1.2	0.8	2.2	3.7	7.6	12.5	3.9	0.3	14.18
(n=1,554)	CI	65.2–69.9	0.7–1.8	0.5–1.4	1.6–3.1	2.9–4.8	6.4–9.1	11.0–14.3	3.1–5.0	0.1–0.8	14.1–14.3
12th	%	66.4	0.7	0.5	1.1	2.3	3.1	5.8	10.5	9.6	15.40
(n=1,057)	CI	63.5–69.2	0.3–1.4	0.2–1.1	0.6–2.0	1.5–3.4	2.2–4.3	4.6–7.4	8.8–12.5	8.0–11.5	15.2–15.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	79.3	1.4	1.0	2.2	3.2	3.6	4.0	3.1	2.1	14.00
(n=5,183)	CI	78.1–80.5	1.1–1.8	0.8–1.4	1.8–2.6	2.8–3.8	3.1–4.2	3.5–4.7	2.6–3.7	1.7–2.6	13.9–14.1
Male	%	79.7	1.7	1.4	2.5	3.0	3.4	3.9	2.6	1.9	13.81
(n=2,593)	CI	78.0–81.4	1.2–2.3	1.0–1.9	1.9–3.2	2.4–3.8	2.7–4.2	3.1–4.9	2.0–3.5	1.3–2.6	13.6–14.0
Female	%	79.2	1.0	0.7	1.8	3.4	3.8	4.1	3.6	2.4	14.24
(n=2,465)	CI	77.4–80.9	0.7–1.5	0.5–1.2	1.3–2.4	2.7–4.2	3.0–4.7	3.3–5.1	2.8–4.5	1.7–3.2	14.1–14.4
White	%	80.5	1.1	0.9	2.3	2.9	3.2	4.0	2.8	2.2	14.07
(n=3,603)	CI	79.0–81.9	0.8–1.5	0.7–1.4	1.9–2.9	2.4–3.5	2.6–3.9	3.3–4.8	2.3–3.6	1.7–2.9	13.9–14.2
Black	%	72.6	2.6	1.0	2.3	3.9	4.0	4.5	5.5	3.6	14.15
(n=493)	CI	68.1–76.7	1.5–4.7	0.4–2.5	1.3–4.3	2.5–6.0	2.4–6.4	2.8–7.1	3.5–8.4	2.1–6.3	13.7–14.6
Hispanic	%	84.8	0.4	0.5	1.0	2.9	3.2	3.7	2.6	1.0	14.27
(n=469)	CI	80.7–88.1	0.1–1.6	0.1–2.1	0.3–2.7	1.6–5.0	1.8–5.6	2.1–6.3	1.4–5.0	0.3–3.1	13.8–14.7
8th	%	89.9	1.2	1.1	2.5	3.9	1.4	0.0	0.0	0.0	12.32
(n=2,555)	CI	88.6–91.0	0.9–1.7	0.8–1.6	1.9–3.1	3.2–4.7	1.0–1.9	0.0–0.3	–	–	12.2–12.5
10th	%	77.1	1.7	1.1	2.4	3.1	5.4	7.1	2.1	0.0	13.70
(n=1,564)	CI	74.9–79.1	1.2–2.5	0.7–1.8	1.7–3.2	2.4–4.1	4.4–6.6	5.9–8.5	1.5–3.0	–	13.5–13.9
12th	%	70.3	1.3	0.8	1.7	2.6	4.1	5.1	7.5	6.6	14.87
(n=1,064)	CI	67.5–73.0	0.8–2.2	0.4–1.6	1.1–2.7	1.8–3.8	3.1–5.5	3.9–6.5	6.1–9.2	5.2–8.2	14.7–15.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.15. Past Month Prescription Drug Misuse

		During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	96.4	1.9	0.9	0.3	0.1	0.1	0.2	3.6
(n=5,191)	CI	95.8–96.9	1.5–2.3	0.7–1.3	0.2–0.6	0.1–0.3	0.0–0.2	0.1–0.4	3.1–4.2
Male	%	96.8	1.5	1.0	0.3	0.1	0.1	0.2	3.2
(n=2,599)	CI	95.9–97.4	1.1–2.1	0.6–1.5	0.1–0.7	0.0–0.3	0.0–0.4	0.1–0.5	2.6–4.1
Female	%	96.5	2.1	0.7	0.2	0.2	0.1	0.1	3.5
(n=2,466)	CI	95.7–97.2	1.6–2.8	0.5–1.2	0.1–0.5	0.1–0.4	0.0–0.3	0.0–0.4	2.8–4.3
White	%	96.9	1.8	0.6	0.2	0.1	0.1	0.2	3.1
(n=3,610)	CI	96.3–97.5	1.4–2.3	0.4–0.9	0.1–0.5	0.1–0.3	0.0–0.3	0.1–0.5	2.5–3.7
Black	%	95.8	2.7	1.1	0.4	0.0	0.0	0.0	4.2
(n=493)	CI	93.3–97.4	1.4–4.8	0.5–2.7	0.1–1.9	–	–	–	2.6–6.7
Hispanic	%	97.1	0.8	1.5	0.5	0.0	0.0	0.0	2.9
(n=470)	CI	95.0–98.4	0.3–2.2	0.7–3.5	0.2–1.7	–	–	–	1.6–5.0
8th	%	96.9	1.5	0.8	0.3	0.2	0.2	0.1	3.1
(n=2,555)	CI	96.1–97.5	1.1–2.1	0.5–1.2	0.2–0.6	0.1–0.5	0.1–0.5	0.0–0.3	2.5–3.9
10th	%	96.4	1.9	1.1	0.3	0.2	0.0	0.2	3.6
(n=1,568)	CI	95.3–97.2	1.3–2.7	0.7–1.7	0.1–0.7	0.1–0.6	–	0.1–0.6	2.8–4.7
12th	%	96.0	2.2	0.9	0.5	0.0	0.1	0.3	4.0
(n=1,068)	CI	94.6–97.0	1.5–3.3	0.5–1.7	0.2–1.1	–	0.0–0.7	0.1–0.9	3.0–5.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.16. Past Month Alcohol Use

		During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	85.2	8.3	3.0	1.3	1.3	0.4	0.5	14.8
(n=5,176)	CI	84.1–86.2	7.6–9.2	2.5–3.5	1.0–1.7	1.0–1.7	0.3–0.7	0.3–0.7	13.8–15.9
Male	%	86.5	7.3	2.5	1.5	1.5	0.3	0.4	13.5
(n=2,589)	CI	85.0–87.9	6.2–8.4	1.9–3.3	1.1–2.1	1.1–2.2	0.1–0.7	0.2–0.8	12.1–15.0
Female	%	84.2	9.4	3.4	1.2	0.9	0.5	0.5	15.8
(n=2,460)	CI	82.5–85.7	8.2–10.8	2.7–4.3	0.8–1.7	0.5–1.4	0.3–0.9	0.2–0.9	14.3–17.5
White	%	85.1	8.3	2.9	1.3	1.4	0.5	0.5	14.9
(n=3,601)	CI	83.8–86.3	7.3–9.3	2.3–3.6	0.9–1.8	1.0–1.9	0.3–0.9	0.3–0.9	13.7–16.2
Black	%	88.0	9.2	1.3	1.3	0.2	0.0	0.0	12.0
(n=494)	CI	84.5–90.8	6.7–12.5	0.6–2.8	0.5–3.0	0.0–1.5	–	–	9.2–15.5
Hispanic	%	87.5	7.8	2.7	1.1	0.3	0.0	0.6	12.5
(n=468)	CI	83.8–90.4	5.6–10.8	1.4–5.1	0.4–3.1	0.0–2.3	–	0.1–2.3	9.6–16.2
8th	%	91.0	5.9	1.2	0.8	0.6	0.2	0.3	9.0
(n=2,547)	CI	89.8–92.0	5.0–6.9	0.9–1.7	0.5–1.2	0.4–1.0	0.1–0.5	0.1–0.6	8.0–10.2
10th	%	85.6	8.4	2.6	1.3	1.2	0.4	0.6	14.4
(n=1,563)	CI	83.8–87.3	7.1–9.9	1.9–3.5	0.8–2.0	0.7–1.8	0.2–0.9	0.3–1.2	12.7–16.2
12th	%	78.6	10.9	5.2	2.0	2.1	0.7	0.6	21.4
(n=1,066)	CI	76.1–81.0	9.2–12.9	4.0–6.7	1.3–3.0	1.4–3.1	0.3–1.4	0.3–1.2	19.0–23.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.17. Past Month Binge Drinking

		During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	92.7	3.7	1.6	0.8	0.6	0.3	0.3	7.3
(n=5,182)	CI	91.9–93.5	3.2–4.4	1.3–2.1	0.5–1.1	0.4–0.9	0.2–0.5	0.2–0.5	6.5–8.1
Male	%	92.3	4.2	1.7	0.9	0.5	0.2	0.2	7.7
(n=2,595)	CI	91.0–93.3	3.4–5.2	1.2–2.4	0.6–1.4	0.3–0.9	0.1–0.6	0.1–0.6	6.7–9.0
Female	%	93.3	3.4	1.5	0.6	0.7	0.3	0.2	6.7
(n=2,460)	CI	92.1–94.4	2.7–4.3	1.0–2.2	0.3–1.0	0.4–1.2	0.2–0.7	0.1–0.5	5.6–7.9
White	%	92.3	3.8	1.7	0.8	0.8	0.3	0.3	7.7
(n=3,605)	CI	91.2–93.2	3.1–4.6	1.3–2.2	0.6–1.3	0.5–1.2	0.2–0.6	0.1–0.6	6.8–8.8
Black	%	95.7	2.7	1.2	0.2	0.2	0.0	0.0	4.3
(n=492)	CI	93.2–97.3	1.5–4.8	0.5–2.9	0.0–1.6	0.0–1.6	–	–	2.7–6.8
Hispanic	%	95.4	2.1	1.5	0.3	0.0	0.0	0.6	4.6
(n=470)	CI	92.6–97.2	1.0–4.5	0.7–3.5	0.0–2.3	–	–	0.1–2.3	2.8–7.4
8th	%	97.3	1.4	0.7	0.2	0.2	0.2	0.2	2.7
(n=2,551)	CI	96.5–97.8	1.0–1.9	0.4–1.1	0.1–0.5	0.1–0.4	0.1–0.4	0.1–0.4	2.2–3.5
10th	%	93.0	4.1	1.1	0.7	0.4	0.3	0.3	7.0
(n=1,564)	CI	91.7–94.2	3.2–5.2	0.7–1.7	0.4–1.3	0.2–0.9	0.1–0.8	0.1–0.8	5.8–8.3
12th	%	87.5	5.9	3.2	1.4	1.2	0.5	0.4	12.5
(n=1,067)	CI	85.4–89.4	4.6–7.5	2.3–4.4	0.8–2.3	0.7–2.1	0.2–1.1	0.1–1.0	10.6–14.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.18. Past Month Cigarette Use

		During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	91.4	3.4	1.6	0.7	0.9	0.7	1.4	8.6
(n=5,177)	CI	90.5–92.2	2.9–3.9	1.2–2.0	0.5–1.0	0.7–1.2	0.5–1.0	1.1–1.9	7.8–9.5
Male	%	91.9	2.7	1.6	0.6	0.8	0.7	1.6	8.1
(n=2,592)	CI	90.7–93.0	2.1–3.5	1.1–2.3	0.4–1.1	0.5–1.3	0.4–1.1	1.1–2.3	7.0–9.3
Female	%	91.2	3.8	1.6	0.7	1.0	0.5	1.1	8.8
(n=2,458)	CI	89.9–92.4	3.1–4.7	1.1–2.3	0.4–1.2	0.7–1.6	0.3–1.0	0.7–1.7	7.6–10.1
White	%	90.5	3.5	1.8	0.8	1.0	0.9	1.5	9.5
(n=3,601)	CI	89.4–91.5	2.9–4.2	1.4–2.4	0.5–1.2	0.7–1.5	0.6–1.3	1.1–2.0	8.5–10.6
Black	%	95.1	2.0	1.0	0.5	0.3	0.2	0.9	4.9
(n=493)	CI	92.5–96.9	1.0–3.9	0.3–2.6	0.1–2.1	0.0–2.1	0.0–1.5	0.3–2.8	3.1–7.5
Hispanic	%	95.0	3.0	0.8	0.1	0.7	0.0	0.3	5.0
(n=470)	CI	92.3–96.8	1.7–5.3	0.3–2.1	0.0–1.1	0.2–2.3	–	0.0–2.3	3.2–7.7
8th	%	94.6	2.9	0.7	0.4	0.7	0.2	0.4	5.4
(n=2,549)	CI	93.7–95.4	2.3–3.6	0.5–1.2	0.2–0.7	0.5–1.2	0.1–0.5	0.2–0.7	4.6–6.3
10th	%	91.9	3.0	1.4	0.8	0.7	1.0	1.3	8.1
(n=1,564)	CI	90.4–93.1	2.3–4.0	0.9–2.1	0.4–1.3	0.4–1.3	0.6–1.6	0.8–2.0	6.9–9.6
12th	%	87.3	4.3	2.6	0.8	1.3	0.9	2.7	12.7
(n=1,064)	CI	85.1–89.1	3.2–5.7	1.8–3.8	0.4–1.6	0.8–2.2	0.5–1.7	1.9–3.9	10.9–14.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	82.9	5.1	2.8	1.5	2.1	1.5	4.2	17.1
(n=5,173)	CI	81.7–83.9	4.5–5.8	2.4–3.3	1.2–1.9	1.7–2.5	1.2–1.9	3.6–4.9	16.1–18.3
Male	%	83.4	4.5	2.2	1.4	1.6	2.0	4.8	16.6
(n=2,586)	CI	81.8–84.9	3.7–5.4	1.7–2.9	1.0–2.1	1.2–2.2	1.5–2.7	4.0–5.9	15.1–18.2
Female	%	82.5	5.6	3.5	1.6	2.4	0.9	3.5	17.5
(n=2,460)	CI	80.8–84.1	4.7–6.7	2.8–4.3	1.1–2.2	1.8–3.2	0.5–1.4	2.8–4.5	15.9–19.2
White	%	81.0	5.4	3.0	1.7	2.2	1.8	4.9	19.0
(n=3,595)	CI	79.6–82.4	4.7–6.3	2.4–3.6	1.2–2.2	1.8–2.8	1.4–2.4	4.1–5.8	17.6–20.4
Black	%	92.5	2.8	0.7	0.8	1.5	0.2	1.5	7.5
(n=494)	CI	89.5–94.7	1.6–4.8	0.2–2.4	0.3–2.2	0.7–3.3	0.0–1.5	0.7–3.3	5.3–10.5
Hispanic	%	90.2	3.0	3.1	0.9	1.2	0.7	0.9	9.8
(n=468)	CI	86.9–92.7	1.7–5.2	1.8–5.5	0.3–2.4	0.5–2.7	0.2–2.3	0.3–2.4	7.3–13.1
8th	%	89.5	3.9	2.3	0.8	1.5	0.9	1.2	10.5
(n=2,548)	CI	88.2–90.6	3.2–4.7	1.8–2.9	0.5–1.3	1.1–2.0	0.6–1.3	0.8–1.7	9.4–11.8
10th	%	80.1	6.1	3.6	1.7	2.4	1.7	4.4	19.9
(n=1,560)	CI	78.0–82.0	5.0–7.4	2.8–4.7	1.2–2.5	1.7–3.3	1.2–2.5	3.4–5.5	18.0–22.0
12th	%	78.7	5.3	2.4	2.1	2.3	1.9	7.3	21.3
(n=1,065)	CI	76.1–81.0	4.1–6.8	1.7–3.5	1.4–3.1	1.6–3.4	1.2–2.9	5.9–9.0	19.0–23.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	95.0	1.4	0.5	0.3	0.8	0.3	1.8	5.0
(n=5,188)	CI	94.3–95.6	1.1–1.8	0.3–0.7	0.1–0.5	0.6–1.1	0.2–0.6	1.4–2.2	4.4–5.7
Male	%	92.2	1.6	0.8	0.3	1.4	0.5	3.1	7.8
(n=2,598)	CI	91.0–93.3	1.2–2.2	0.5–1.3	0.2–0.7	1.0–2.1	0.3–0.9	2.4–3.9	6.7–9.0
Female	%	98.1	1.1	0.1	0.2	0.1	0.1	0.2	1.9
(n=2,463)	CI	97.5–98.6	0.7–1.7	0.0–0.4	0.1–0.5	0.0–0.3	0.0–0.3	0.1–0.6	1.4–2.5
White	%	94.0	1.5	0.5	0.3	1.0	0.4	2.2	6.0
(n=3,608)	CI	93.1–94.8	1.1–2.0	0.3–0.8	0.2–0.6	0.7–1.4	0.3–0.8	1.7–2.8	5.2–6.9
Black	%	98.7	0.7	0.3	0.0	0.0	0.0	0.2	1.3
(n=493)	CI	97.0–99.5	0.2–2.3	0.0–2.1	–	–	–	0.0–1.5	0.5–3.0
Hispanic	%	98.3	1.2	0.0	0.2	0.0	0.0	0.2	1.7
(n=469)	CI	96.4–99.2	0.5–3.0	–	0.0–1.7	–	–	0.0–1.7	0.8–3.6
8th	%	97.8	1.0	0.3	0.1	0.2	0.2	0.4	2.2
(n=2,554)	CI	97.2–98.3	0.7–1.4	0.2–0.6	0.0–0.4	0.1–0.5	0.1–0.4	0.2–0.7	1.7–2.8
10th	%	94.2	1.5	0.6	0.3	0.9	0.4	2.0	5.8
(n=1,567)	CI	92.9–95.3	1.0–2.3	0.3–1.2	0.1–0.8	0.5–1.5	0.2–0.8	1.4–2.9	4.7–7.1
12th	%	92.7	1.7	0.5	0.4	1.3	0.5	3.0	7.3
(n=1,067)	CI	91.0–94.1	1.1–2.7	0.2–1.1	0.1–1.0	0.8–2.2	0.2–1.1	2.1–4.2	5.9–9.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	88.1	4.1	1.9	1.3	1.2	1.0	2.5	11.9
(n=5,177)	CI	87.1-89.0	3.5-4.7	1.6-2.4	1.0-1.7	0.9-1.5	0.8-1.4	2.0-3.0	11.0-12.9
Male	%	88.8	3.0	1.5	1.4	0.8	1.1	3.4	11.2
(n=2,593)	CI	87.4-90.1	2.3-3.7	1.1-2.1	1.0-2.0	0.5-1.3	0.7-1.6	2.6-4.3	9.9-12.6
Female	%	87.6	5.0	2.3	1.2	1.4	1.0	1.5	12.4
(n=2,458)	CI	86.0-88.9	4.2-6.1	1.7-3.1	0.8-1.8	1.0-2.1	0.6-1.6	1.0-2.1	11.1-14.0
White	%	89.5	3.3	1.7	1.1	1.1	1.0	2.2	10.5
(n=3,603)	CI	88.4-90.6	2.7-4.0	1.3-2.2	0.8-1.6	0.8-1.6	0.7-1.4	1.8-2.9	9.4-11.6
Black	%	81.1	6.9	3.3	1.7	1.8	1.2	4.0	18.9
(n=489)	CI	76.8-84.7	4.8-10.0	1.9-5.6	0.8-3.6	0.8-3.8	0.5-3.0	2.5-6.5	15.3-23.2
Hispanic	%	91.5	3.8	1.5	1.1	0.3	0.3	1.5	8.5
(n=470)	CI	88.2-94.0	2.2-6.2	0.7-3.5	0.4-2.9	0.1-1.2	0.0-2.3	0.6-3.5	6.0-11.8
8th	%	93.6	3.1	1.2	0.4	0.9	0.5	0.4	6.4
(n=2,550)	CI	92.6-94.5	2.5-3.8	0.8-1.7	0.2-0.7	0.6-1.3	0.3-0.9	0.2-0.7	5.5-7.4
10th	%	87.1	4.5	1.8	1.7	1.0	0.9	3.1	12.9
(n=1,562)	CI	85.3-88.7	3.6-5.6	1.2-2.6	1.2-2.5	0.6-1.6	0.5-1.5	2.3-4.1	11.3-14.7
12th	%	83.3	4.7	2.9	1.8	1.7	1.7	4.0	16.7
(n=1,065)	CI	80.9-85.4	3.6-6.1	2.0-4.1	1.1-2.8	1.1-2.7	1.1-2.7	3.0-5.4	14.6-19.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.2	0.4	0.1	0.0	0.0	0.1	0.1	0.8
(n=5,195)	CI	98.9-99.5	0.2-0.6	0.1-0.3	0.0-0.1	0.0-0.2	0.0-0.2	0.0-0.3	0.5-1.1
Male	%	99.3	0.2	0.2	0.0	0.0	0.1	0.0	0.7
(n=2,601)	CI	98.9-99.6	0.1-0.6	0.1-0.6	0.0-0.3	-	0.0-0.4	0.0-0.3	0.4-1.1
Female	%	99.5	0.3	0.1	0.0	0.0	0.0	0.0	0.5
(n=2,467)	CI	99.0-99.7	0.1-0.7	0.0-0.4	-	0.0-0.3	0.0-0.3	0.0-0.2	0.3-1.0
White	%	99.4	0.3	0.1	0.0	0.0	0.0	0.1	0.6
(n=3,611)	CI	99.1-99.7	0.2-0.6	0.0-0.4	-	0.0-0.2	0.0-0.2	0.0-0.2	0.3-0.9
Black	%	99.7	0.0	0.3	0.0	0.0	0.0	0.0	0.3
(n=495)	CI	97.9-100.0	-	0.0-2.1	-	-	-	-	0.0-2.1
Hispanic	%	99.5	0.0	0.3	0.1	0.0	0.0	0.0	0.5
(n=470)	CI	97.9-99.9	-	0.0-2.3	0.0-1.1	-	-	-	0.1-2.1
8th	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=2,555)	CI	99.4-99.8	0.0-0.4	0.0-0.3	0.0-0.3	-	0.0-0.3	0.0-0.3	0.2-0.6
10th	%	99.2	0.3	0.0	0.1	0.1	0.1	0.2	0.8
(n=1,570)	CI	98.7-99.6	0.1-0.8	-	0.0-0.5	0.0-0.5	0.0-0.5	0.1-0.6	0.4-1.3
12th	%	98.8	0.7	0.4	0.0	0.0	0.1	0.1	1.2
(n=1,070)	CI	97.9-99.3	0.3-1.4	0.1-1.0	-	-	0.0-0.7	0.0-0.7	0.7-2.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.4	0.3	0.1	0.1	0.1	0.1	0.1	0.6
(n=5,195)	CI	99.1–99.6	0.1–0.5	0.1–0.3	0.0–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.4–0.9
Male	%	99.4	0.3	0.1	0.1	0.1	0.0	0.0	0.6
(n=2,602)	CI	98.9–99.6	0.1–0.6	0.0–0.3	0.0–0.5	0.0–0.4	–	–	0.4–1.1
Female	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=2,466)	CI	99.4–99.9	0.0–0.2	0.0–0.4	0.0–0.3	0.0–0.2	0.0–0.3	–	0.1–0.6
White	%	99.6	0.2	0.1	0.1	0.0	0.0	0.0	0.4
(n=3,611)	CI	99.3–99.7	0.1–0.4	0.0–0.2	0.0–0.3	0.0–0.3	0.0–0.2	0.0–0.2	0.3–0.7
Black	%	99.7	0.0	0.0	0.3	0.0	0.0	0.0	0.3
(n=495)	CI	97.9–100.0	–	–	0.0–2.1	–	–	–	0.0–2.1
Hispanic	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=470)	CI	98.8–99.9	0.0–1.1	0.0–1.1	–	–	–	–	0.1–1.2
8th	%	99.6	0.2	0.2	0.0	0.1	0.0	0.0	0.4
(n=2,554)	CI	99.2–99.8	0.1–0.4	0.1–0.4	–	0.0–0.3	0.0–0.3	–	0.2–0.8
10th	%	99.2	0.3	0.2	0.1	0.1	0.1	0.1	0.8
(n=1,571)	CI	98.7–99.6	0.1–0.7	0.1–0.6	0.0–0.5	0.0–0.5	0.0–0.5	0.0–0.5	0.4–1.3
12th	%	99.3	0.4	0.0	0.2	0.1	0.0	0.1	0.7
(n=1,070)	CI	98.5–99.6	0.1–1.0	–	0.0–0.7	0.0–0.7	–	0.0–0.7	0.4–1.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.9	0.5	0.3	0.2	0.1	0.1	0.1	1.1
(n=5,194)	CI	98.5–99.1	0.3–0.8	0.1–0.5	0.1–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.9–1.5
Male	%	98.6	0.5	0.4	0.2	0.1	0.1	0.0	1.4
(n=2,602)	CI	98.0–99.1	0.3–1.0	0.2–0.8	0.1–0.6	0.0–0.4	0.0–0.4	0.0–0.2	0.9–2.0
Female	%	99.4	0.3	0.2	0.0	0.0	0.0	0.0	0.6
(n=2,465)	CI	98.9–99.7	0.1–0.7	0.0–0.5	0.0–0.3	0.0–0.3	–	0.0–0.2	0.3–1.1
White	%	99.1	0.4	0.2	0.1	0.1	0.0	0.0	0.9
(n=3,610)	CI	98.6–99.4	0.3–0.8	0.1–0.5	0.0–0.4	0.0–0.2	0.0–0.3	0.0–0.2	0.6–1.4
Black	%	99.7	0.0	0.0	0.3	0.0	0.0	0.0	0.3
(n=495)	CI	97.9–100.0	–	–	0.0–2.1	–	–	–	0.0–2.1
Hispanic	%	99.0	0.3	0.5	0.1	0.0	0.0	0.0	1.0
(n=470)	CI	97.3–99.7	0.0–2.3	0.1–2.1	0.0–1.1	–	–	–	0.3–2.7
8th	%	99.6	0.2	0.0	0.0	0.0	0.0	0.1	0.4
(n=2,555)	CI	99.3–99.8	0.1–0.5	0.0–0.3	0.0–0.3	–	–	0.0–0.3	0.2–0.7
10th	%	98.9	0.4	0.2	0.1	0.3	0.1	0.1	1.1
(n=1,571)	CI	98.3–99.3	0.2–0.9	0.1–0.6	0.0–0.5	0.1–0.7	0.0–0.5	0.0–0.5	0.7–1.7
12th	%	97.9	0.9	0.6	0.4	0.0	0.1	0.1	2.1
(n=1,068)	CI	96.9–98.6	0.5–1.7	0.3–1.2	0.1–1.0	–	0.0–0.7	0.0–0.7	1.4–3.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.5	0.2	0.1	0.1	0.1	0.0	0.1	0.5
(n=5,195)	CI	99.3–99.7	0.1–0.4	0.0–0.2	0.0–0.2	0.0–0.2	0.0–0.1	0.0–0.3	0.3–0.7
Male	%	99.6	0.0	0.1	0.1	0.1	0.0	0.0	0.4
(n=2,601)	CI	99.2–99.8	0.0–0.3	0.0–0.4	0.0–0.4	0.0–0.4	0.0–0.2	0.0–0.3	0.2–0.8
Female	%	99.7	0.2	0.0	0.0	0.0	0.0	0.0	0.3
(n=2,467)	CI	99.4–99.9	0.1–0.5	0.0–0.3	–	0.0–0.2	–	–	0.1–0.6
White	%	99.6	0.2	0.1	0.0	0.0	0.0	0.1	0.4
(n=3,612)	CI	99.3–99.8	0.1–0.4	0.0–0.3	0.0–0.3	–	–	0.0–0.2	0.2–0.7
Black	%	99.7	0.0	0.0	0.0	0.3	0.0	0.0	0.3
(n=495)	CI	97.9–100.0	–	–	–	0.0–2.1	–	–	0.0–2.1
Hispanic	%	99.7	0.0	0.1	0.0	0.1	0.0	0.0	0.3
(n=469)	CI	98.8–99.9	–	0.0–1.1	–	0.0–1.1	–	–	0.1–1.2
8th	%	99.7	0.1	0.1	0.0	0.1	0.0	0.0	0.3
(n=2,555)	CI	99.4–99.9	0.0–0.3	0.0–0.3	–	0.0–0.3	0.0–0.3	–	0.1–0.6
10th	%	99.4	0.3	0.1	0.1	0.0	0.0	0.2	0.6
(n=1,571)	CI	98.8–99.7	0.1–0.8	0.0–0.5	0.0–0.5	–	–	0.1–0.6	0.3–1.2
12th	%	99.4	0.2	0.1	0.1	0.1	0.0	0.1	0.6
(n=1,069)	CI	98.8–99.7	0.0–0.7	0.0–0.7	0.0–0.7	0.0–0.7	–	0.0–0.7	0.3–1.2

Note: The confidence interval (CI) is 95 percent.

Table BH3.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.6	0.2	0.1	0.1	0.0	0.1	0.1	0.4
(n=5,195)	CI	99.3–99.7	0.1–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.0–0.2	0.0–0.2	0.3–0.7
Male	%	99.6	0.1	0.1	0.1	0.0	0.1	0.0	0.4
(n=2,601)	CI	99.3–99.8	0.0–0.3	0.0–0.4	0.0–0.3	0.0–0.3	0.0–0.4	–	0.2–0.7
Female	%	99.8	0.1	0.0	0.0	0.0	0.0	0.0	0.2
(n=2,467)	CI	99.6–99.9	0.0–0.2	0.0–0.3	–	0.0–0.3	0.0–0.2	–	0.1–0.4
White	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=3,612)	CI	99.5–99.9	0.1–0.3	0.0–0.3	–	0.0–0.2	–	0.0–0.2	0.1–0.5
Black	%	99.7	0.0	0.0	0.0	0.0	0.3	0.0	0.3
(n=495)	CI	97.9–100.0	–	–	–	–	0.0–2.1	–	0.0–2.1
Hispanic	%	99.9	0.0	0.0	0.1	0.0	0.0	0.0	0.1
(n=469)	CI	98.9–100.0	–	–	0.0–1.1	–	–	–	0.0–1.1
8th	%	99.8	0.1	0.0	0.0	0.0	0.1	0.0	0.2
(n=2,555)	CI	99.5–99.9	0.0–0.3	–	0.0–0.3	–	0.0–0.3	–	0.1–0.5
10th	%	99.4	0.2	0.1	0.1	0.1	0.0	0.1	0.6
(n=1,571)	CI	98.9–99.7	0.1–0.6	0.0–0.5	0.0–0.5	0.0–0.5	–	0.0–0.5	0.3–1.1
12th	%	99.5	0.2	0.1	0.0	0.0	0.1	0.1	0.5
(n=1,069)	CI	98.9–99.8	0.0–0.7	0.0–0.7	–	–	0.0–0.7	0.0–0.7	0.2–1.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you <u>use ecstasy (MDMA)</u> ?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.2	0.4	0.2	0.1	0.0	0.0	0.1	0.8
(n=5,196)	CI	98.9-99.5	0.2-0.6	0.1-0.4	0.0-0.2	0.0-0.2	0.0-0.1	0.1-0.3	0.5-1.1
Male	%	99.0	0.5	0.2	0.1	0.1	0.0	0.1	1.0
(n=2,602)	CI	98.5-99.4	0.3-1.0	0.1-0.6	0.0-0.4	0.0-0.4	-	0.0-0.4	0.6-1.5
Female	%	99.7	0.1	0.1	0.0	0.0	0.0	0.0	0.3
(n=2,467)	CI	99.4-99.9	0.0-0.4	0.0-0.4	0.0-0.3	-	0.0-0.2	-	0.1-0.6
White	%	99.5	0.4	0.0	0.1	0.0	0.0	0.1	0.5
(n=3,612)	CI	99.1-99.7	0.2-0.7	0.0-0.1	0.0-0.3	-	-	0.0-0.2	0.3-0.9
Black	%	99.7	0.0	0.0	0.0	0.0	0.0	0.3	0.3
(n=495)	CI	97.9-100.0	-	-	-	-	-	0.0-2.1	0.0-2.1
Hispanic	%	99.0	0.7	0.3	0.0	0.0	0.0	0.0	1.0
(n=470)	CI	97.3-99.7	0.2-2.6	0.1-1.2	-	-	-	-	0.3-2.7
8th	%	99.8	0.0	0.1	0.0	0.0	0.0	0.0	0.2
(n=2,555)	CI	99.5-99.9	0.0-0.3	0.0-0.4	-	-	0.0-0.3	-	0.1-0.5
10th	%	99.3	0.3	0.1	0.1	0.0	0.0	0.2	0.7
(n=1,571)	CI	98.7-99.6	0.1-0.8	0.0-0.5	0.0-0.5	-	-	0.1-0.6	0.4-1.3
12th	%	98.5	0.8	0.3	0.1	0.1	0.0	0.2	1.5
(n=1,070)	CI	97.6-99.1	0.4-1.6	0.1-0.9	0.0-0.7	0.0-0.7	-	0.0-0.7	0.9-2.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you <u>use methamphetamines (crank, etc.)</u> ?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.4	0.2	0.1	0.0	0.1	0.0	0.2	0.6
(n=5,193)	CI	99.1-99.6	0.1-0.4	0.0-0.2	0.0-0.1	0.0-0.3	-	0.1-0.4	0.4-0.9
Male	%	99.4	0.2	0.0	0.0	0.1	0.0	0.1	0.6
(n=2,601)	CI	99.0-99.7	0.1-0.6	0.0-0.3	0.0-0.3	0.0-0.5	-	0.0-0.5	0.3-1.0
Female	%	99.7	0.1	0.0	0.0	0.1	0.0	0.2	0.3
(n=2,466)	CI	99.3-99.8	0.0-0.4	0.0-0.2	-	0.0-0.2	-	0.0-0.5	0.2-0.7
White	%	99.6	0.1	0.0	0.0	0.1	0.0	0.1	0.4
(n=3,609)	CI	99.3-99.8	0.1-0.4	0.0-0.1	-	0.0-0.3	-	0.1-0.4	0.2-0.7
Black	%	99.5	0.2	0.0	0.0	0.0	0.0	0.3	0.5
(n=495)	CI	97.9-99.9	0.0-1.5	-	-	-	-	0.0-2.1	0.1-2.1
Hispanic	%	99.9	0.0	0.0	0.1	0.0	0.0	0.0	0.1
(n=470)	CI	98.9-100.0	-	-	0.0-1.1	-	-	-	0.0-1.1
8th	%	99.8	0.0	0.0	0.0	0.1	0.0	0.0	0.2
(n=2,552)	CI	99.5-99.9	-	0.0-0.3	0.0-0.3	0.0-0.4	-	0.0-0.3	0.1-0.5
10th	%	99.4	0.2	0.1	0.1	0.0	0.0	0.2	0.6
(n=1,571)	CI	98.9-99.7	0.1-0.6	0.0-0.5	0.0-0.5	-	-	0.1-0.6	0.3-1.1
12th	%	99.0	0.5	0.0	0.0	0.2	0.0	0.4	1.0
(n=1,070)	CI	98.2-99.4	0.2-1.1	-	-	0.0-0.7	-	0.1-1.0	0.6-1.8

Note: The confidence interval (CI) is 95 percent.

Table BH3.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	29.9	24.7	19.5	25.9	45.4
(n=1,564)	CI	27.5–32.4	22.5–27.1	17.5–21.6	23.7–28.2	42.8–48.0
Male	%	28.9	21.5	20.0	29.6	49.6
(n=703)	CI	25.5–32.7	18.5–24.9	17.0–23.3	26.2–33.2	45.7–53.4
Female	%	29.0	27.9	19.4	23.8	43.1
(n=794)	CI	25.7–32.4	24.7–31.3	16.6–22.5	20.9–27.0	39.5–46.8
White	%	27.5	24.0	20.1	28.4	48.5
(n=1,160)	CI	24.9–30.4	21.5–26.7	17.7–22.6	25.8–31.2	45.5–51.5
Black	%	40.4	25.9	20.4	13.3	33.7
(n=111)	CI	31.0–50.5	18.1–35.7	13.6–29.5	7.9–21.3	25.0–43.5
Hispanic	%	30.4	23.3	23.2	23.1	46.3
(n=104)	CI	21.7–40.7	15.8–33.1	15.7–32.8	15.7–32.5	36.4–56.5
8th	%	23.9	23.6	16.7	35.8	52.5
(n=728)	CI	20.9–27.1	20.6–26.8	14.2–19.6	32.4–39.4	48.9–56.2
10th	%	32.2	25.9	19.8	22.1	42.0
(n=478)	CI	28.1–36.5	22.1–30.0	16.5–23.7	18.6–26.1	37.6–46.5
12th	%	33.2	24.7	21.6	20.5	42.1
(n=358)	CI	28.6–38.3	20.5–29.4	17.6–26.2	16.6–25.0	37.1–47.3

Note: The confidence interval (CI) is 95 percent.

Table BH3.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	26.5	37.5	19.1	16.9	36.0
(n=2,443)	CI	24.6–28.4	35.5–39.6	17.5–20.8	15.5–18.5	34.0–38.0
Male	%	27.7	34.9	20.8	16.6	37.4
(n=1,132)	CI	25.0–30.6	32.1–37.9	18.4–23.4	14.5–18.9	34.5–40.4
Female	%	24.4	39.7	18.5	17.3	35.9
(n=1,228)	CI	21.9–27.0	36.9–42.7	16.4–20.9	15.3–19.6	33.1–38.7
White	%	25.0	38.2	19.4	17.4	36.8
(n=1,824)	CI	23.0–27.2	35.9–40.6	17.6–21.3	15.7–19.2	34.5–39.1
Black	%	33.3	32.6	14.7	19.4	34.1
(n=165)	CI	26.0–41.5	25.5–40.7	9.9–21.2	13.9–26.5	26.9–42.1
Hispanic	%	29.6	35.1	18.8	16.5	35.3
(n=161)	CI	22.5–37.8	27.7–43.4	13.3–25.8	11.5–23.1	28.1–43.3
8th	%	20.5	31.9	21.2	26.3	47.5
(n=1,036)	CI	18.2–23.1	29.1–34.8	18.8–23.8	23.7–29.1	44.5–50.6
10th	%	23.2	41.9	20.6	14.3	34.9
(n=809)	CI	20.4–26.2	38.5–45.3	18.0–23.5	12.1–16.9	31.7–38.3
12th	%	34.2	37.7	15.9	12.1	28.1
(n=598)	CI	30.5–38.1	33.9–41.7	13.2–19.1	9.7–15.0	24.6–31.8

Note: The confidence interval (CI) is 95 percent.

Table BH3.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	43.5	30.4	12.6	13.6	26.2
(n=2,340)	CI	41.3–45.6	28.5–32.3	11.3–14.0	12.3–15.0	24.4–28.0
Male	%	46.7	27.6	12.1	13.6	25.7
(n=1,138)	CI	43.6–49.7	25.0–30.4	10.3–14.1	11.8–15.7	23.2–28.3
Female	%	39.0	33.7	13.3	13.9	27.2
(n=1,126)	CI	36.1–42.1	30.9–36.7	11.4–15.5	12.0–16.0	24.6–29.9
White	%	43.7	30.2	12.3	13.8	26.2
(n=1,796)	CI	41.3–46.1	28.0–32.4	10.9–13.9	12.3–15.5	24.2–28.3
Black	%	49.1	25.8	13.6	11.5	25.1
(n=138)	CI	40.4–57.9	18.9–34.2	8.7–20.6	7.2–17.9	18.4–33.1
Hispanic	%	37.9	32.7	14.1	15.4	29.5
(n=131)	CI	29.3–47.2	24.8–41.7	9.0–21.4	10.2–22.4	22.2–38.0
8th	%	25.6	30.7	17.9	25.8	43.7
(n=930)	CI	22.9–28.5	27.8–33.8	15.6–20.5	23.1–28.7	40.6–46.9
10th	%	39.2	36.6	12.4	11.8	24.3
(n=803)	CI	35.9–42.6	33.3–40.0	10.3–14.9	9.8–14.2	21.4–27.3
12th	%	59.8	24.2	9.0	7.0	16.0
(n=607)	CI	55.9–63.6	20.9–27.7	7.0–11.5	5.2–9.3	13.3–19.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	33.8	28.5	14.8	22.9	37.7
(n=1,865)	CI	31.5–36.1	26.4–30.7	13.2–16.6	21.0–24.8	35.5–40.0
Male	%	34.7	25.7	15.7	23.8	39.5
(n=882)	CI	31.5–38.2	22.8–28.9	13.3–18.4	21.2–26.8	36.2–42.9
Female	%	31.7	30.7	14.5	23.1	37.6
(n=917)	CI	28.6–35.1	27.6–33.9	12.3–17.0	20.5–25.9	34.4–40.9
White	%	31.1	28.2	14.8	26.0	40.8
(n=1,346)	CI	28.5–33.8	25.7–30.8	12.9–16.9	23.7–28.4	38.1–43.5
Black	%	48.2	26.5	14.5	10.9	25.4
(n=166)	CI	40.2–56.2	19.9–34.2	9.9–20.7	7.0–16.6	19.3–32.6
Hispanic	%	33.4	29.6	20.1	16.8	37.0
(n=127)	CI	25.2–42.8	22.0–38.6	13.8–28.4	11.2–24.4	28.7–46.0
8th	%	17.5	20.8	15.6	46.1	61.7
(n=678)	CI	14.8–20.6	17.9–24.0	13.1–18.5	42.4–49.9	58.0–65.3
10th	%	34.7	31.9	14.8	18.6	33.4
(n=670)	CI	31.2–38.4	28.5–35.5	12.3–17.7	15.9–21.8	29.9–37.1
12th	%	42.4	29.9	14.4	13.2	27.6
(n=517)	CI	38.2–46.7	26.2–34.0	11.6–17.7	10.6–16.4	24.0–31.7

Note: The confidence interval (CI) is 95 percent.

Table BH3.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	2.0	98.0
(n=5,187)	CI	1.6–2.4	97.6–98.4
Male	%	2.2	97.8
(n=2,605)	CI	1.6–2.9	97.1–98.4
Female	%	1.6	98.4
(n=2,458)	CI	1.1–2.2	97.8–98.9
White	%	1.5	98.5
(n=3,604)	CI	1.2–2.0	98.0–98.8
Black	%	2.3	97.7
(n=492)	CI	1.2–4.3	95.7–98.8
Hispanic	%	1.9	98.1
(n=470)	CI	0.9–3.7	96.3–99.1
8th	%	1.6	98.4
(n=2,555)	CI	1.2–2.2	97.8–98.8
10th	%	1.9	98.1
(n=1,566)	CI	1.3–2.7	97.3–98.7
12th	%	2.4	97.6
(n=1,066)	CI	1.7–3.5	96.5–98.3

Note: The confidence interval (CI) is 95 percent.

Table BH3.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	12.7	87.3
(n=5,168)	CI	11.7–13.7	86.3–88.3
Male	%	11.7	88.3
(n=2,589)	CI	10.4–13.1	86.9–89.6
Female	%	13.4	86.6
(n=2,451)	CI	12.0–15.0	85.0–88.0
White	%	12.8	87.2
(n=3,588)	CI	11.7–14.1	85.9–88.3
Black	%	8.9	91.1
(n=495)	CI	6.5–12.1	87.9–93.5
Hispanic	%	11.3	88.7
(n=470)	CI	8.5–14.8	85.2–91.5
8th	%	7.3	92.7
(n=2,546)	CI	6.4–8.4	91.6–93.6
10th	%	12.6	87.4
(n=1,562)	CI	11.0–14.3	85.7–89.0
12th	%	18.5	81.5
(n=1,060)	CI	16.3–20.9	79.1–83.7

Note: The confidence interval (CI) is 95 percent.

Table BH3.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=97)	Male (n=49)	Female (n=40)	White (n=51)	Black (n=10)	Hispanic (n=8)	8th (n=41)	10th (n=30)	12th (n=26)
Got a prescription from one doctor	%	13.2	15.2	11.3	10.9	6.0	15.8	16.7	16.7	7.7
	CI	7.7–21.7	7.4–28.7	4.3–26.6	4.4–24.3	0.5–44.2	2.5–57.7	7.9–31.8	6.7–35.7	1.7–28.1
Got a prescription from more than one doctor	%	7.6	4.5	12.0	7.0	0.0	7.9	19.0	6.7	0.0
	CI	4.0–14.0	1.3–14.0	5.1–25.6	2.8–16.8	–	0.6–53.9	9.5–34.4	1.5–24.7	–
Bought on the Internet	%	11.0	16.2	5.8	6.8	26.6	17.5	4.8	6.7	19.2
	CI	5.7–20.4	7.6–31.3	1.2–23.2	2.0–20.6	5.1–70.8	1.5–74.3	1.1–18.0	1.5–24.7	7.7–40.4
Bought from a friend, relative, or someone you know	%	32.9	32.5	37.1	36.7	6.0	49.1	28.6	30.0	38.5
	CI	23.7–43.6	19.8–48.3	22.6–54.3	23.6–52.0	0.5–44.2	14.6–84.4	16.6–44.6	15.8–49.5	21.1–59.3
Got from a friend, relative, or someone you know for free	%	46.1	45.4	48.7	53.5	13.3	71.5	45.2	46.7	46.2
	CI	35.8–56.7	31.0–60.5	32.6–65.2	38.8–67.7	1.2–65.7	29.3–93.8	30.5–60.9	29.0–65.2	27.3–66.2
Took from a friend, relative, or someone you know without asking	%	26.3	27.9	21.4	25.2	22.9	20.6	33.3	33.3	15.4
	CI	18.3–36.3	16.5–43.1	11.2–37.1	14.8–39.6	4.2–67.1	3.3–66.5	20.4–49.4	18.3–52.8	5.5–36.3
Took from someone's house or workplace without asking	%	12.3	11.2	13.1	6.1	0.0	15.8	19.0	16.7	3.8
	CI	7.2–20.4	4.8–24.1	5.6–27.7	1.9–17.8	–	2.5–57.7	9.5–34.4	6.7–35.7	0.5–25.4
Got it at a party	%	27.2	23.9	35.0	24.3	22.9	25.4	19.0	30.0	30.8
	CI	18.7–37.7	13.4–39.1	20.5–52.8	13.8–39.1	4.2–67.1	4.0–73.4	9.5–34.4	15.8–49.5	15.4–52.0
Got it some other way	%	35.0	37.7	29.8	21.7	55.5	58.8	23.8	26.7	50.0
	CI	25.4–45.8	24.2–53.3	16.6–47.7	11.6–37.0	20.6–85.7	19.9–89.1	13.0–39.6	13.3–46.2	30.5–69.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=578)	Male (n=262)	Female (n=295)	White (n=403)	Black (n=39)	Hispanic (n=47)	8th (n=186)	10th (n=197)	12th (n=195)
I bought it in a store	%	5.1	7.9	1.7	3.3	10.7	7.2	1.6	3.6	7.6
	CI	3.4–7.4	4.9–12.4	0.7–4.1	1.8–5.9	3.7–27.0	2.1–21.7	0.5–4.9	1.7–7.3	4.6–12.3
I bought it in a gas station	%	6.2	7.6	3.7	5.8	3.4	4.3	2.7	5.6	8.1
	CI	4.4–8.7	4.7–12.0	2.0–6.8	3.7–8.8	0.4–21.8	0.9–17.6	1.1–6.3	3.1–9.8	5.0–12.9
I bought it in a restaurant	%	1.4	1.8	0.3	1.1	0.0	2.9	0.0	2.0	1.5
	CI	0.7–2.9	0.7–4.7	0.0–2.4	0.4–2.8	–	0.4–19.2	–	0.8–5.3	0.5–4.7
I bought it in a bar or club	%	2.7	3.3	1.1	1.6	0.0	2.9	1.1	3.0	3.0
	CI	1.6–4.5	1.6–6.5	0.3–3.7	0.7–3.7	–	0.4–19.2	0.3–4.2	1.4–6.7	1.4–6.7
I bought it at a public event	%	2.6	3.3	1.2	0.6	4.9	4.3	2.7	2.0	3.0
	CI	1.6–4.4	1.6–6.6	0.4–3.4	0.2–2.0	1.0–20.1	0.9–17.6	1.1–6.3	0.8–5.3	1.4–6.7
I gave someone money to buy it for me	%	29.4	28.8	30.5	31.4	14.1	32.8	11.8	25.9	39.6
	CI	25.6–33.6	23.2–35.1	25.2–36.4	26.8–36.5	5.7–30.8	20.0–48.8	7.8–17.3	20.2–32.5	32.9–46.6
I got it from a sibling (under 21)	%	7.6	7.5	7.7	5.8	15.0	10.7	8.6	9.1	6.1
	CI	5.7–10.1	4.7–11.6	5.2–11.4	3.9–8.5	6.1–32.4	4.2–24.8	5.3–13.6	5.8–14.1	3.5–10.5
I got it from a friend (under 21)	%	23.5	19.2	27.1	23.5	22.9	24.5	24.6	23.9	22.8
	CI	20.1–27.3	14.7–24.7	22.1–32.7	19.4–28.1	11.6–40.2	13.9–39.5	18.9–31.3	18.4–30.4	17.5–29.3
I got it at a party	%	40.4	38.5	42.2	36.3	52.3	57.4	27.3	38.1	47.7
	CI	36.3–44.7	32.5–44.9	36.4–48.3	31.5–41.4	35.5–68.5	42.1–71.4	21.3–34.2	31.5–45.1	40.8–54.8
I took it from a store or family member	%	10.7	11.6	9.1	9.8	7.3	11.2	17.6	8.6	9.1
	CI	8.4–13.5	8.1–16.1	6.3–13.0	7.2–13.1	2.1–22.3	4.6–24.7	12.8–23.8	5.4–13.5	5.8–14.1
I took it home without permission	%	18.5	18.5	17.5	20.0	12.8	13.3	24.6	24.9	11.2
	CI	15.5–21.9	14.2–23.9	13.6–22.3	16.3–24.2	5.5–27.2	6.1–26.8	18.9–31.3	19.3–31.4	7.4–16.4
I got it from my parents with permission	%	19.4	20.6	18.1	20.8	10.7	14.1	24.6	22.8	14.7
	CI	16.3–22.9	16.0–26.2	14.1–23.0	17.0–25.1	3.7–27.0	6.4–28.3	18.9–31.3	17.5–29.3	10.4–20.4
I got it from an older sibling (over 21)	%	13.7	12.3	14.9	11.9	25.7	16.3	11.8	12.2	15.7
	CI	11.0–17.0	8.6–17.2	11.0–19.7	8.9–15.7	13.5–43.4	7.8–31.0	7.8–17.3	8.3–17.6	11.3–21.6
I got it from an older friend (over 21)	%	25.4	22.9	28.7	23.9	30.6	32.6	13.9	21.8	33.0
	CI	21.8–29.4	17.9–28.8	23.5–34.6	19.7–28.6	17.1–48.4	19.5–49.0	9.6–19.7	16.6–28.2	26.7–39.9
I got it from some other adult	%	15.8	12.4	18.7	14.8	12.2	12.0	12.3	19.3	14.7
	CI	12.9–19.2	8.8–17.2	14.4–23.9	11.5–18.8	4.8–27.7	5.0–25.9	8.3–17.9	14.3–25.5	10.4–20.4
I got it some other way	%	17.4	20.9	14.2	15.5	22.0	20.2	24.1	17.3	14.7
	CI	14.5–20.8	16.2–26.4	10.6–18.8	12.3–19.5	11.2–38.7	10.7–34.8	18.4–30.8	12.6–23.2	10.4–20.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?				
		I bought or tried to buy alcohol	Yes, asked for proof of age	No, not asked for proof of age
Total	%	3.4	22.6	77.4
(n=130)	CI	2.9–4.1	15.8–31.2	68.8–84.2
Male	%	4.2	23.8	76.2
(n=75)	CI	3.3–5.2	14.9–35.7	64.3–85.1
Female	%	2.1	21.4	78.6
(n=43)	CI	1.5–2.8	10.6–38.5	61.5–89.4
White	%	2.3	17.6	82.4
(n=66)	CI	1.8–3.0	9.8–29.6	70.4–90.2
Black	%	5.4	26.6	73.4
(n=15)	CI	3.4–8.5	8.8–57.5	42.5–91.2
Hispanic	%	4.5	23.3	76.7
(n=15)	CI	2.8–7.3	6.5–57.3	42.7–93.5
8th	%	2.2	17.0	83.0
(n=46)	CI	1.7–2.9	8.5–31.1	68.9–91.5
10th	%	3.1	15.0	85.0
(n=40)	CI	2.3–4.1	6.7–30.4	69.6–93.3
12th	%	5.1	29.8	70.2
(n=44)	CI	3.9–6.7	18.1–44.8	55.2–81.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?					
		I bought or tried to buy alcohol	No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	3.4	43.4	28.7	27.8
(n=130)	CI	2.9–4.1	34.9–52.4	21.3–37.6	20.4–36.7
Male	%	4.2	40.6	32.6	26.8
(n=75)	CI	3.3–5.2	29.5–52.7	22.3–44.9	17.5–38.8
Female	%	2.1	56.7	20.7	22.6
(n=43)	CI	1.5–2.8	40.5–71.6	10.5–36.8	11.6–39.3
White	%	2.3	34.4	31.7	33.9
(n=66)	CI	1.8–3.0	23.2–47.7	20.7–45.2	22.4–47.6
Black	%	5.4	64.1	20.8	15.1
(n=15)	CI	3.4–8.5	36.7–84.6	6.2–51.2	4.1–42.8
Hispanic	%	4.5	59.4	27.8	12.7
(n=15)	CI	2.8–7.3	31.8–82.2	9.6–58.3	2.5–45.4
8th	%	2.2	55.3	23.4	21.3
(n=46)	CI	1.7–2.9	40.5–69.2	13.2–38.1	11.6–35.8
10th	%	3.1	50.0	27.5	22.5
(n=40)	CI	2.3–4.1	34.3–65.7	15.5–43.9	11.8–38.7
12th	%	5.1	34.0	31.9	34.0
(n=44)	CI	3.9–6.7	21.6–49.1	19.9–47.0	21.6–49.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.39. Ever Ridden with a Prescription Drugs Impaired Driver

		Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		
		No	Yes	Not sure
Total	%	80.9	8.5	10.6
(n=4,807)	CI	79.7–82.0	7.7–9.4	9.7–11.5
Male	%	82.7	7.3	10.0
(n=2,385)	CI	81.0–84.3	6.3–8.5	8.8–11.3
Female	%	79.8	9.5	10.7
(n=2,309)	CI	78.1–81.5	8.3–10.8	9.5–12.1
White	%	81.3	8.4	10.3
(n=3,386)	CI	79.9–82.6	7.5–9.4	9.3–11.4
Black	%	79.6	6.5	13.8
(n=413)	CI	75.2–83.5	4.4–9.7	10.7–17.7
Hispanic	%	86.7	6.1	7.2
(n=436)	CI	82.9–89.7	4.1–9.2	5.0–10.2
8th	%	80.4	8.0	11.7
(n=2,374)	CI	78.7–81.9	6.9–9.1	10.4–13.0
10th	%	80.2	8.7	11.0
(n=1,451)	CI	78.1–82.2	7.4–10.3	9.5–12.7
12th	%	82.1	8.9	9.0
(n=982)	CI	79.5–84.3	7.3–10.9	7.4–11.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.40. Frequency Riding with a Prescription Drugs Impaired Driver

		During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?					
		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	94.9	1.8	1.4	0.5	1.3	5.1
(n=5,034)	CI	94.3–95.5	1.5–2.2	1.1–1.8	0.3–0.8	1.0–1.7	4.5–5.7
Male	%	95.6	1.3	1.3	0.4	1.3	4.4
(n=2,507)	CI	94.7–96.4	0.9–1.9	0.9–1.9	0.2–0.8	0.9–1.9	3.6–5.3
Female	%	94.7	2.2	1.7	0.6	0.8	5.3
(n=2,407)	CI	93.7–95.5	1.7–2.9	1.2–2.3	0.4–1.0	0.5–1.3	4.5–6.3
White	%	95.6	1.5	1.5	0.4	1.0	4.4
(n=3,521)	CI	94.8–96.2	1.2–2.0	1.1–2.0	0.2–0.7	0.7–1.4	3.8–5.2
Black	%	94.4	1.7	1.1	0.6	2.2	5.6
(n=466)	CI	91.6–96.3	0.8–3.7	0.4–2.6	0.2–1.9	1.1–4.4	3.7–8.4
Hispanic	%	96.2	2.2	0.8	0.8	0.0	3.8
(n=454)	CI	93.7–97.7	1.1–4.2	0.3–2.7	0.3–2.3	–	2.3–6.3
8th	%	95.1	2.0	1.1	0.7	1.0	4.9
(n=2,480)	CI	94.2–95.9	1.5–2.6	0.8–1.6	0.5–1.1	0.7–1.5	4.1–5.8
10th	%	95.1	1.6	1.4	0.3	1.5	4.9
(n=1,520)	CI	93.9–96.1	1.1–2.4	0.9–2.1	0.1–0.8	1.0–2.3	3.9–6.1
12th	%	94.5	1.8	1.8	0.5	1.3	5.5
(n=1,034)	CI	93.0–95.8	1.2–2.8	1.2–2.8	0.2–1.2	0.8–2.3	4.2–7.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	69.1	20.3	10.6
(n=5,062)	CI	67.8–70.5	19.1–21.5	9.7–11.5
Male	%	72.1	16.9	11.0
(n=2,519)	CI	70.2–74.0	15.3–18.5	9.8–12.3
Female	%	66.7	23.6	9.7
(n=2,420)	CI	64.7–68.6	21.8–25.4	8.6–11.0
White	%	69.9	20.1	10.0
(n=3,531)	CI	68.2–71.4	18.8–21.6	9.0–11.1
Black	%	64.4	21.4	14.2
(n=474)	CI	59.7–68.9	17.7–25.6	11.2–17.8
Hispanic	%	75.6	14.6	9.8
(n=460)	CI	71.1–79.6	11.4–18.4	7.3–13.1
8th	%	71.1	16.6	12.3
(n=2,502)	CI	69.3–72.8	15.2–18.1	11.1–13.7
10th	%	69.2	20.8	10.0
(n=1,525)	CI	66.9–71.5	18.9–22.9	8.5–11.6
12th	%	67.0	23.7	9.3
(n=1,035)	CI	64.0–69.8	21.2–26.4	7.7–11.2

Note: The confidence interval (CI) is 95 percent.

Table BH3.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	88.9	4.3	3.5	1.2	2.2	11.1
(n=5,048)	CI	87.9–89.8	3.7–4.9	3.0–4.1	0.9–1.5	1.8–2.7	10.2–12.1
Male	%	90.9	3.2	2.8	0.8	2.2	9.1
(n=2,513)	CI	89.6–92.1	2.6–4.1	2.2–3.6	0.5–1.3	1.7–3.0	7.9–10.4
Female	%	87.1	5.4	4.3	1.6	1.7	12.9
(n=2,414)	CI	85.6–88.5	4.5–6.4	3.5–5.2	1.1–2.2	1.2–2.3	11.5–14.4
White	%	89.8	3.8	3.4	1.4	1.7	10.2
(n=3,523)	CI	88.6–90.8	3.2–4.5	2.8–4.1	1.0–1.9	1.3–2.2	9.2–11.4
Black	%	84.7	7.4	4.0	0.4	3.5	15.3
(n=472)	CI	80.9–87.9	5.2–10.3	2.5–6.4	0.1–1.5	2.1–6.0	12.1–19.1
Hispanic	%	92.2	2.9	3.9	0.3	0.7	7.8
(n=458)	CI	89.1–94.4	1.7–5.1	2.4–6.3	0.1–1.2	0.3–1.9	5.6–10.9
8th	%	90.9	3.7	3.0	0.9	1.5	9.1
(n=2,495)	CI	89.7–91.9	3.0–4.5	2.4–3.8	0.6–1.3	1.1–2.1	8.1–10.3
10th	%	89.3	4.3	3.4	1.2	1.7	10.7
(n=1,519)	CI	87.6–90.7	3.4–5.5	2.6–4.5	0.8–2.0	1.2–2.5	9.3–12.4
12th	%	86.3	4.9	4.0	1.3	3.4	13.7
(n=1,034)	CI	84.1–88.3	3.7–6.4	3.0–5.4	0.8–2.3	2.4–4.7	11.7–15.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.7	2.0	5.1	91.1	96.2
(n=5,041)	CI	1.4–2.2	1.6–2.5	4.5–5.8	90.2–91.9	95.6–96.8
Male	%	1.9	1.4	4.8	91.9	96.7
(n=2,508)	CI	1.4–2.6	1.0–2.0	4.0–5.8	90.7–93.0	95.9–97.4
Female	%	1.2	2.5	5.4	90.9	96.3
(n=2,409)	CI	0.8–1.7	1.9–3.3	4.5–6.4	89.6–92.0	95.4–97.0
White	%	1.1	1.9	5.0	92.0	97.0
(n=3,515)	CI	0.8–1.6	1.5–2.4	4.3–5.8	91.0–92.9	96.3–97.5
Black	%	3.5	1.9	5.3	89.3	94.6
(n=468)	CI	2.0–5.9	0.9–4.0	3.5–8.0	85.8–92.0	91.8–96.5
Hispanic	%	1.3	2.3	4.4	91.9	96.3
(n=460)	CI	0.5–3.2	1.3–4.3	2.7–7.0	88.8–94.3	94.0–97.8
8th	%	1.2	1.9	4.8	92.1	96.9
(n=2,492)	CI	0.9–1.8	1.4–2.5	4.0–5.7	90.9–93.1	96.1–97.5
10th	%	1.7	2.2	5.1	91.0	96.1
(n=1,520)	CI	1.2–2.5	1.6–3.1	4.1–6.3	89.5–92.3	95.0–96.9
12th	%	2.3	1.9	5.5	90.2	95.7
(n=1,029)	CI	1.6–3.4	1.2–3.0	4.3–7.1	88.3–91.9	94.3–96.8

Note: The confidence interval (CI) is 95 percent.

Table BH3.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.2	4.5	10.7	82.6	93.3
(n=5,062)	CI	1.8–2.7	3.9–5.1	9.8–11.7	81.4–83.7	92.5–94.0
Male	%	2.7	4.3	11.6	81.4	93.0
(n=2,518)	CI	2.1–3.5	3.6–5.3	10.3–13.0	79.7–83.0	91.8–94.0
Female	%	1.2	4.6	9.8	84.4	94.3
(n=2,419)	CI	0.8–1.8	3.8–5.6	8.6–11.2	82.8–85.9	93.1–95.2
White	%	1.9	4.9	11.5	81.7	93.2
(n=3,533)	CI	1.5–2.5	4.2–5.7	10.4–12.7	80.3–83.0	92.2–94.0
Black	%	3.3	2.3	7.9	86.5	94.4
(n=469)	CI	1.9–5.7	1.2–4.4	5.6–10.9	82.8–89.5	91.6–96.3
Hispanic	%	1.8	2.5	9.2	86.6	95.7
(n=461)	CI	0.8–3.9	1.3–4.7	6.6–12.6	82.6–89.7	93.1–97.4
8th	%	0.8	3.7	8.5	87.0	95.6
(n=2,498)	CI	0.5–1.2	3.0–4.5	7.5–9.7	85.7–88.3	94.7–96.3
10th	%	2.2	4.2	11.0	82.7	93.7
(n=1,528)	CI	1.5–3.0	3.3–5.3	9.5–12.6	80.7–84.5	92.3–94.8
12th	%	3.8	5.7	12.8	77.7	90.5
(n=1,036)	CI	2.8–5.2	4.4–7.2	10.9–14.9	75.1–80.2	88.6–92.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.45. Parents Feel Regularly Drinking Alcohol is Wrong

How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.2	8.7	11.5	74.6	86.1
(n=5,053)	CI	4.6–5.9	7.9–9.6	10.6–12.4	73.3–75.9	85.0–87.1
Male	%	5.4	8.1	11.7	74.7	86.5
(n=2,513)	CI	4.5–6.5	7.0–9.3	10.4–13.2	72.9–76.5	84.9–87.9
Female	%	4.6	9.3	11.1	75.1	86.2
(n=2,415)	CI	3.7–5.6	8.1–10.6	9.8–12.5	73.2–76.9	84.6–87.6
White	%	4.8	9.6	12.1	73.5	85.6
(n=3,527)	CI	4.1–5.7	8.6–10.7	11.0–13.3	71.9–75.0	84.3–86.8
Black	%	4.7	6.0	7.8	81.6	89.4
(n=468)	CI	2.9–7.4	4.0–8.8	5.6–10.8	77.4–85.1	85.8–92.1
Hispanic	%	3.8	5.6	11.8	78.8	90.6
(n=460)	CI	2.2–6.5	3.6–8.4	8.9–15.5	74.4–82.7	87.1–93.2
8th	%	2.9	6.4	9.9	80.8	90.7
(n=2,495)	CI	2.3–3.7	5.5–7.4	8.7–11.1	79.2–82.3	89.5–91.7
10th	%	5.0	8.4	11.4	75.1	86.6
(n=1,523)	CI	4.1–6.3	7.1–9.9	9.9–13.1	72.9–77.3	84.7–88.2
12th	%	7.9	11.4	13.3	67.4	80.7
(n=1,035)	CI	6.4–9.7	9.6–13.5	11.3–15.5	64.5–70.2	78.2–83.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.46. Parents Feel Smoking Tobacco is Wrong

How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.8	3.8	9.3	84.2	93.4
(n=5,055)	CI	2.3–3.3	3.3–4.4	8.4–10.2	83.1–85.3	92.6–94.2
Male	%	3.2	3.6	9.2	84.0	93.2
(n=2,514)	CI	2.5–4.1	2.8–4.5	8.0–10.5	82.4–85.5	92.0–94.3
Female	%	1.8	4.1	9.2	85.0	94.1
(n=2,416)	CI	1.3–2.5	3.3–5.1	8.0–10.5	83.3–86.5	93.0–95.1
White	%	2.6	4.4	10.3	82.7	93.0
(n=3,526)	CI	2.1–3.3	3.7–5.3	9.2–11.4	81.3–84.0	91.9–93.9
Black	%	2.3	3.6	7.7	86.5	94.1
(n=470)	CI	1.2–4.5	2.1–6.0	5.4–10.7	82.7–89.5	91.2–96.1
Hispanic	%	1.8	1.5	4.7	92.0	96.7
(n=461)	CI	0.8–3.9	0.6–3.4	2.9–7.5	88.7–94.4	94.3–98.1
8th	%	1.1	1.8	6.6	90.5	97.1
(n=2,498)	CI	0.7–1.6	1.4–2.4	5.7–7.6	89.3–91.6	96.3–97.7
10th	%	2.4	3.4	8.4	85.8	94.2
(n=1,524)	CI	1.8–3.3	2.6–4.4	7.1–9.9	83.9–87.4	92.9–95.2
12th	%	4.9	6.4	13.1	75.6	88.7
(n=1,033)	CI	3.7–6.4	5.0–8.0	11.2–15.3	72.9–78.2	86.7–90.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.8	1.7	5.6	90.9	96.5
(n=5,050)	CI	1.5–2.3	1.3–2.1	4.9–6.3	90.0–91.8	95.9–97.0
Male	%	2.2	1.6	5.2	91.0	96.2
(n=2,514)	CI	1.6–2.9	1.1–2.3	4.3–6.3	89.7–92.2	95.3–97.0
Female	%	0.9	1.8	5.9	91.3	97.3
(n=2,412)	CI	0.6–1.5	1.3–2.5	5.0–7.1	90.0–92.5	96.4–97.9
White	%	1.3	1.8	6.0	90.9	96.9
(n=3,525)	CI	1.0–1.8	1.4–2.4	5.2–7.0	89.7–91.9	96.2–97.5
Black	%	2.8	2.2	5.0	90.0	94.9
(n=469)	CI	1.5–5.2	1.1–4.4	3.2–7.6	86.5–92.6	92.1–96.8
Hispanic	%	1.5	0.5	3.8	94.3	98.0
(n=461)	CI	0.6–3.4	0.1–2.1	2.2–6.3	91.3–96.2	96.0–99.1
8th	%	0.9	0.8	3.0	95.4	98.4
(n=2,497)	CI	0.6–1.3	0.5–1.2	2.4–3.7	94.5–96.1	97.8–98.8
10th	%	2.2	1.0	5.2	91.6	96.8
(n=1,523)	CI	1.5–3.0	0.6–1.7	4.2–6.4	90.1–92.9	95.8–97.6
12th	%	2.5	3.4	8.8	85.3	94.1
(n=1,030)	CI	1.7–3.7	2.4–4.7	7.2–10.7	83.0–87.4	92.5–95.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.48. Parents Feel Using Electronic Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.9	7.6	12.1	75.3	87.5
(n=5,046)	CI	4.3–5.7	6.8–8.4	11.2–13.1	74.0–76.6	86.4–88.4
Male	%	5.4	6.9	12.0	75.7	87.7
(n=2,508)	CI	4.5–6.5	5.9–8.1	10.7–13.4	73.8–77.4	86.2–89.0
Female	%	4.0	8.2	12.0	75.8	87.8
(n=2,413)	CI	3.2–5.0	7.1–9.5	10.7–13.5	73.9–77.6	86.3–89.1
White	%	4.9	8.2	13.4	73.5	86.9
(n=3,522)	CI	4.2–5.8	7.2–9.2	12.3–14.7	71.9–75.1	85.7–88.1
Black	%	4.2	4.8	7.9	83.2	91.0
(n=469)	CI	2.5–6.7	3.1–7.3	5.7–10.8	79.2–86.5	87.8–93.5
Hispanic	%	2.3	4.0	8.4	85.4	93.7
(n=461)	CI	1.2–4.5	2.3–6.6	6.0–11.6	81.4–88.6	90.7–95.8
8th	%	2.6	6.1	10.1	81.2	91.3
(n=2,493)	CI	2.0–3.3	5.3–7.1	9.0–11.4	79.6–82.7	90.1–92.3
10th	%	4.9	6.8	12.9	75.4	88.3
(n=1,521)	CI	3.9–6.1	5.7–8.2	11.3–14.7	73.1–77.5	86.6–89.8
12th	%	7.6	10.0	13.4	69.0	82.4
(n=1,032)	CI	6.1–9.4	8.3–12.0	11.4–15.6	66.1–71.7	79.9–84.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.3	5.0	7.9	81.8	89.8
(n=5,049)	CI	4.6–6.0	4.3–5.7	7.2–8.8	80.6–83.0	88.8–90.6
Male	%	5.5	4.3	7.9	82.3	90.2
(n=2,514)	CI	4.6–6.6	3.5–5.3	6.8–9.1	80.6–83.9	88.9–91.4
Female	%	4.6	5.5	7.9	81.9	89.8
(n=2,411)	CI	3.8–5.7	4.6–6.6	6.8–9.2	80.2–83.5	88.4–91.1
White	%	4.4	4.8	8.2	82.6	90.8
(n=3,525)	CI	3.8–5.3	4.1–5.6	7.3–9.2	81.2–83.9	89.7–91.8
Black	%	7.6	5.9	8.8	77.7	86.5
(n=466)	CI	5.3–10.9	3.9–8.8	6.3–12.0	73.3–81.6	82.6–89.6
Hispanic	%	3.7	2.5	5.5	88.3	93.8
(n=461)	CI	2.1–6.3	1.3–4.8	3.6–8.5	84.5–91.2	90.7–95.9
8th	%	2.4	2.8	5.6	89.2	94.8
(n=2,497)	CI	1.9–3.1	2.2–3.5	4.8–6.6	87.9–90.3	93.9–95.6
10th	%	5.4	5.6	8.2	80.8	89.0
(n=1,519)	CI	4.4–6.6	4.5–6.9	6.9–9.7	78.7–82.7	87.3–90.5
12th	%	8.3	6.6	10.1	75.0	85.1
(n=1,033)	CI	6.7–10.1	5.3–8.3	8.4–12.1	72.2–77.5	82.8–87.1

Note: The confidence interval (CI) is 95 percent.

Table BH3.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.9	3.4	7.0	84.7	91.7
(n=5,043)	CI	4.3–5.6	2.9–4.0	6.3–7.8	83.6–85.7	90.8–92.5
Male	%	4.9	3.1	7.1	84.9	92.0
(n=2,510)	CI	4.1–6.0	2.4–3.9	6.1–8.3	83.3–86.3	90.7–93.1
Female	%	4.4	3.7	6.8	85.2	92.0
(n=2,409)	CI	3.5–5.4	2.9–4.6	5.8–7.9	83.6–86.7	90.7–93.1
White	%	4.1	3.3	7.1	85.6	92.7
(n=3,520)	CI	3.4–4.8	2.7–4.0	6.2–8.0	84.3–86.8	91.6–93.6
Black	%	7.5	4.2	8.2	80.1	88.3
(n=469)	CI	5.2–10.8	2.5–6.7	5.9–11.4	75.8–83.8	84.6–91.2
Hispanic	%	3.2	2.3	4.4	90.0	94.4
(n=460)	CI	1.8–5.7	1.2–4.7	2.8–7.0	86.5–92.7	91.4–96.4
8th	%	2.2	1.9	4.8	91.1	95.9
(n=2,495)	CI	1.7–2.8	1.5–2.5	4.0–5.7	89.9–92.2	95.1–96.6
10th	%	5.1	3.3	7.9	83.7	91.7
(n=1,521)	CI	4.1–6.3	2.5–4.3	6.7–9.4	81.8–85.5	90.2–93.0
12th	%	7.7	5.2	8.3	78.7	87.0
(n=1,027)	CI	6.3–9.5	4.0–6.8	6.8–10.2	76.1–81.1	84.8–88.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.51. Friends Feel Misusing Prescription Drugs is Wrong

How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	8.0	5.9	15.2	70.9	86.1
(n=5,028)	CI	7.2–8.8	5.3–6.7	14.2–16.3	69.5–72.2	85.0–87.1
Male	%	8.7	6.2	15.1	70.1	85.1
(n=2,495)	CI	7.5–10.0	5.3–7.3	13.6–16.6	68.1–72.0	83.6–86.6
Female	%	6.6	5.6	15.4	72.4	87.7
(n=2,409)	CI	5.6–7.8	4.7–6.7	13.9–16.9	70.4–74.2	86.3–89.1
White	%	7.2	5.8	15.8	71.2	87.0
(n=3,511)	CI	6.4–8.2	5.0–6.7	14.5–17.1	69.6–72.8	85.7–88.1
Black	%	8.2	4.6	12.8	74.3	87.1
(n=466)	CI	5.9–11.4	3.0–7.1	9.9–16.4	69.8–78.3	83.5–90.1
Hispanic	%	8.1	7.3	15.2	69.4	84.6
(n=458)	CI	5.7–11.3	5.1–10.4	12.0–19.1	64.7–73.7	80.6–87.9
8th	%	5.6	5.8	14.6	74.0	88.6
(n=2,482)	CI	4.8–6.6	4.9–6.7	13.3–16.1	72.2–75.7	87.3–89.8
10th	%	8.2	5.7	15.9	70.2	86.1
(n=1,518)	CI	6.9–9.6	4.7–7.0	14.2–17.8	67.8–72.4	84.3–87.8
12th	%	10.3	6.4	15.1	68.3	83.4
(n=1,028)	CI	8.5–12.3	5.0–8.0	13.0–17.4	65.4–71.0	81.0–85.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.52. Friends Feel Drinking Alcohol is Wrong

How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	15.0	14.9	18.5	51.6	70.1
(n=5,035)	CI	14.0–16.2	13.8–15.9	17.4–19.7	50.1–53.1	68.7–71.5
Male	%	16.4	14.3	18.2	51.1	69.4
(n=2,505)	CI	14.8–18.0	12.9–15.8	16.7–19.9	49.1–53.2	67.4–71.3
Female	%	13.0	15.6	18.9	52.4	71.4
(n=2,406)	CI	11.6–14.6	14.1–17.3	17.4–20.6	50.3–54.5	69.4–73.3
White	%	15.0	16.2	18.9	49.9	68.8
(n=3,516)	CI	13.8–16.4	14.9–17.6	17.6–20.3	48.1–51.6	67.1–70.4
Black	%	12.3	8.5	17.1	62.1	79.2
(n=466)	CI	9.4–16.0	6.1–11.6	13.7–21.0	57.3–66.7	74.9–82.9
Hispanic	%	13.6	13.1	18.9	54.4	73.3
(n=459)	CI	10.4–17.5	10.0–17.0	15.4–22.9	49.5–59.2	68.6–77.5
8th	%	8.8	11.3	18.9	61.1	80.0
(n=2,487)	CI	7.7–9.9	10.1–12.6	17.4–20.5	59.1–63.0	78.4–81.5
10th	%	14.9	15.5	21.3	48.3	69.6
(n=1,517)	CI	13.2–16.8	13.8–17.4	19.3–23.5	45.7–50.8	67.2–71.9
12th	%	22.0	18.0	15.0	44.9	60.0
(n=1,031)	CI	19.6–24.6	15.8–20.5	13.0–17.4	41.9–48.0	57.0–62.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	15.1	11.1	18.3	55.5	73.9
(n=5035)	CI	14.0–16.2	10.2–12.0	17.2–19.5	54.1–57.0	72.5–75.2
Male	%	16.6	10.5	18.5	54.3	72.9
(n=2,505)	CI	15.1–18.3	9.3–11.9	17.0–20.2	52.3–56.4	70.9–74.7
Female	%	13.0	11.9	18.0	57.1	75.2
(n=2,406)	CI	11.5–14.5	10.6–13.4	16.4–19.7	55.0–59.2	73.2–77.0
White	%	15.9	11.8	19.1	53.2	72.3
(n=3,518)	CI	14.6–17.3	10.7–13.0	17.8–20.5	51.4–54.9	70.7–73.9
Black	%	10.8	7.6	15.1	66.5	81.6
(n=467)	CI	8.1–14.2	5.4–10.6	12.0–18.9	61.8–70.9	77.5–85.1
Hispanic	%	10.3	9.6	20.8	59.3	80.1
(n=457)	CI	7.6–13.9	7.0–13.0	17.1–25.0	54.4–64.0	75.7–83.8
8th	%	7.8	9.2	17.2	65.7	83.0
(n=2,486)	CI	6.8–9.0	8.1–10.4	15.8–18.8	63.9–67.6	81.5–84.4
10th	%	14.2	11.0	20.4	54.3	74.8
(n=1,519)	CI	12.5–16.0	9.6–12.7	18.5–22.6	51.8–56.8	72.5–76.9
12th	%	23.8	13.1	17.2	45.8	63.0
(n=1,030)	CI	21.3–26.5	11.2–15.3	15.0–19.6	42.8–48.9	60.0–65.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	26.0	14.5	13.8	45.7	59.5
(n=5,019)	CI	24.7–27.3	13.5–15.6	12.8–14.8	44.3–47.2	58.1–61.0
Male	%	25.7	14.3	14.0	46.1	60.0
(n=2,494)	CI	23.8–27.6	12.9–15.8	12.6–15.4	44.0–48.1	58.0–62.1
Female	%	25.7	14.9	13.7	45.8	59.5
(n=2,404)	CI	23.8–27.6	13.4–16.4	12.3–15.2	43.7–47.9	57.4–61.5
White	%	27.4	15.8	13.9	42.8	56.7
(n=3,508)	CI	25.8–29.1	14.6–17.2	12.8–15.2	41.1–44.5	55.0–58.5
Black	%	16.8	8.7	13.8	60.7	74.5
(n=466)	CI	13.5–20.7	6.4–11.8	10.7–17.5	55.9–65.3	70.1–78.4
Hispanic	%	17.7	12.1	18.4	51.8	70.2
(n=454)	CI	14.2–21.9	9.2–15.8	14.9–22.4	46.9–56.7	65.4–74.6
8th	%	16.4	13.1	14.9	55.6	70.5
(n=2,476)	CI	15.0–17.9	11.8–14.5	13.6–16.4	53.6–57.5	68.7–72.3
10th	%	26.4	16.7	15.3	41.6	56.9
(n=1,516)	CI	24.2–28.6	14.9–18.7	13.6–17.3	39.1–44.1	54.4–59.4
12th	%	35.8	13.6	10.9	39.6	50.5
(n=1,027)	CI	32.9–38.8	11.7–15.9	9.2–13.0	36.6–42.6	47.5–53.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	21.1	11.7	14.7	52.4	67.1
(n=5,030)	CI	19.9–22.4	10.8–12.7	13.7–15.8	50.9–53.9	65.7–68.5
Male	%	20.5	11.0	15.4	53.1	68.5
(n=2,500)	CI	18.9–22.3	9.7–12.4	14.0–16.9	51.0–55.2	66.5–70.4
Female	%	21.1	12.5	14.1	52.3	66.4
(n=2,406)	CI	19.3–22.9	11.2–14.0	12.7–15.6	50.2–54.4	64.3–68.4
White	%	20.0	12.4	15.4	52.2	67.6
(n=3,516)	CI	18.6–21.5	11.3–13.6	14.2–16.7	50.5–54.0	65.9–69.3
Black	%	25.3	8.0	11.0	55.7	66.6
(n=467)	CI	21.3–29.9	5.8–11.0	8.3–14.4	50.8–60.4	61.9–71.1
Hispanic	%	16.3	11.3	17.3	55.1	72.5
(n=459)	CI	12.8–20.4	8.5–14.9	13.9–21.3	50.2–60.0	67.7–76.7
8th	%	12.1	8.5	13.7	65.7	79.4
(n=2,482)	CI	10.9–13.5	7.4–9.6	12.4–15.1	63.8–67.6	77.8–81.0
10th	%	22.5	13.0	16.3	48.3	64.5
(n=1,517)	CI	20.5–24.7	11.4–14.8	14.5–18.2	45.7–50.8	62.1–66.9
12th	%	29.4	14.0	14.2	42.4	56.6
(n=1,031)	CI	26.7–32.3	12.0–16.2	12.2–16.4	39.4–45.5	53.6–59.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	89.3	4.8	2.2	0.9	2.7	10.7
(n=5,039)	CI	88.3–90.2	4.2–5.5	1.8–2.7	0.7–1.3	2.3–3.3	9.8–11.7
Male	%	90.7	4.0	2.3	0.7	2.4	9.3
(n=2,509)	CI	89.3–91.9	3.2–4.9	1.7–3.0	0.4–1.1	1.8–3.1	8.1–10.7
Female	%	88.6	5.5	2.1	1.1	2.7	11.4
(n=2,409)	CI	87.1–89.8	4.6–6.5	1.6–2.8	0.8–1.7	2.1–3.5	10.2–12.9
White	%	89.4	5.0	2.3	0.9	2.4	10.6
(n=3,518)	CI	88.3–90.5	4.3–5.8	1.8–2.9	0.6–1.3	1.9–3.0	9.5–11.7
Black	%	91.3	3.4	0.9	1.6	2.8	8.7
(n=472)	CI	88.1–93.7	2.0–5.6	0.3–2.5	0.7–3.6	1.5–5.0	6.3–11.9
Hispanic	%	91.6	3.2	2.3	1.0	1.9	8.4
(n=458)	CI	88.3–94.0	1.7–5.7	1.2–4.3	0.3–2.8	1.0–3.9	6.0–11.7
8th	%	92.0	4.3	1.2	0.9	1.6	8.0
(n=2,484)	CI	90.9–93.0	3.5–5.1	0.9–1.8	0.6–1.3	1.2–2.2	7.0–9.1
10th	%	88.6	5.4	2.4	0.7	2.8	11.4
(n=1,518)	CI	86.9–90.1	4.4–6.7	1.8–3.3	0.4–1.3	2.1–3.8	9.9–13.1
12th	%	87.2	4.7	3.1	1.2	3.8	12.8
(n=1,037)	CI	85.0–89.1	3.6–6.2	2.2–4.3	0.7–2.1	2.8–5.2	10.9–15.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	56.8	12.7	10.7	5.0	14.7	43.2
(n=5,066)	CI	55.3–58.2	11.8–13.7	9.9–11.7	4.4–5.7	13.6–15.8	41.8–44.7
Male	%	61.7	10.2	9.6	4.2	14.3	38.3
(n=2,521)	CI	59.7–63.7	9.0–11.5	8.4–10.9	3.4–5.1	12.9–15.9	36.3–40.3
Female	%	51.9	15.6	12.1	5.8	14.6	48.1
(n=2,421)	CI	49.8–54.0	14.1–17.1	10.8–13.6	4.9–7.0	13.1–16.2	46.0–50.2
White	%	53.6	13.5	11.5	5.8	15.6	46.4
(n=3,538)	CI	51.9–55.3	12.3–14.7	10.5–12.7	5.0–6.7	14.3–17.0	44.7–48.1
Black	%	72.4	8.8	6.9	2.9	9.1	27.6
(n=476)	CI	67.9–76.5	6.5–11.9	4.9–9.7	1.6–5.1	6.6–12.3	23.5–32.1
Hispanic	%	66.1	12.1	7.8	3.1	10.8	33.9
(n=462)	CI	61.3–70.7	9.2–15.8	5.5–11.0	1.7–5.6	8.0–14.4	29.3–38.7
8th	%	70.2	11.6	8.6	3.0	6.6	29.8
(n=2,494)	CI	68.4–72.0	10.4–12.9	7.6–9.8	2.4–3.7	5.7–7.6	28.0–31.6
10th	%	52.1	14.9	11.9	5.4	15.8	47.9
(n=1,531)	CI	49.5–54.6	13.2–16.7	10.3–13.6	4.4–6.7	14.0–17.7	45.4–50.5
12th	%	47.4	11.7	11.8	6.9	22.3	52.6
(n=1,041)	CI	44.4–50.4	9.8–13.7	10.0–13.9	5.5–8.6	19.8–24.9	49.6–55.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	72.0	9.0	6.2	3.2	9.7	28.0
(n=5,056)	CI	70.6–73.3	8.2–9.8	5.5–6.9	2.7–3.8	8.8–10.6	26.7–29.4
Male	%	73.8	7.8	5.5	3.0	9.8	26.2
(n=2,517)	CI	71.9–75.6	6.8–9.0	4.6–6.6	2.4–3.9	8.6–11.2	24.4–28.1
Female	%	70.6	10.2	6.8	3.4	9.0	29.4
(n=2,417)	CI	68.6–72.5	9.0–11.6	5.8–8.0	2.7–4.3	7.8–10.4	27.5–31.4
White	%	69.3	9.7	6.7	3.8	10.5	30.7
(n=3,532)	CI	67.6–70.9	8.7–10.8	5.8–7.7	3.1–4.5	9.4–11.7	29.1–32.4
Black	%	86.4	3.7	3.4	1.3	5.2	13.6
(n=474)	CI	82.6–89.5	2.2–6.1	2.0–5.7	0.5–3.1	3.4–7.9	10.5–17.4
Hispanic	%	79.9	8.0	5.3	2.0	4.9	20.1
(n=459)	CI	75.4–83.7	5.6–11.3	3.4–8.1	1.0–4.1	3.0–7.9	16.3–24.6
8th	%	86.3	6.7	3.4	0.8	2.8	13.7
(n=2,489)	CI	84.9–87.6	5.8–7.8	2.7–4.2	0.5–1.2	2.2–3.5	12.4–15.1
10th	%	68.4	10.7	6.7	4.1	10.1	31.6
(n=1,525)	CI	66.1–70.7	9.3–12.4	5.5–8.0	3.2–5.2	8.7–11.7	29.3–33.9
12th	%	60.4	9.4	8.6	5.0	16.6	39.6
(n=1,042)	CI	57.4–63.3	7.8–11.4	7.0–10.4	3.8–6.5	14.5–19.0	36.7–42.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have smoked part, or all of a cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	72.6	9.8	5.3	3.1	9.2	27.4
(n=5,039)	CI	71.3–73.9	9.0–10.7	4.6–6.0	2.6–3.7	8.3–10.1	26.1–28.7
Male	%	75.5	8.6	4.6	2.6	8.7	24.5
(n=2,506)	CI	73.6–77.3	7.5–9.9	3.7–5.5	2.0–3.4	7.5–10.0	22.7–26.4
Female	%	70.1	11.2	5.9	3.7	9.2	29.9
(n=2,410)	CI	68.1–72.0	9.9–12.6	4.9–7.0	2.9–4.6	8.0–10.6	28.0–31.9
White	%	69.9	11.1	5.8	3.6	9.7	30.1
(n=3,514)	CI	68.2–71.5	10.0–12.2	5.0–6.7	2.9–4.3	8.6–10.8	28.5–31.8
Black	%	84.6	5.4	2.1	1.5	6.4	15.4
(n=473)	CI	80.8–87.8	3.6–8.0	1.0–4.2	0.7–3.0	4.4–9.4	12.2–19.2
Hispanic	%	81.9	6.0	4.6	1.9	5.6	18.1
(n=461)	CI	77.8–85.4	4.0–8.8	3.0–7.1	0.9–4.0	3.7–8.4	14.6–22.2
8th	%	81.0	9.3	3.3	1.8	4.5	19.0
(n=2,489)	CI	79.5–82.5	8.2–10.5	2.7–4.1	1.4–2.4	3.8–5.4	17.5–20.5
10th	%	72.7	10.3	4.9	3.1	9.0	27.3
(n=1,515)	CI	70.4–74.9	8.9–11.9	4.0–6.2	2.3–4.1	7.6–10.5	25.1–29.6
12th	%	63.4	9.9	7.7	4.5	14.5	36.6
(n=1,035)	CI	60.4–66.3	8.2–11.9	6.2–9.5	3.4–6.0	12.5–16.8	33.7–39.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used an e-cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	60.6	10.5	7.2	4.6	17.1	39.4
(n=5,046)	CI	59.1–62.0	9.6–11.4	6.5–8.0	4.0–5.3	16.0–18.3	38.0–40.9
Male	%	62.3	9.9	6.3	4.3	17.3	37.7
(n=2,513)	CI	60.2–64.3	8.8–11.2	5.3–7.4	3.5–5.2	15.7–19.0	35.7–39.8
Female	%	59.1	11.2	8.0	5.0	16.6	40.9
(n=2,409)	CI	57.0–61.1	10.0–12.6	6.9–9.3	4.2–6.1	15.1–18.3	38.9–43.0
White	%	56.4	11.4	8.4	5.6	18.3	43.6
(n=3,523)	CI	54.6–58.1	10.3–12.5	7.4–9.4	4.8–6.5	17.0–19.7	41.9–45.4
Black	%	81.5	5.4	3.2	1.7	8.2	18.5
(n=474)	CI	77.6–84.8	3.7–7.8	1.9–5.2	0.8–3.4	6.0–11.2	15.2–22.4
Hispanic	%	72.1	8.6	3.2	3.6	12.5	27.9
(n=458)	CI	67.4–76.3	6.3–11.8	1.8–5.4	2.1–6.0	9.5–16.2	23.7–32.6
8th	%	69.8	11.5	5.9	2.9	10.0	30.2
(n=2,489)	CI	67.9–71.5	10.3–12.8	5.1–6.9	2.3–3.6	8.8–11.2	28.5–32.1
10th	%	55.5	11.3	7.8	5.4	20.0	44.5
(n=1,521)	CI	53.0–58.0	9.8–13.0	6.6–9.3	4.4–6.6	18.1–22.1	42.0–47.0
12th	%	56.0	8.5	8.0	5.7	21.8	44.0
(n=1,036)	CI	53.0–59.0	7.0–10.4	6.5–9.8	4.4–7.2	19.4–24.4	41.0–47.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	68.6	9.2	6.2	3.6	12.5	31.4
(n=5,054)	CI	67.2–69.9	8.4–10.1	5.5–7.0	3.0–4.2	11.5–13.5	30.1–32.8
Male	%	71.7	7.8	5.3	3.3	11.9	28.3
(n=2,511)	CI	69.8–73.6	6.7–8.9	4.5–6.4	2.6–4.1	10.6–13.4	26.4–30.2
Female	%	66.0	10.4	6.9	3.9	12.7	34.0
(n=2,419)	CI	64.0–68.0	9.2–11.8	5.9–8.1	3.2–4.9	11.3–14.3	32.0–36.0
White	%	69.2	9.3	6.3	3.7	11.4	30.8
(n=3,526)	CI	67.6–70.8	8.3–10.4	5.5–7.3	3.1–4.5	10.3–12.6	29.2–32.4
Black	%	66.3	7.6	6.5	3.3	16.3	33.7
(n=470)	CI	61.6–70.8	5.4–10.5	4.4–9.3	2.0–5.5	12.9–20.4	29.2–38.4
Hispanic	%	75.2	7.4	5.2	2.3	9.9	24.8
(n=462)	CI	70.6–79.3	5.2–10.6	3.4–8.0	1.2–4.3	7.3–13.5	20.7–29.4
8th	%	81.0	7.3	4.1	2.1	5.6	19.0
(n=2,495)	CI	79.4–82.5	6.3–8.3	3.4–5.0	1.6–2.7	4.7–6.5	17.5–20.6
10th	%	64.3	10.6	6.4	4.5	14.2	35.7
(n=1,521)	CI	61.8–66.7	9.1–12.2	5.2–7.7	3.6–5.7	12.6–16.1	33.3–38.2
12th	%	59.8	9.8	8.2	4.1	18.1	40.2
(n=1,038)	CI	56.8–62.7	8.1–11.7	6.7–10.1	3.1–5.5	15.9–20.6	37.3–43.2

Note: The confidence interval (CI) is 95 percent.

Table BH3.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	11.1	6.3	12.5	70.2	82.6
(n=4,977)	CI	10.2–12.0	5.6–7.0	11.6–13.5	68.8–71.5	81.5–83.7
Male	%	12.7	6.4	13.0	68.0	81.0
(n=2,470)	CI	11.3–14.1	5.4–7.5	11.6–14.4	66.0–69.9	79.3–82.6
Female	%	9.1	6.3	11.7	72.9	84.6
(n=2,387)	CI	7.9–10.4	5.4–7.4	10.4–13.1	71.0–74.7	83.0–86.0
White	%	7.6	5.7	12.1	74.6	86.7
(n=3,482)	CI	6.7–8.6	5.0–6.6	11.0–13.3	73.1–76.1	85.5–87.8
Black	%	28.4	6.8	11.1	53.6	64.8
(n=460)	CI	24.2–33.0	4.7–9.8	8.4–14.6	48.8–58.4	60.0–69.3
Hispanic	%	14.9	8.6	16.5	60.0	76.5
(n=453)	CI	11.7–18.7	6.2–11.8	13.2–20.5	55.1–64.7	72.1–80.4
8th	%	9.4	7.2	13.4	70.0	83.4
(n=2,459)	CI	8.3–10.6	6.3–8.3	12.1–14.8	68.2–71.8	81.8–84.8
10th	%	11.7	6.0	11.1	71.2	82.3
(n=1,497)	CI	10.2–13.5	4.9–7.3	9.6–12.8	68.8–73.4	80.2–84.1
12th	%	12.2	5.6	13.0	69.2	82.3
(n=1,021)	CI	10.3–14.3	4.3–7.1	11.1–15.3	66.3–72.0	79.8–84.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.63. Perceived Risk of Regularly Drinking Alcohol

		How much do you think people risk harming themselves physically and in other ways when they <u>take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	14.8	15.5	22.6	47.0	69.7
(n=4,978)	CI	13.8–15.9	14.5–16.6	21.4–23.9	45.6–48.5	68.3–71.0
Male	%	17.1	16.2	23.1	43.5	66.6
(n=2,474)	CI	15.6–18.8	14.7–17.8	21.4–24.9	41.5–45.6	64.6–68.6
Female	%	11.9	14.9	22.1	51.1	73.2
(n=2,383)	CI	10.6–13.4	13.5–16.5	20.4–24.0	48.9–53.2	71.3–75.0
White	%	12.0	16.0	23.8	48.3	72.1
(n=3,484)	CI	10.9–13.2	14.7–17.3	22.3–25.3	46.5–50.0	70.5–73.6
Black	%	29.5	12.7	18.7	39.1	57.8
(n=459)	CI	25.3–34.2	9.8–16.2	15.2–22.7	34.5–44.0	53.0–62.5
Hispanic	%	17.6	14.2	20.5	47.7	68.2
(n=452)	CI	14.2–21.7	11.0–18.1	16.8–24.8	42.8–52.6	63.4–72.6
8th	%	12.5	14.8	21.6	51.0	72.6
(n=2,452)	CI	11.3–13.9	13.5–16.3	20.0–23.3	49.1–53.0	70.8–74.4
10th	%	15.1	15.1	22.8	47.0	69.9
(n=1,504)	CI	13.4–17.0	13.4–17.0	20.8–25.0	44.5–49.5	67.5–72.1
12th	%	17.0	16.6	23.5	42.8	66.3
(n=1,022)	CI	14.8–19.4	14.5–19.0	21.0–26.2	39.8–45.9	63.4–69.2

Note: The confidence interval (CI) is 95 percent.

Table BH3.64. Perceived Risk of Binge Drinking

		How much do you think people risk harming themselves physically and in other ways when they <u>take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	13.1	10.4	20.3	56.2	76.5
(n=4,991)	CI	12.1–14.1	9.5–11.4	19.2–21.5	54.8–57.7	75.3–77.8
Male	%	15.2	11.2	20.8	52.8	73.6
(n=2,479)	CI	13.8–16.8	9.9–12.6	19.1–22.5	50.7–54.8	71.7–75.4
Female	%	10.4	9.6	19.6	60.4	80.0
(n=2,392)	CI	9.2–11.8	8.4–10.9	18.0–21.3	58.3–62.4	78.2–81.6
White	%	9.9	10.9	20.7	58.4	79.2
(n=3,495)	CI	8.9–11.0	9.9–12.1	19.4–22.2	56.7–60.1	77.7–80.6
Black	%	29.0	10.0	16.0	45.0	61.0
(n=457)	CI	24.8–33.6	7.4–13.4	12.8–19.8	40.3–49.9	56.2–65.7
Hispanic	%	16.4	9.3	17.7	56.6	74.3
(n=453)	CI	13.1–20.4	6.7–12.7	14.3–21.7	51.7–61.4	69.7–78.4
8th	%	10.8	8.5	20.6	60.0	80.6
(n=2,457)	CI	9.7–12.1	7.5–9.7	19.0–22.2	58.1–62.0	79.0–82.2
10th	%	13.4	10.3	19.8	56.5	76.3
(n=1,508)	CI	11.7–15.2	8.9–12.0	17.9–21.9	54.0–59.0	74.1–78.4
12th	%	15.1	12.5	20.5	51.8	72.4
(n=1,026)	CI	13.1–17.4	10.6–14.7	18.2–23.1	48.8–54.9	69.6–75.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	11.3	7.3	16.9	64.6	81.4
(n=5,005)	CI	10.4–12.2	6.6–8.1	15.8–18.0	63.2–66.0	80.3–82.6
Male	%	12.6	7.2	18.2	62.0	80.2
(n=2,485)	CI	11.3–14.1	6.1–8.3	16.7–19.9	59.9–64.0	78.5–81.8
Female	%	9.4	7.3	15.3	68.0	83.3
(n=2,398)	CI	8.2–10.7	6.2–8.4	13.9–16.9	66.0–69.9	81.7–84.9
White	%	8.0	7.0	17.4	67.6	85.0
(n=3,507)	CI	7.1–9.0	6.1–7.9	16.1–18.8	66.0–69.3	83.7–86.3
Black	%	28.1	7.8	14.7	49.3	64.1
(n=462)	CI	23.9–32.7	5.5–11.0	11.6–18.5	44.5–54.1	59.3–68.6
Hispanic	%	15.6	8.8	19.5	56.1	75.6
(n=453)	CI	12.4–19.4	6.4–12.1	16.0–23.7	51.1–60.9	71.1–79.6
8th	%	9.4	6.6	17.3	66.7	84.0
(n=2,468)	CI	8.3–10.7	5.7–7.7	15.9–18.8	64.8–68.5	82.5–85.4
10th	%	11.7	6.5	15.5	66.2	81.7
(n=1,513)	CI	10.2–13.5	5.4–7.9	13.8–17.4	63.8–68.5	79.7–83.6
12th	%	12.7	8.8	17.9	60.6	78.4
(n=1,024)	CI	10.8–14.9	7.2–10.7	15.6–20.3	57.6–63.5	75.8–80.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	15.6	20.9	26.0	37.5	63.5
(n=4,993)	CI	14.6–16.7	19.8–22.1	24.7–27.3	36.1–38.9	62.0–64.9
Male	%	17.6	21.5	25.2	35.7	60.9
(n=2,483)	CI	16.0–19.2	19.8–23.3	23.4–27.1	33.8–37.7	58.9–62.9
Female	%	12.9	20.3	26.9	39.9	66.8
(n=2,389)	CI	11.5–14.4	18.7–22.1	25.0–28.8	37.8–42.0	64.7–68.7
White	%	12.8	22.1	27.2	37.9	65.1
(n=3,498)	CI	11.6–14.0	20.7–23.6	25.7–28.8	36.2–39.6	63.4–66.8
Black	%	30.9	18.3	18.3	32.5	50.8
(n=459)	CI	26.6–35.6	14.8–22.3	14.8–22.4	28.2–37.2	46.0–55.6
Hispanic	%	16.7	17.1	26.9	39.2	66.1
(n=451)	CI	13.4–20.7	13.7–21.3	22.8–31.5	34.5–44.1	61.3–70.6
8th	%	13.0	20.3	23.2	43.5	66.7
(n=2,467)	CI	11.8–14.4	18.7–21.9	21.6–24.9	41.6–45.5	64.8–68.5
10th	%	15.3	20.8	27.6	36.2	63.9
(n=1,507)	CI	13.6–17.2	18.8–22.9	25.4–29.9	33.9–38.7	61.4–66.3
12th	%	18.7	21.8	27.3	32.2	59.5
(n=1,019)	CI	16.5–21.2	19.3–24.4	24.7–30.1	29.4–35.1	56.5–62.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	30.2	21.5	14.7	33.6	48.3
(n=4,970)	CI	28.8–31.6	20.3–22.8	13.7–15.8	32.2–34.9	46.8–49.8
Male	%	32.4	19.9	14.0	33.7	47.7
(n=2,472)	CI	30.5–34.4	18.3–21.6	12.6–15.4	31.8–35.7	45.6–49.8
Female	%	27.1	23.5	15.5	33.9	49.5
(n=2,379)	CI	25.2–29.1	21.7–25.3	14.1–17.1	32.0–35.9	47.3–51.6
White	%	28.3	22.5	14.2	35.0	49.2
(n=3,478)	CI	26.8–30.0	21.0–24.0	13.0–15.4	33.4–36.7	47.4–50.9
Black	%	40.4	20.6	13.3	25.6	39.0
(n=461)	CI	35.7–45.2	17.0–24.9	10.4–16.9	21.7–30.0	34.4–43.7
Hispanic	%	26.9	16.8	21.4	34.9	56.3
(n=452)	CI	22.7–31.5	13.4–20.8	17.6–25.7	30.4–39.6	51.4–61.1
8th	%	21.9	17.7	16.5	43.9	60.5
(n=2,450)	CI	20.3–23.6	16.2–19.2	15.1–18.0	42.0–45.9	58.5–62.4
10th	%	32.3	23.9	14.2	29.6	43.8
(n=1,500)	CI	30.0–34.7	21.8–26.1	12.5–16.0	27.4–32.0	41.3–46.3
12th	%	36.8	23.2	13.4	26.6	40.0
(n=1,020)	CI	33.9–39.8	20.7–25.9	11.4–15.6	24.0–29.4	37.0–43.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	23.3	16.3	18.9	41.5	60.4
(n=4,976)	CI	22.1–24.6	15.2–17.5	17.7–20.0	40.1–42.9	58.9–61.8
Male	%	25.5	15.8	18.4	40.3	58.7
(n=2,475)	CI	23.7–27.4	14.3–17.4	16.8–20.0	38.3–42.4	56.6–60.7
Female	%	20.5	16.7	19.4	43.4	62.8
(n=2,381)	CI	18.8–22.3	15.2–18.4	17.8–21.1	41.3–45.5	60.7–64.8
White	%	20.2	16.2	19.3	44.2	63.6
(n=3,486)	CI	18.8–21.7	14.9–17.6	18.0–20.7	42.5–46.0	61.8–65.2
Black	%	37.4	16.2	18.3	28.1	46.4
(n=459)	CI	32.8–42.2	12.9–20.2	14.9–22.3	24.0–32.5	41.6–51.2
Hispanic	%	23.9	15.2	20.5	40.4	60.9
(n=449)	CI	20.0–28.4	11.9–19.2	16.8–24.8	35.7–45.2	55.9–65.6
8th	%	17.2	12.3	17.5	53.1	70.5
(n=2,453)	CI	15.7–18.7	11.1–13.7	16.0–19.0	51.1–55.0	68.7–72.3
10th	%	23.5	17.0	21.1	38.4	59.5
(n=1,499)	CI	21.4–25.7	15.2–19.0	19.1–23.3	35.9–40.9	57.0–62.0
12th	%	29.7	19.9	18.0	32.4	50.4
(n=1,024)	CI	27.0–32.6	17.6–22.5	15.7–20.4	29.6–35.4	47.3–53.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.1	4.7	11.3	80.9	92.2
(n=4,996)	CI	2.6–3.6	4.1–5.4	10.4–12.3	79.7–82.0	91.4–93.0
Male	%	3.4	4.2	11.2	81.1	92.4
(n=2,485)	CI	2.7–4.3	3.5–5.2	9.9–12.6	79.4–82.7	91.1–93.4
Female	%	2.4	4.9	11.3	81.5	92.8
(n=2,389)	CI	1.8–3.1	4.0–5.9	10.0–12.7	79.8–83.1	91.6–93.8
White	%	2.2	4.4	10.8	82.7	93.4
(n=3,502)	CI	1.7–2.8	3.7–5.2	9.7–11.9	81.3–84.0	92.5–94.3
Black	%	5.0	4.6	12.0	78.5	90.5
(n=461)	CI	3.1–7.8	2.9–7.3	9.1–15.6	74.1–82.3	87.0–93.1
Hispanic	%	2.8	5.5	13.4	78.3	91.7
(n=447)	CI	1.5–5.0	3.6–8.3	10.3–17.2	74.0–82.2	88.5–94.1
8th	%	1.7	3.8	10.0	84.5	94.5
(n=2,461)	CI	1.3–2.3	3.1–4.6	8.9–11.2	83.0–85.9	93.5–95.3
10th	%	3.2	5.0	10.6	81.2	91.8
(n=1,512)	CI	2.5–4.3	4.0–6.2	9.1–12.2	79.2–83.1	90.3–93.1
12th	%	4.3	5.4	13.6	76.7	90.3
(n=1,023)	CI	3.2–5.7	4.2–7.0	11.6–15.8	74.0–79.2	88.3–91.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.7	11.1	19.4	63.8	83.2
(n=5,025)	CI	5.0–6.5	10.2–12.0	18.2–20.6	62.4–65.3	82.1–84.3
Male	%	6.9	11.0	18.3	63.8	82.1
(n=2,499)	CI	5.8–8.1	9.7–12.4	16.8–20.0	61.8–65.8	80.4–83.7
Female	%	3.9	11.2	20.5	64.3	84.8
(n=2,403)	CI	3.2–4.9	9.9–12.7	18.9–22.3	62.2–66.3	83.2–86.3
White	%	5.6	11.5	19.7	63.2	82.9
(n=3,520)	CI	4.8–6.5	10.4–12.7	18.3–21.2	61.4–64.8	81.5–84.2
Black	%	5.3	7.5	16.1	71.2	87.2
(n=465)	CI	3.4–8.2	5.3–10.5	12.8–20.0	66.5–75.4	83.5–90.2
Hispanic	%	4.2	8.8	19.1	67.9	87.0
(n=450)	CI	2.5–6.9	6.3–12.2	15.5–23.3	63.0–72.4	83.1–90.1
8th	%	2.2	7.6	15.0	75.2	90.2
(n=2,480)	CI	1.7–2.9	6.6–8.7	13.7–16.5	73.4–76.8	89.0–91.3
10th	%	6.3	11.2	20.6	61.9	82.5
(n=1,519)	CI	5.2–7.7	9.7–12.9	18.7–22.8	59.4–64.3	80.5–84.3
12th	%	8.8	14.7	22.8	53.7	76.5
(n=1,026)	CI	7.2–10.7	12.7–17.0	20.3–25.4	50.6–56.7	73.8–78.9

Note: The confidence interval (CI) is 95 percent.

Table BH3.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.4	9.5	18.6	64.5	83.1
(n=5,021)	CI	6.6–8.2	8.7–10.5	17.5–19.8	63.0–65.9	81.9–84.2
Male	%	8.3	9.6	16.7	65.4	82.1
(n=2,499)	CI	7.1–9.6	8.4–10.9	15.2–18.4	63.3–67.4	80.4–83.7
Female	%	5.8	9.6	20.5	64.1	84.6
(n=2,399)	CI	4.9–7.0	8.3–10.9	18.8–22.3	62.0–66.2	82.9–86.1
White	%	7.5	10.2	19.1	63.2	82.3
(n=3,520)	CI	6.5–8.5	9.1–11.3	17.8–20.6	61.5–64.9	80.9–83.7
Black	%	5.5	6.4	18.2	69.9	88.2
(n=463)	CI	3.5–8.4	4.3–9.3	14.7–22.3	65.2–74.3	84.4–91.1
Hispanic	%	4.5	7.7	16.1	71.7	87.8
(n=450)	CI	2.7–7.2	5.4–11.0	12.7–20.1	67.0–76.0	84.0–90.8
8th	%	2.1	5.6	14.4	78.0	92.3
(n=2,478)	CI	1.6–2.7	4.7–6.5	13.0–15.8	76.3–79.6	91.2–93.3
10th	%	6.7	9.8	19.8	63.7	83.5
(n=1,517)	CI	5.6–8.1	8.4–11.4	17.9–21.9	61.2–66.0	81.5–85.3
12th	%	13.8	13.6	22.0	50.7	72.7
(n=1,026)	CI	11.8–16.0	11.6–15.8	19.6–24.6	47.6–53.7	69.9–75.3

Note: The confidence interval (CI) is 95 percent.

Table BH3.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.6	15.5	17.2	52.7	69.9
(n=5,014)	CI	13.6–15.8	14.4–16.6	16.1–18.4	51.2–54.2	68.5–71.3
Male	%	15.7	13.4	17.1	53.8	70.9
(n=2,495)	CI	14.2–17.4	12.0–14.9	15.5–18.7	51.7–55.9	68.9–72.7
Female	%	13.1	17.2	17.5	52.3	69.7
(n=2,397)	CI	11.7–14.6	15.6–18.9	15.9–19.1	50.1–54.4	67.7–71.6
White	%	15.5	16.5	17.1	50.9	68.0
(n=3,516)	CI	14.2–16.9	15.2–17.8	15.9–18.5	49.1–52.6	66.3–69.7
Black	%	8.6	11.5	15.8	64.1	79.9
(n=461)	CI	6.2–11.8	8.7–15.0	12.6–19.8	59.3–68.6	75.7–83.6
Hispanic	%	9.0	11.9	18.9	60.3	79.1
(n=449)	CI	6.5–12.3	9.0–15.5	15.3–23.1	55.3–65.0	74.8–82.9
8th	%	8.2	12.7	13.9	65.2	79.1
(n=2,476)	CI	7.2–9.4	11.4–14.0	12.6–15.3	63.3–67.1	77.4–80.6
10th	%	13.7	17.0	19.1	50.2	69.3
(n=1,513)	CI	12.0–15.5	15.2–19.0	17.2–21.1	47.7–52.7	66.9–71.6
12th	%	22.6	16.8	18.8	41.8	60.6
(n=1,025)	CI	20.1–25.3	14.6–19.2	16.5–21.3	38.8–44.8	57.6–63.6

Note: The confidence interval (CI) is 95 percent.

Table BH3.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	13.3	11.7	14.2	60.9	75.1
(n=5,005)	CI	12.3–14.3	10.7–12.7	13.2–15.3	59.4–62.3	73.7–76.4
Male	%	14.0	10.4	14.0	61.6	75.6
(n=2,489)	CI	12.5–15.6	9.2–11.8	12.6–15.5	59.6–63.7	73.7–77.4
Female	%	11.9	12.7	14.5	60.9	75.4
(n=2,395)	CI	10.5–13.4	11.3–14.3	13.1–16.1	58.8–63.0	73.5–77.2
White	%	12.9	11.1	14.1	62.0	76.0
(n=3,503)	CI	11.7–14.2	10.0–12.3	12.9–15.3	60.2–63.7	74.5–77.6
Black	%	12.6	15.2	13.4	58.8	72.2
(n=462)	CI	9.6–16.4	12.0–19.1	10.4–17.2	53.9–63.5	67.5–76.4
Hispanic	%	9.4	11.5	14.4	64.8	79.1
(n=450)	CI	6.8–13.0	8.6–15.2	11.2–18.2	59.8–69.4	74.6–83.0
8th	%	5.1	7.2	12.2	75.4	87.7
(n=2,470)	CI	4.3–6.1	6.2–8.3	11.0–13.6	73.7–77.1	86.3–88.9
10th	%	15.0	12.8	14.0	58.1	72.2
(n=1,510)	CI	13.3–16.9	11.2–14.6	12.4–15.9	55.6–60.6	69.8–74.4
12th	%	20.2	15.2	16.5	48.1	64.6
(n=1,025)	CI	17.8–22.7	13.2–17.6	14.3–18.9	45.1–51.2	61.6–67.5

Note: The confidence interval (CI) is 95 percent.

Table BH3.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	65.3	24.4	10.4	34.7
(n=4,988)	CI	63.9–66.6	23.1–25.6	9.5–11.3	33.4–36.1
Male	%	66.7	23.6	9.7	33.3
(n=2,477)	CI	64.8–68.6	21.9–25.4	8.5–10.9	31.4–35.2
Female	%	63.6	25.3	11.1	36.4
(n=2,390)	CI	61.5–65.6	23.5–27.1	9.9–12.5	34.4–38.5
White	%	65.8	24.4	9.9	34.2
(n=3,496)	CI	64.1–67.4	22.9–25.9	8.9–10.9	32.6–35.9
Black	%	70.8	20.1	9.0	29.2
(n=454)	CI	66.3–75.1	16.5–24.3	6.6–12.2	24.9–33.7
Hispanic	%	59.2	28.8	11.9	40.8
(n=454)	CI	54.4–63.9	24.6–33.4	9.2–15.5	36.1–45.6
8th	%	61.3	26.9	11.9	38.7
(n=2,460)	CI	59.3–63.2	25.1–28.6	10.6–13.2	36.8–40.7
10th	%	66.2	23.8	10.0	33.8
(n=1,510)	CI	63.8–68.5	21.7–26.0	8.6–11.6	31.5–36.2
12th	%	68.6	22.3	9.2	31.4
(n=1,018)	CI	65.6–71.3	19.8–24.9	7.6–11.1	28.7–34.4

Note: The confidence interval (CI) is 95 percent.

Table BH3.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	50.5	32.5	16.9	49.5
(n=4,991)	CI	49.1–52.0	31.2–33.9	15.9–18.0	48.0–50.9
Male	%	52.2	32.3	15.6	47.8
(n=2,473)	CI	50.1–54.2	30.4–34.2	14.1–17.1	45.8–49.9
Female	%	48.5	33.1	18.3	51.5
(n=2,397)	CI	46.4–50.7	31.2–35.1	16.8–20.0	49.3–53.6
White	%	49.4	33.5	17.0	50.6
(n=3,501)	CI	47.7–51.2	31.9–35.2	15.8–18.3	48.8–52.3
Black	%	60.6	26.6	12.8	39.4
(n=452)	CI	55.8–65.2	22.6–31.0	9.9–16.4	34.8–44.2
Hispanic	%	45.3	35.5	19.2	54.7
(n=454)	CI	40.5–50.3	31.0–40.2	15.7–23.3	49.7–59.5
8th	%	45.1	35.0	19.9	54.9
(n=2,459)	CI	43.1–47.1	33.2–36.9	18.3–21.5	52.9–56.9
10th	%	49.0	34.3	16.6	51.0
(n=1,512)	CI	46.5–51.5	32.0–36.8	14.9–18.6	48.5–53.5
12th	%	58.1	27.9	14.0	41.9
(n=1,020)	CI	55.0–61.1	25.2–30.7	12.0–16.3	38.9–45.0

Note: The confidence interval (CI) is 95 percent.

Table BH3.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	41.6	31.8	26.6	58.4
(n=4,994)	CI	40.2–43.1	30.4–33.1	25.3–27.9	56.9–59.8
Male	%	44.5	29.5	26.1	55.5
(n=2,475)	CI	42.4–46.6	27.6–31.4	24.3–27.9	53.4–57.6
Female	%	38.3	34.5	27.2	61.7
(n=2,398)	CI	36.3–40.4	32.5–36.5	25.4–29.1	59.6–63.7
White	%	40.4	33.3	26.4	59.6
(n=3,501)	CI	38.7–42.1	31.6–34.9	24.9–27.9	57.9–61.3
Black	%	52.8	24.0	23.2	47.2
(n=454)	CI	47.9–57.6	20.1–28.4	19.4–27.5	42.4–52.1
Hispanic	%	39.7	32.4	27.8	60.3
(n=454)	CI	35.0–44.7	28.1–37.1	23.6–32.5	55.3–65.0
8th	%	39.1	31.7	29.2	60.9
(n=2,458)	CI	37.2–41.0	29.9–33.5	27.5–31.1	59.0–62.8
10th	%	40.7	32.9	26.4	59.3
(n=1,511)	CI	38.3–43.2	30.6–35.3	24.2–28.7	56.8–61.7
12th	%	45.4	30.6	24.0	54.6
(n=1,025)	CI	42.4–48.5	27.9–33.5	21.4–26.7	51.5–57.6

Note: The confidence interval (CI) is 95 percent.

Behavioral Health Planning Region 5 Results Summary

Table BH5.1. Lifetime Prescription Drug Misuse

		On how many occasions (if any) have you ever used prescription drugs not prescribed to you?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	90.4	5.0	1.8	1.0	0.5	0.3	1.0	9.6
(n=4,312)	CI	89.4–91.3	4.3–5.7	1.4–2.3	0.7–1.4	0.3–0.8	0.2–0.5	0.7–1.4	8.7–10.6
Male	%	90.3	4.3	1.9	1.3	0.6	0.4	1.2	9.7
(n=2,092)	CI	88.9–91.5	3.5–5.3	1.4–2.6	0.9–2.0	0.3–1.1	0.2–0.8	0.8–1.8	8.5–11.1
Female	%	91.4	5.2	1.6	0.7	0.4	0.2	0.5	8.6
(n=2,099)	CI	90.1–92.6	4.3–6.3	1.2–2.3	0.4–1.1	0.2–0.7	0.1–0.5	0.3–1.0	7.4–9.9
White	%	90.8	5.0	1.7	0.9	0.4	0.2	0.9	9.2
(n=3,506)	CI	89.8–91.8	4.3–5.8	1.3–2.3	0.6–1.3	0.2–0.7	0.1–0.4	0.6–1.3	8.2–10.2
Black	%	86.9	3.6	1.7	1.9	0.0	2.7	3.2	13.1
(n=130)	CI	79.5–91.9	1.5–8.6	0.4–6.8	0.5–7.3	–	0.9–8.2	1.2–8.4	8.1–20.5
Hispanic	%	90.2	3.1	2.4	3.2	0.0	0.0	1.1	9.8
(n=103)	CI	82.3–94.8	1.2–8.2	0.6–9.1	1.0–9.9	–	–	0.2–7.8	5.2–17.7
8th	%	95.0	3.3	0.6	0.4	0.2	0.1	0.3	5.0
(n=1,892)	CI	94.0–95.9	2.6–4.2	0.4–1.1	0.2–0.8	0.1–0.6	0.0–0.4	0.1–0.7	4.1–6.0
10th	%	89.0	5.4	2.0	1.0	0.5	0.3	1.8	11.0
(n=1,299)	CI	87.1–90.5	4.3–6.7	1.4–2.9	0.6–1.7	0.3–1.1	0.1–0.8	1.2–2.7	9.5–12.9
12th	%	86.9	6.3	2.8	1.7	0.8	0.5	0.9	13.1
(n=1,121)	CI	84.8–88.8	5.0–7.9	2.0–4.0	1.1–2.6	0.4–1.5	0.2–1.2	0.5–1.6	11.2–15.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.2. Lifetime Alcohol Use

		On how many occasions (if any) have you ever had one or more drinks of an alcoholic beverage?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	59.6	16.0	7.3	4.4	4.4	2.7	5.6	40.4
(n=4,323)	CI	58.1–61.1	14.9–17.2	6.5–8.1	3.8–5.1	3.8–5.1	2.2–3.3	4.9–6.4	38.9–41.9
Male	%	59.5	15.4	6.7	4.0	4.5	2.9	7.1	40.5
(n=2,100)	CI	57.3–61.7	13.8–17.0	5.6–7.9	3.2–5.0	3.7–5.6	2.2–3.7	6.0–8.3	38.3–42.7
Female	%	60.5	16.9	7.8	4.6	4.2	2.5	3.5	39.5
(n=2,101)	CI	58.3–62.6	15.3–18.6	6.7–9.1	3.7–5.6	3.4–5.2	1.9–3.4	2.8–4.5	37.4–41.7
White	%	59.0	16.1	7.3	4.6	4.5	2.6	5.9	41.0
(n=3,512)	CI	57.3–60.6	14.9–17.4	6.5–8.3	3.9–5.4	3.8–5.3	2.1–3.3	5.1–6.7	39.4–42.7
Black	%	57.8	16.8	10.9	3.6	4.5	2.6	3.7	42.2
(n=132)	CI	48.9–66.2	11.3–24.4	6.4–18.1	1.4–9.4	1.9–10.4	0.8–8.0	1.4–9.5	33.8–51.1
Hispanic	%	64.8	12.4	4.7	4.3	5.8	1.9	6.0	35.2
(n=103)	CI	54.7–73.8	7.2–20.6	1.9–11.1	1.6–11.1	2.4–13.4	0.5–7.7	2.5–13.8	26.2–45.3
8th	%	78.8	13.6	3.4	1.8	0.8	0.8	0.8	21.2
(n=1,903)	CI	76.9–80.6	12.1–15.2	2.7–4.3	1.3–2.5	0.5–1.4	0.5–1.3	0.5–1.4	19.4–23.1
10th	%	55.2	17.8	8.2	5.0	5.8	2.8	5.1	44.8
(n=1,295)	CI	52.5–57.9	15.9–20.0	6.9–9.9	3.9–6.3	4.7–7.3	2.0–3.8	4.0–6.4	42.1–47.5
12th	%	43.6	16.7	10.4	6.6	6.7	4.7	11.3	56.4
(n=1,125)	CI	40.7–46.5	14.6–18.9	8.7–12.3	5.3–8.3	5.4–8.4	3.6–6.1	9.6–13.3	53.5–59.3

Note: The confidence interval (CI) is 95 percent.

Table BH5.3. Lifetime Binge Drinking

		On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.3	6.8	4.0	2.5	2.7	1.8	2.9	20.7
(n=4,324)	CI	78.0–80.5	6.1–7.6	3.4–4.6	2.1–3.1	2.2–3.3	1.4–2.3	2.4–3.5	19.5–22.0
Male	%	77.9	5.5	3.8	2.9	3.4	2.5	4.0	22.1
(n=2,094)	CI	75.9–79.7	4.6–6.6	3.0–4.7	2.3–3.8	2.6–4.3	1.9–3.3	3.2–5.0	20.3–24.1
Female	%	81.7	8.0	3.9	1.9	2.1	0.9	1.5	18.3
(n=2,108)	CI	79.9–83.4	6.8–9.3	3.1–4.9	1.4–2.7	1.5–2.9	0.5–1.4	1.1–2.2	16.6–20.1
White	%	79.0	6.8	4.0	2.5	2.8	1.8	3.1	21.0
(n=3,515)	CI	77.5–80.4	6.0–7.7	3.4–4.8	2.0–3.1	2.2–3.4	1.4–2.3	2.5–3.8	19.6–22.5
Black	%	78.6	7.3	4.4	2.7	2.7	1.5	2.8	21.4
(n=131)	CI	70.0–85.2	3.7–14.0	1.8–10.3	0.9–8.2	0.9–8.2	0.3–5.9	0.9–8.4	14.8–30.0
Hispanic	%	82.0	5.8	1.9	4.2	2.4	1.2	2.4	18.0
(n=103)	CI	72.5–88.7	2.4–13.4	0.5–7.7	1.5–10.9	0.6–9.4	0.2–8.4	0.6–9.4	11.3–27.5
8th	%	94.7	2.6	1.4	0.6	0.3	0.4	0.1	5.3
(n=1,899)	CI	93.6–95.7	2.0–3.4	0.9–2.0	0.3–1.0	0.1–0.6	0.2–0.8	0.0–0.4	4.3–6.4
10th	%	77.9	7.8	4.3	3.0	2.5	1.8	2.6	22.1
(n=1,299)	CI	75.6–80.1	6.5–9.4	3.3–5.5	2.2–4.1	1.8–3.5	1.2–2.7	1.9–3.6	19.9–24.4
12th	%	64.2	10.3	6.4	4.1	5.6	3.3	6.3	35.8
(n=1,126)	CI	61.3–66.9	8.6–12.2	5.1–8.0	3.1–5.4	4.4–7.1	2.4–4.5	5.0–7.9	33.1–38.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.4. Lifetime Cigarette Use

		On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	82.8	7.6	2.8	1.4	1.2	1.1	3.0	17.2
(n=4,324)	CI	81.6–83.9	6.8–8.5	2.4–3.4	1.1–1.9	0.9–1.6	0.8–1.5	2.5–3.6	16.1–18.4
Male	%	81.6	7.3	2.8	1.6	1.6	1.4	3.7	18.4
(n=2,101)	CI	79.8–83.3	6.2–8.6	2.1–3.7	1.1–2.3	1.1–2.2	0.9–2.1	2.9–4.7	16.7–20.2
Female	%	85.0	7.7	2.7	1.1	0.9	0.6	2.0	15.0
(n=2,101)	CI	83.3–86.6	6.6–9.0	2.1–3.6	0.7–1.7	0.5–1.4	0.3–1.0	1.4–2.7	13.4–16.7
White	%	81.8	8.2	2.9	1.6	1.2	1.1	3.0	18.2
(n=3,517)	CI	80.5–83.1	7.3–9.2	2.4–3.6	1.2–2.1	0.9–1.7	0.8–1.6	2.5–3.7	16.9–19.5
Black	%	87.2	3.6	2.5	0.9	0.9	1.8	3.2	12.8
(n=131)	CI	79.7–92.2	1.3–9.2	0.8–7.6	0.1–6.0	0.1–6.0	0.4–7.0	1.2–8.5	7.8–20.3
Hispanic	%	83.6	7.0	3.9	1.1	2.0	0.0	2.4	16.4
(n=103)	CI	74.4–90.0	3.1–14.8	1.4–10.3	0.2–7.8	0.5–8.1	–	0.6–9.1	10.0–25.6
8th	%	94.2	3.3	0.9	0.4	0.6	0.2	0.5	5.8
(n=1,899)	CI	93.0–95.1	2.6–4.2	0.6–1.4	0.2–0.8	0.4–1.1	0.1–0.6	0.2–0.9	4.9–7.0
10th	%	80.9	8.7	3.2	1.2	1.1	1.1	3.6	19.1
(n=1,300)	CI	78.7–83.0	7.3–10.4	2.4–4.3	0.8–2.0	0.7–1.9	0.7–1.9	2.7–4.8	17.0–21.3
12th	%	72.5	11.1	4.5	2.8	1.9	1.9	5.1	27.5
(n=1,125)	CI	69.9–75.1	9.4–13.0	3.4–5.9	2.0–4.0	1.3–2.9	1.3–2.9	4.0–6.6	24.9–30.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	67.8	9.1	3.9	2.2	2.2	1.9	12.9	32.2
(n=4,316)	CI	66.4-69.3	8.3-10.0	3.3-4.5	1.8-2.8	1.7-2.7	1.5-2.4	11.9-14.0	30.7-33.6
Male	%	65.2	9.1	4.2	1.9	1.6	1.8	16.2	34.8
(n=2,094)	CI	63.1-67.3	7.9-10.5	3.4-5.2	1.4-2.6	1.1-2.3	1.3-2.5	14.6-17.9	32.7-36.9
Female	%	71.1	9.4	3.6	2.5	2.6	2.0	8.9	28.9
(n=2,100)	CI	69.0-73.1	8.2-10.8	2.8-4.5	1.9-3.3	1.9-3.4	1.4-2.7	7.7-10.3	26.9-31.0
White	%	66.9	9.4	3.9	2.2	2.3	2.0	13.3	33.1
(n=3,509)	CI	65.3-68.5	8.5-10.4	3.3-4.6	1.8-2.8	1.8-2.8	1.6-2.6	12.2-14.6	31.5-34.7
Black	%	64.2	7.6	4.4	1.5	3.6	1.8	16.8	35.8
(n=132)	CI	55.3-72.3	4.1-13.8	1.9-9.7	0.4-6.1	1.4-9.4	0.4-6.9	11.0-24.7	27.7-44.7
Hispanic	%	76.6	3.0	2.7	2.4	0.8	2.4	12.1	23.4
(n=103)	CI	66.8-84.2	1.0-9.3	0.8-8.3	0.6-9.4	0.1-5.5	0.6-9.1	6.8-20.8	15.8-33.2
8th	%	85.6	6.4	2.4	1.4	0.9	0.4	2.8	14.4
(n=1,901)	CI	84.0-87.1	5.4-7.6	1.8-3.2	1.0-2.1	0.6-1.5	0.2-0.8	2.2-3.7	12.9-16.0
10th	%	62.3	11.3	4.4	1.9	2.4	2.2	15.5	37.7
(n=1,293)	CI	59.7-64.9	9.7-13.2	3.4-5.7	1.3-2.8	1.7-3.4	1.5-3.1	13.6-17.6	35.1-40.3
12th	%	54.5	9.7	5.0	3.5	3.2	3.2	21.0	45.5
(n=1,122)	CI	51.6-57.4	8.1-11.6	3.8-6.4	2.5-4.7	2.3-4.4	2.3-4.4	18.7-23.4	42.6-48.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1-2 times	3-5 times	6-9 times	10-19 times	20-39 times	40+ times	Any Use
Total	%	88.5	3.8	1.4	0.9	0.9	0.8	3.7	11.5
(n=4,318)	CI	87.5-89.5	3.2-4.4	1.1-1.8	0.7-1.3	0.7-1.3	0.5-1.1	3.1-4.3	10.5-12.5
Male	%	82.2	5.0	2.4	1.5	1.6	1.0	6.3	17.8
(n=2,095)	CI	80.4-83.8	4.1-6.1	1.8-3.2	1.0-2.2	1.1-2.2	0.7-1.6	5.3-7.5	16.2-19.6
Female	%	95.8	2.5	0.3	0.2	0.3	0.3	0.7	4.2
(n=2,101)	CI	94.8-96.6	1.8-3.3	0.2-0.7	0.1-0.6	0.1-0.7	0.1-0.6	0.4-1.2	3.4-5.2
White	%	88.0	3.8	1.6	0.9	0.9	0.9	3.8	12.0
(n=3,510)	CI	86.8-89.1	3.2-4.6	1.2-2.1	0.6-1.3	0.7-1.3	0.6-1.3	3.2-4.6	10.9-13.2
Black	%	88.1	4.1	0.0	2.4	0.0	0.0	5.4	11.9
(n=132)	CI	80.7-92.9	1.7-9.8	-	0.7-7.3	-	-	2.4-11.6	7.1-19.3
Hispanic	%	85.8	4.5	3.1	3.2	1.1	0.0	2.3	14.2
(n=103)	CI	76.9-91.7	1.6-11.6	1.0-9.3	1.0-9.9	0.2-7.8	-	0.6-8.8	8.3-23.1
8th	%	96.0	2.3	0.6	0.2	0.3	0.2	0.4	4.0
(n=1,902)	CI	95.0-96.8	1.7-3.0	0.3-1.0	0.1-0.6	0.1-0.7	0.1-0.6	0.2-0.8	3.2-5.0
10th	%	87.4	3.5	1.8	0.8	1.2	0.7	4.7	12.6
(n=1,292)	CI	85.4-89.1	2.6-4.6	1.2-2.7	0.5-1.5	0.7-1.9	0.4-1.3	3.7-6.0	10.9-14.6
12th	%	81.7	5.7	2.0	1.8	1.3	1.4	6.1	18.3
(n=1,124)	CI	79.4-83.9	4.5-7.2	1.3-2.9	1.1-2.7	0.8-2.2	0.9-2.3	4.9-7.7	16.1-20.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	78.8	5.6	3.1	2.0	2.5	1.7	6.4	21.2
(n=4,316)	CI	77.5–80.1	4.9–6.3	2.6–3.7	1.6–2.5	2.0–3.0	1.3–2.1	5.7–7.2	19.9–22.5
Male	%	78.6	4.9	3.3	1.5	2.3	1.9	7.6	21.4
(n=2,092)	CI	76.7–80.4	4.0–6.0	2.6–4.2	1.0–2.1	1.7–3.1	1.3–2.6	6.4–8.9	19.6–23.3
Female	%	80.0	6.5	2.8	2.4	2.5	1.3	4.6	20.0
(n=2,103)	CI	78.1–81.7	5.5–7.8	2.1–3.6	1.8–3.2	1.9–3.3	0.8–1.9	3.7–5.7	18.3–21.9
White	%	78.5	5.7	3.2	2.0	2.5	1.6	6.5	21.5
(n=3,510)	CI	77.0–79.9	5.0–6.6	2.6–3.9	1.5–2.5	2.0–3.2	1.2–2.1	5.6–7.4	20.1–23.0
Black	%	74.4	4.1	2.7	3.2	3.2	5.2	7.0	25.6
(n=132)	CI	65.8–81.5	1.7–9.8	0.9–8.1	1.2–8.5	1.2–8.5	2.3–11.3	3.5–13.5	18.5–34.2
Hispanic	%	80.6	7.5	2.4	0.0	0.0	1.1	8.4	19.4
(n=102)	CI	71.1–87.6	3.7–14.6	0.6–9.2	–	–	0.2–7.9	4.0–16.7	12.4–28.9
8th	%	95.5	2.2	0.6	0.5	0.5	0.3	0.4	4.5
(n=1,898)	CI	94.5–96.4	1.6–2.9	0.4–1.1	0.3–1.0	0.3–1.0	0.1–0.6	0.2–0.8	3.6–5.5
10th	%	76.2	6.1	3.0	2.4	2.7	2.1	7.5	23.8
(n=1,298)	CI	73.8–78.4	4.9–7.5	2.2–4.1	1.7–3.4	1.9–3.7	1.5–3.1	6.2–9.1	21.6–26.2
12th	%	63.6	8.7	5.8	3.1	4.4	2.7	11.7	36.4
(n=1,120)	CI	60.8–66.4	7.2–10.5	4.6–7.3	2.2–4.3	3.3–5.7	1.9–3.8	10.0–13.8	33.6–39.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	98.4	0.8	0.2	0.2	0.1	0.1	0.3	1.6
(n=4,318)	CI	97.9–98.7	0.6–1.1	0.1–0.4	0.1–0.4	0.0–0.2	0.0–0.2	0.2–0.6	1.3–2.1
Male	%	98.3	0.9	0.2	0.3	0.0	0.0	0.3	1.7
(n=2,096)	CI	97.6–98.8	0.6–1.5	0.1–0.5	0.1–0.6	0.0–0.3	–	0.1–0.7	1.2–2.4
Female	%	99.0	0.6	0.2	0.1	0.1	0.0	0.1	1.0
(n=2,101)	CI	98.5–99.4	0.3–1.1	0.1–0.5	0.0–0.4	0.0–0.4	–	0.0–0.4	0.6–1.5
White	%	98.4	0.9	0.2	0.2	0.1	0.1	0.2	1.6
(n=3,510)	CI	98.0–98.8	0.6–1.3	0.1–0.4	0.1–0.4	0.0–0.3	0.0–0.3	0.1–0.4	1.2–2.0
Black	%	93.8	1.8	0.9	0.9	0.0	0.0	2.7	6.2
(n=132)	CI	87.5–97.0	0.4–6.9	0.1–6.0	0.1–6.0	–	–	0.9–8.1	3.0–12.5
Hispanic	%	98.1	0.8	0.0	0.0	0.0	0.0	1.1	1.9
(n=103)	CI	92.3–99.5	0.1–5.5	–	–	–	–	0.2–7.8	0.5–7.7
8th	%	99.5	0.2	0.0	0.1	0.1	0.0	0.1	0.5
(n=1,897)	CI	99.0–99.7	0.1–0.6	–	0.0–0.4	0.0–0.4	–	0.0–0.4	0.3–1.0
10th	%	98.0	1.1	0.2	0.3	0.1	0.1	0.2	2.0
(n=1,300)	CI	97.1–98.6	0.6–1.8	0.1–0.7	0.1–0.8	0.0–0.5	0.0–0.5	0.1–0.7	1.4–2.9
12th	%	97.5	1.2	0.4	0.2	0.1	0.1	0.6	2.5
(n=1,121)	CI	96.4–98.3	0.7–2.0	0.1–0.9	0.0–0.7	0.0–0.6	0.0–0.6	0.3–1.3	1.7–3.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	96.2	1.8	0.6	0.4	0.3	0.1	0.6	3.8
(n=4,323)	CI	95.5–96.8	1.4–2.3	0.4–1.0	0.2–0.6	0.2–0.6	0.0–0.3	0.4–0.8	3.2–4.5
Male	%	95.2	2.2	0.7	0.6	0.5	0.1	0.6	4.8
(n=2,098)	CI	94.2–96.1	1.6–3.0	0.4–1.2	0.3–1.1	0.3–1.0	0.0–0.4	0.3–1.1	3.9–5.8
Female	%	97.9	1.1	0.5	0.1	0.1	0.1	0.2	2.1
(n=2,104)	CI	97.1–98.5	0.7–1.7	0.3–0.9	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.6	1.5–2.9
White	%	96.1	2.0	0.7	0.4	0.3	0.1	0.4	3.9
(n=3,514)	CI	95.3–96.7	1.5–2.5	0.4–1.1	0.2–0.7	0.2–0.6	0.1–0.4	0.2–0.7	3.3–4.7
Black	%	94.7	0.0	0.9	0.0	0.9	0.0	3.6	5.3
(n=132)	CI	88.5–97.6	–	0.1–6.0	–	0.1–6.4	–	1.3–9.2	2.4–11.5
Hispanic	%	95.7	3.1	0.0	0.0	0.0	0.0	1.1	4.3
(n=103)	CI	88.9–98.4	1.0–9.6	–	–	–	–	0.2–7.8	1.6–11.1
8th	%	99.2	0.5	0.1	0.0	0.0	0.0	0.3	0.8
(n=1,896)	CI	98.6–99.5	0.3–1.0	0.0–0.4	–	–	–	0.1–0.6	0.5–1.4
10th	%	95.7	1.8	0.8	0.6	0.3	0.1	0.6	4.3
(n=1,301)	CI	94.5–96.7	1.2–2.7	0.5–1.5	0.3–1.2	0.1–0.8	0.0–0.5	0.3–1.2	3.3–5.5
12th	%	93.5	3.1	1.1	0.5	0.7	0.3	0.8	6.5
(n=1,126)	CI	91.9–94.8	2.2–4.3	0.6–1.9	0.2–1.2	0.4–1.4	0.1–0.8	0.4–1.5	5.2–8.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.10. Age of First Prescription Drug Misuse

		How old were you when you first <u>used prescription drugs not prescribed to you</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	91.4	1.1	0.5	0.7	0.9	1.2	2.0	1.3	0.9	13.90
(n=4,337)	CI	90.5–92.3	0.8–1.5	0.3–0.7	0.5–1.0	0.7–1.2	0.9–1.6	1.6–2.5	1.0–1.7	0.6–1.3	13.7–14.1
Male	%	91.3	1.0	0.6	0.7	0.7	0.8	2.4	1.4	1.0	14.04
(n=2,108)	CI	90.0–92.5	0.7–1.6	0.3–1.0	0.4–1.2	0.5–1.2	0.5–1.4	1.8–3.2	0.9–2.1	0.6–1.6	13.7–14.4
Female	%	92.3	1.0	0.4	0.6	0.9	1.3	1.6	1.1	0.8	13.87
(n=2,107)	CI	91.0–93.4	0.6–1.5	0.2–0.8	0.3–1.0	0.6–1.4	0.9–1.9	1.1–2.3	0.7–1.7	0.5–1.4	13.5–14.2
White	%	91.6	1.0	0.4	0.6	0.9	1.2	2.0	1.3	1.0	14.01
(n=3,525)	CI	90.6–92.5	0.7–1.4	0.3–0.7	0.4–0.9	0.6–1.2	0.9–1.7	1.6–2.6	0.9–1.7	0.7–1.4	13.8–14.3
Black	%	90.0	1.8	0.0	2.4	0.6	2.6	2.6	0.0	0.0	13.02
(n=132)	CI	83.0–94.3	0.4–6.9	–	0.7–7.3	0.1–4.2	0.8–8.0	0.8–8.0	–	–	11.8–14.2
Hispanic	%	93.0	1.6	0.8	0.0	0.0	1.1	1.2	1.2	1.1	13.78
(n=103)	CI	85.6–96.7	0.4–6.2	0.1–5.5	–	–	0.2–7.8	0.2–8.4	0.2–8.4	0.2–7.8	11.3–16.3
8th	%	95.9	1.0	0.5	0.8	1.7	0.1	0.0	0.0	0.1	11.92
(n=1,905)	CI	94.9–96.7	0.6–1.6	0.2–0.9	0.5–1.3	1.2–2.4	0.0–0.4	–	–	0.0–0.4	11.6–12.2
10th	%	90.1	1.7	0.7	0.8	0.4	2.1	3.7	0.5	0.1	13.39
(n=1,304)	CI	88.3–91.6	1.1–2.5	0.4–1.3	0.5–1.5	0.2–0.9	1.5–3.1	2.8–4.8	0.2–1.0	0.0–0.5	13.0–13.7
12th	%	88.1	0.6	0.3	0.4	0.6	1.4	2.4	3.6	2.7	15.09
(n=1,128)	CI	86.1–89.8	0.3–1.3	0.1–0.8	0.1–0.9	0.3–1.3	0.9–2.3	1.6–3.5	2.7–4.9	1.9–3.8	14.8–15.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.11. Age of First Alcohol Use

How old were you when you first had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	63.2	4.1	2.1	3.3	4.8	5.9	7.4	5.6	3.5	13.87
(n=4,317)	CI	61.7–64.7	3.5–4.7	1.7–2.6	2.8–3.9	4.2–5.5	5.2–6.7	6.6–8.3	4.9–6.4	3.0–4.2	13.8–14.0
Male	%	63.3	4.9	2.4	3.5	4.7	5.8	6.5	5.1	3.8	13.72
(n=2,094)	CI	61.1–65.4	4.0–5.9	1.8–3.1	2.8–4.4	3.8–5.7	4.8–7.0	5.5–7.8	4.2–6.2	3.0–4.8	13.6–13.9
Female	%	63.8	3.0	1.6	2.9	5.0	5.9	8.4	6.1	3.4	14.09
(n=2,102)	CI	61.6–65.9	2.3–3.8	1.1–2.2	2.2–3.7	4.1–6.0	4.9–7.0	7.2–9.8	5.1–7.3	2.7–4.4	13.9–14.2
White	%	62.5	3.7	1.8	3.5	4.8	6.4	8.0	5.8	3.5	13.95
(n=3,510)	CI	60.8–64.1	3.1–4.3	1.4–2.3	2.9–4.2	4.1–5.6	5.6–7.3	7.1–9.0	5.1–6.7	2.9–4.2	13.8–14.1
Black	%	63.2	8.2	6.8	1.8	5.7	1.5	3.6	5.5	3.7	12.99
(n=131)	CI	54.2–71.3	4.4–14.8	3.6–12.5	0.4–7.0	2.7–11.7	0.4–5.9	1.3–9.3	2.5–11.7	1.4–9.6	12.2–13.8
Hispanic	%	67.7	5.5	5.2	2.0	6.1	3.9	3.5	2.4	3.6	13.11
(n=102)	CI	57.5–76.5	2.4–12.1	2.1–12.2	0.5–8.1	2.7–13.3	1.4–10.4	1.1–10.5	0.6–9.2	1.1–10.7	12.2–14.0
8th	%	83.1	4.4	2.6	3.3	5.0	1.6	0.1	0.0	0.0	11.83
(n=1,896)	CI	81.3–84.7	3.5–5.4	2.0–3.4	2.6–4.2	4.1–6.1	1.2–2.3	0.0–0.4	–	–	11.7–12.0
10th	%	59.2	4.4	2.0	3.5	6.1	9.6	11.2	3.9	0.2	13.58
(n=1,297)	CI	56.5–61.9	3.4–5.6	1.4–2.9	2.6–4.6	4.9–7.5	8.1–11.3	9.6–13.0	3.0–5.1	0.0–0.6	13.4–13.7
12th	%	46.2	3.5	1.7	3.2	3.4	6.6	11.3	13.3	11.0	14.78
(n=1,124)	CI	43.3–49.1	2.5–4.7	1.1–2.6	2.3–4.4	2.5–4.6	5.3–8.2	9.5–13.2	11.4–15.4	9.3–13.0	14.6–14.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.12. Age of First Cigarette Use

		How old were you when you first <u>smoked a cigarette, even just a puff?</u>									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	83.3	2.5	1.2	1.6	2.1	2.4	2.8	2.1	2.0	13.65
(n=4,329)	CI	82.1–84.4	2.1–3.1	0.9–1.6	1.2–2.0	1.7–2.6	2.0–3.0	2.3–3.3	1.7–2.7	1.6–2.5	13.5–13.8
Male	%	82.2	2.9	1.6	1.9	2.2	2.2	2.7	2.0	2.3	13.52
(n=2,103)	CI	80.4–83.9	2.2–3.8	1.1–2.2	1.4–2.6	1.6–2.9	1.6–3.0	2.0–3.6	1.5–2.8	1.7–3.2	13.3–13.8
Female	%	85.3	1.7	0.8	1.1	2.1	2.6	2.8	2.0	1.7	13.90
(n=2,105)	CI	83.6–86.8	1.2–2.3	0.5–1.3	0.7–1.6	1.5–2.8	2.0–3.5	2.1–3.7	1.4–2.8	1.2–2.4	13.6–14.1
White	%	82.4	2.6	1.2	1.7	2.3	2.5	2.9	2.4	2.1	13.67
(n=3,521)	CI	81.0–83.7	2.1–3.2	0.9–1.6	1.3–2.2	1.8–2.9	2.0–3.1	2.3–3.5	1.9–3.0	1.6–2.7	13.5–13.9
Black	%	89.9	3.3	1.5	0.9	0.9	0.9	1.7	0.0	0.9	12.46
(n=131)	CI	82.9–94.2	1.2–8.5	0.3–5.9	0.1–6.5	0.1–6.0	0.1–6.5	0.4–6.7	–	0.1–6.5	10.9–14.0
Hispanic	%	83.6	2.0	5.0	0.0	1.2	1.2	4.5	1.2	1.1	13.14
(n=103)	CI	74.4–90.0	0.5–8.1	2.1–11.9	–	0.2–8.4	0.2–8.4	1.7–11.6	0.2–8.4	0.2–7.8	11.8–14.5
8th	%	94.4	1.5	1.2	1.1	1.5	0.3	0.0	0.1	0.0	11.65
(n=1,905)	CI	93.3–95.4	1.0–2.1	0.8–1.8	0.7–1.7	1.0–2.1	0.1–0.6	–	0.0–0.4	–	11.4–11.9
10th	%	81.5	3.5	1.6	2.1	3.0	3.6	3.8	0.8	0.2	12.90
(n=1,298)	CI	79.3–83.5	2.7–4.7	1.1–2.5	1.4–3.0	2.2–4.1	2.7–4.8	2.9–4.9	0.4–1.4	0.0–0.6	12.7–13.1
12th	%	73.2	2.7	0.7	1.5	1.8	3.5	4.7	5.8	6.2	14.65
(n=1,126)	CI	70.5–75.7	1.9–3.8	0.4–1.4	0.9–2.4	1.1–2.7	2.5–4.7	3.6–6.1	4.6–7.4	4.9–7.8	14.4–14.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	69.0	1.0	1.3	2.0	3.8	5.6	7.4	5.9	4.0	14.51
(n=4,309)	CI	67.6–70.4	0.7–1.3	1.0–1.6	1.7–2.5	3.3–4.4	4.9–6.3	6.6–8.3	5.2–6.7	3.4–4.7	14.4–14.6
Male	%	66.5	1.2	1.4	2.8	4.5	6.2	7.6	5.8	4.0	14.36
(n=2,093)	CI	64.4–68.6	0.8–1.8	0.9–2.0	2.2–3.6	3.7–5.5	5.2–7.4	6.5–8.9	4.8–7.0	3.1–5.0	14.2–14.5
Female	%	72.1	0.6	1.0	1.2	3.0	4.8	7.1	6.0	4.2	14.75
(n=2,096)	CI	70.0–74.0	0.3–1.1	0.6–1.5	0.8–1.7	2.4–3.8	3.9–5.9	6.0–8.4	5.0–7.3	3.3–5.2	14.6–14.9
White	%	68.0	0.9	1.0	2.1	3.7	6.0	7.8	6.2	4.1	14.56
(n=3,505)	CI	66.4–69.6	0.6–1.3	0.7–1.4	1.7–2.7	3.2–4.4	5.2–6.8	6.9–8.8	5.4–7.1	3.5–4.9	14.5–14.7
Black	%	66.7	3.3	2.0	3.6	3.2	6.3	5.0	6.3	3.7	13.97
(n=132)	CI	57.8–74.5	1.2–8.6	0.6–6.3	1.4–8.5	1.2–8.3	3.1–12.3	2.2–10.9	3.0–12.6	1.4–9.5	13.3–14.7
Hispanic	%	77.5	0.8	2.7	0.8	4.0	2.3	6.0	2.4	3.6	14.21
(n=102)	CI	67.7–84.9	0.1–5.6	0.8–8.4	0.1–5.6	1.4–10.4	0.6–8.9	2.5–13.7	0.6–9.2	1.2–10.8	13.3–15.1
8th	%	86.8	1.0	1.3	3.0	5.5	2.3	0.2	0.0	0.0	12.55
(n=1,897)	CI	85.2–88.2	0.6–1.6	0.8–1.9	2.3–3.9	4.5–6.6	1.7–3.1	0.1–0.5	–	–	12.4–12.7
10th	%	63.2	1.0	1.9	1.9	3.5	10.6	14.2	3.5	0.2	14.14
(n=1,292)	CI	60.6–65.8	0.6–1.7	1.2–2.7	1.2–2.7	2.6–4.6	9.1–12.4	12.4–16.2	2.7–4.7	0.1–0.7	14.0–14.3
12th	%	56.0	0.9	0.6	1.2	2.4	3.6	8.1	14.8	12.4	15.47
(n=1,120)	CI	53.1–58.9	0.5–1.6	0.3–1.3	0.7–2.0	1.7–3.5	2.7–4.9	6.6–9.8	12.8–17.0	10.6–14.4	15.3–15.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	79.3	1.1	0.8	1.6	2.4	3.4	5.1	3.7	2.6	14.39
(n=4,334)	CI	78.0–80.5	0.8–1.5	0.5–1.1	1.2–2.0	1.9–2.9	2.9–4.0	4.4–5.9	3.1–4.4	2.1–3.2	14.3–14.5
Male	%	79.2	1.3	0.9	1.9	2.5	3.4	5.1	3.0	2.6	14.23
(n=2,105)	CI	77.3–81.0	0.9–1.9	0.6–1.5	1.3–2.5	1.9–3.3	2.7–4.3	4.2–6.3	2.3–4.0	2.0–3.5	14.0–14.4
Female	%	80.4	0.7	0.5	1.0	2.1	3.3	4.9	4.4	2.6	14.66
(n=2,107)	CI	78.5–82.1	0.4–1.2	0.3–1.0	0.7–1.6	1.6–2.9	2.6–4.3	4.0–6.0	3.5–5.4	2.0–3.5	14.5–14.8
White	%	78.9	0.9	0.8	1.5	2.4	3.6	5.2	3.9	2.7	14.46
(n=3,523)	CI	77.5–80.3	0.6–1.3	0.5–1.1	1.2–2.0	1.9–3.0	3.0–4.4	4.5–6.1	3.3–4.7	2.2–3.3	14.3–14.6
Black	%	76.3	3.6	0.0	3.8	1.4	1.5	6.1	4.5	2.8	14.00
(n=132)	CI	67.7–83.1	1.3–9.2	–	1.6–9.1	0.3–5.8	0.4–6.1	2.9–12.4	1.9–10.4	0.9–8.3	13.1–14.9
Hispanic	%	79.7	0.8	1.9	2.8	4.4	4.6	1.1	2.4	2.4	13.71
(n=103)	CI	70.2–86.8	0.1–5.5	0.5–7.7	0.9–8.7	1.6–11.3	1.7–11.8	0.2–7.8	0.6–9.1	0.6–9.1	12.8–14.7
8th	%	95.6	0.6	0.4	1.1	1.7	0.5	0.1	0.0	0.0	12.28
(n=1,906)	CI	94.6–96.5	0.4–1.1	0.2–0.8	0.7–1.7	1.2–2.4	0.2–0.9	0.0–0.4	–	–	12.0–12.5
10th	%	76.9	1.8	0.8	2.2	3.0	5.4	8.0	1.8	0.2	13.80
(n=1,302)	CI	74.5–79.1	1.2–2.6	0.4–1.4	1.5–3.2	2.2–4.1	4.3–6.7	6.7–9.6	1.2–2.7	0.0–0.6	13.6–14.0
12th	%	64.2	1.0	1.2	1.4	2.4	4.5	7.4	9.7	8.1	15.07
(n=1,126)	CI	61.4–67.0	0.5–1.8	0.7–2.1	0.9–2.3	1.6–3.5	3.4–5.9	6.0–9.1	8.1–11.6	6.6–9.8	14.9–15.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.15. Past Month Prescription Drug Misuse

		During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	96.5	2.0	0.5	0.2	0.4	0.1	0.2	3.5
(n=4,345)	CI	95.9–97.1	1.6–2.5	0.3–0.8	0.1–0.5	0.3–0.7	0.0–0.3	0.1–0.4	2.9–4.1
Male	%	96.8	1.8	0.3	0.2	0.5	0.2	0.2	3.2
(n=2,112)	CI	95.9–97.5	1.3–2.5	0.1–0.7	0.1–0.5	0.2–0.9	0.1–0.6	0.1–0.5	2.5–4.1
Female	%	97.0	2.1	0.4	0.2	0.3	0.0	0.1	3.0
(n=2,105)	CI	96.1–97.6	1.5–2.9	0.2–0.8	0.1–0.6	0.1–0.6	–	0.0–0.4	2.4–3.9
White	%	96.8	2.0	0.4	0.2	0.3	0.1	0.1	3.2
(n=3,529)	CI	96.2–97.4	1.6–2.5	0.2–0.7	0.1–0.5	0.2–0.6	0.0–0.3	0.0–0.3	2.6–3.8
Black	%	94.7	0.9	0.9	0.9	0.9	0.0	1.7	5.3
(n=132)	CI	88.5–97.6	0.1–6.0	0.1–6.4	0.1–6.4	0.1–6.4	–	0.4–6.7	2.4–11.5
Hispanic	%	94.3	1.2	1.1	0.0	2.3	0.0	1.1	5.7
(n=105)	CI	86.9–97.6	0.2–8.2	0.2–7.6	–	0.6–8.9	–	0.2–7.6	2.4–13.1
8th	%	98.5	0.9	0.2	0.1	0.1	0.1	0.2	1.5
(n=1,907)	CI	97.8–98.9	0.6–1.4	0.1–0.6	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.5	1.1–2.2
10th	%	95.4	3.0	0.3	0.2	0.7	0.2	0.4	4.6
(n=1,308)	CI	94.1–96.4	2.2–4.0	0.1–0.8	0.0–0.6	0.4–1.3	0.0–0.6	0.2–0.9	3.6–5.9
12th	%	95.7	2.2	1.0	0.5	0.4	0.1	0.1	4.3
(n=1,130)	CI	94.3–96.7	1.5–3.2	0.5–1.7	0.2–1.2	0.2–1.1	0.0–0.6	0.0–0.6	3.3–5.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.16. Past Month Alcohol Use

		During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	82.3	9.2	4.8	1.5	1.3	0.4	0.5	17.7
(n=4,332)	CI	81.1–83.5	8.3–10.1	4.1–5.5	1.2–1.9	1.0–1.7	0.3–0.7	0.4–0.8	16.5–18.9
Male	%	81.6	8.4	5.1	1.8	1.7	0.5	0.8	18.4
(n=2,105)	CI	79.8–83.3	7.3–9.8	4.2–6.2	1.3–2.6	1.2–2.4	0.3–1.0	0.4–1.3	16.7–20.2
Female	%	84.1	9.5	4.3	1.0	0.6	0.2	0.3	15.9
(n=2,101)	CI	82.4–85.7	8.3–10.9	3.4–5.3	0.6–1.6	0.3–1.1	0.1–0.5	0.1–0.7	14.3–17.6
White	%	82.2	9.2	5.1	1.4	1.3	0.4	0.5	17.8
(n=3,515)	CI	80.9–83.5	8.3–10.3	4.3–5.9	1.0–1.8	0.9–1.7	0.2–0.7	0.3–0.8	16.5–19.1
Black	%	77.3	10.3	3.7	4.4	1.8	1.8	0.8	22.7
(n=133)	CI	68.7–84.0	5.9–17.4	1.4–9.4	1.8–10.2	0.4–6.9	0.4–6.9	0.1–5.9	16.0–31.3
Hispanic	%	84.5	5.8	5.0	0.0	2.4	1.2	1.1	15.5
(n=105)	CI	75.6–90.6	2.5–12.6	2.0–11.8	–	0.6–9.2	0.2–8.2	0.2–7.6	9.4–24.4
8th	%	94.3	4.0	1.0	0.2	0.3	0.1	0.2	5.7
(n=1,905)	CI	93.1–95.2	3.2–5.0	0.6–1.6	0.1–0.6	0.1–0.6	0.0–0.4	0.1–0.5	4.8–6.9
10th	%	80.0	11.1	5.4	1.4	1.2	0.5	0.4	20.0
(n=1,299)	CI	77.7–82.1	9.5–12.9	4.3–6.7	0.9–2.2	0.8–2.0	0.3–1.1	0.2–0.9	17.9–22.3
12th	%	72.0	12.6	8.2	3.0	2.4	0.6	1.1	28.0
(n=1,128)	CI	69.3–74.5	10.8–14.7	6.7–10.0	2.2–4.2	1.6–3.5	0.3–1.3	0.7–2.0	25.5–30.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.17. Past Month Binge Drinking

		During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	90.2	4.7	2.3	1.2	1.0	0.2	0.4	9.8
(n=4,337)	CI	89.2–91.1	4.1–5.5	1.9–2.8	0.9–1.6	0.7–1.4	0.1–0.4	0.2–0.6	8.9–10.8
Male	%	88.3	5.0	2.8	1.6	1.6	0.2	0.5	11.7
(n=2,103)	CI	86.7–89.7	4.1–6.1	2.1–3.7	1.1–2.3	1.1–2.3	0.1–0.6	0.3–0.9	10.3–13.3
Female	%	93.1	4.3	1.7	0.4	0.3	0.1	0.2	6.9
(n=2,106)	CI	91.9–94.2	3.4–5.3	1.2–2.4	0.2–0.9	0.1–0.6	0.0–0.4	0.1–0.5	5.8–8.1
White	%	90.1	5.1	2.4	1.0	1.0	0.2	0.3	9.9
(n=3,520)	CI	89.0–91.1	4.4–6.0	1.9–3.0	0.7–1.4	0.7–1.4	0.1–0.4	0.2–0.6	8.9–11.0
Black	%	84.2	4.5	3.7	3.5	2.4	0.0	1.7	15.8
(n=133)	CI	76.3–89.9	1.9–10.5	1.4–9.4	1.3–9.0	0.8–7.5	–	0.4–6.6	10.1–23.7
Hispanic	%	89.9	4.3	1.2	2.3	1.2	0.0	1.1	10.1
(n=105)	CI	81.5–94.7	1.6–11.3	0.2–8.2	0.6–8.9	0.2–8.2	–	0.2–7.6	5.3–18.5
8th	%	98.1	0.7	0.6	0.2	0.2	0.1	0.1	1.9
(n=1,904)	CI	97.4–98.6	0.4–1.2	0.3–1.0	0.1–0.6	0.1–0.6	0.0–0.4	0.0–0.4	1.4–2.6
10th	%	89.4	5.0	2.8	1.2	1.0	0.1	0.4	10.6
(n=1,303)	CI	87.7–91.0	4.0–6.4	2.1–3.9	0.8–2.0	0.6–1.7	0.0–0.5	0.2–0.9	9.0–12.3
12th	%	82.6	8.7	3.6	2.1	1.9	0.4	0.7	17.4
(n=1,130)	CI	80.3–84.7	7.2–10.5	2.7–4.9	1.4–3.1	1.2–2.8	0.1–0.9	0.4–1.4	15.3–19.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.18. Past Month Cigarette Use

		During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	93.1	3.2	1.4	0.4	0.6	0.5	0.8	6.9
(n=4,340)	CI	92.3–93.9	2.7–3.8	1.0–1.8	0.3–0.7	0.4–0.9	0.3–0.7	0.6–1.2	6.1–7.7
Male	%	92.0	3.4	1.6	0.3	1.0	0.7	1.1	8.0
(n=2,106)	CI	90.7–93.1	2.6–4.3	1.1–2.2	0.2–0.8	0.6–1.5	0.4–1.2	0.7–1.7	6.9–9.3
Female	%	95.1	2.7	0.9	0.4	0.2	0.3	0.4	4.9
(n=2,107)	CI	94.0–96.0	2.1–3.6	0.6–1.5	0.2–0.7	0.1–0.5	0.1–0.7	0.2–0.9	4.0–6.0
White	%	93.0	3.4	1.3	0.4	0.7	0.5	0.8	7.0
(n=3,522)	CI	92.0–93.8	2.8–4.1	0.9–1.7	0.3–0.8	0.4–1.0	0.3–0.9	0.5–1.1	6.2–8.0
Black	%	92.4	1.5	4.4	0.0	0.8	0.0	0.8	7.6
(n=133)	CI	86.0–96.1	0.4–6.1	1.8–10.2	–	0.1–5.9	–	0.1–5.9	3.9–14.0
Hispanic	%	91.1	4.3	1.1	0.0	1.2	0.0	2.2	8.9
(n=105)	CI	83.1–95.6	1.6–11.3	0.2–7.6	–	0.2–8.2	–	0.5–8.6	4.4–16.9
8th	%	97.7	1.2	0.6	0.2	0.2	0.0	0.2	2.3
(n=1,909)	CI	96.9–98.3	0.8–1.8	0.4–1.1	0.1–0.5	0.1–0.5	–	0.1–0.5	1.7–3.1
10th	%	92.8	2.8	1.5	0.3	0.9	0.5	1.1	7.2
(n=1,303)	CI	91.3–94.1	2.1–3.9	0.9–2.3	0.1–0.8	0.5–1.6	0.3–1.1	0.7–1.9	5.9–8.7
12th	%	88.6	5.7	2.0	0.9	0.8	0.9	1.1	11.4
(n=1,128)	CI	86.6–90.3	4.5–7.2	1.4–3.0	0.5–1.6	0.4–1.5	0.5–1.6	0.7–2.0	9.7–13.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	78.6	5.1	3.2	1.8	2.6	1.9	6.8	21.4
(n=4,323)	CI	77.3–79.9	4.4–5.8	2.7–3.8	1.4–2.2	2.1–3.1	1.5–2.4	6.0–7.7	20.1–22.7
Male	%	77.0	4.7	3.2	1.3	2.3	2.9	8.5	23.0
(n=2,096)	CI	75.1–78.9	3.9–5.7	2.4–4.1	0.9–1.9	1.7–3.1	2.3–3.8	7.3–9.9	21.1–24.9
Female	%	81.2	5.3	3.2	1.9	2.7	1.0	4.7	18.8
(n=2,102)	CI	79.4–82.9	4.4–6.4	2.5–4.1	1.4–2.7	2.0–3.5	0.6–1.5	3.8–5.8	17.1–20.6
White	%	77.9	5.3	3.2	1.7	2.8	2.1	7.0	22.1
(n=3,507)	CI	76.4–79.3	4.6–6.2	2.6–3.9	1.3–2.2	2.3–3.5	1.6–2.6	6.1–8.0	20.7–23.6
Black	%	74.3	7.1	2.4	2.1	0.9	0.9	12.3	25.7
(n=132)	CI	65.7–81.4	3.7–13.3	0.7–7.3	0.7–6.6	0.1–6.4	0.1–6.4	7.4–19.7	18.6–34.3
Hispanic	%	79.3	3.4	5.0	1.1	2.3	1.2	7.7	20.7
(n=105)	CI	69.7–86.5	1.1–10.2	2.0–11.8	0.2–7.6	0.6–8.9	0.2–8.2	3.6–15.4	13.5–30.3
8th	%	92.2	2.9	1.4	0.8	1.2	0.4	1.0	7.8
(n=1,906)	CI	90.9–93.4	2.3–3.8	1.0–2.1	0.5–1.4	0.8–1.7	0.2–0.8	0.7–1.6	6.6–9.1
10th	%	74.2	7.1	3.8	1.7	3.0	2.7	7.5	25.8
(n=1,295)	CI	71.8–76.5	5.8–8.6	2.9–5.0	1.1–2.6	2.2–4.1	1.9–3.7	6.2–9.1	23.5–28.2
12th	%	68.7	5.2	4.5	2.8	3.7	2.8	12.3	31.3
(n=1,122)	CI	65.9–71.3	4.1–6.7	3.5–5.9	2.0–4.0	2.8–5.0	1.9–3.9	10.5–14.3	28.7–34.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	95.0	1.3	0.8	0.3	0.6	0.5	1.4	5.0
(n=4,342)	CI	94.2–95.6	1.0–1.7	0.6–1.2	0.1–0.5	0.4–0.9	0.4–0.8	1.1–1.9	4.4–5.8
Male	%	91.8	1.8	1.5	0.4	1.0	1.0	2.6	8.2
(n=2,106)	CI	90.5–93.0	1.3–2.5	1.0–2.1	0.2–0.9	0.6–1.6	0.6–1.6	1.9–3.4	7.0–9.5
Female	%	99.0	0.6	0.2	0.1	0.1	0.0	0.2	1.0
(n=2,108)	CI	98.4–99.3	0.3–1.1	0.1–0.5	0.0–0.4	0.0–0.4	–	0.1–0.5	0.7–1.6
White	%	94.9	1.3	0.9	0.2	0.6	0.6	1.5	5.1
(n=3,526)	CI	94.1–95.6	0.9–1.7	0.6–1.3	0.1–0.5	0.4–0.9	0.4–0.9	1.1–2.0	4.4–5.9
Black	%	95.6	1.8	0.0	0.9	0.0	0.9	0.9	4.4
(n=132)	CI	89.7–98.2	0.4–7.0	–	0.1–6.0	–	0.1–6.4	0.1–6.0	1.8–10.3
Hispanic	%	88.9	6.6	1.1	0.0	1.1	0.0	2.2	11.1
(n=105)	CI	80.5–94.0	3.0–14.2	0.2–7.6	–	0.2–7.6	–	0.5–8.6	6.0–19.5
8th	%	98.4	0.7	0.2	0.2	0.1	0.3	0.3	1.6
(n=1,910)	CI	97.7–98.9	0.4–1.2	0.1–0.5	0.1–0.5	0.0–0.4	0.1–0.6	0.1–0.6	1.1–2.3
10th	%	93.9	1.6	0.9	0.4	0.6	0.8	1.8	6.1
(n=1,304)	CI	92.5–95.1	1.0–2.4	0.5–1.6	0.2–0.9	0.3–1.2	0.4–1.4	1.2–2.7	4.9–7.5
12th	%	92.5	1.7	1.5	0.3	1.1	0.6	2.3	7.5
(n=1,128)	CI	90.8–93.9	1.1–2.6	0.9–2.4	0.1–0.8	0.7–2.0	0.3–1.3	1.6–3.4	6.1–9.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	88.1	4.1	2.2	1.1	1.4	1.1	1.9	11.9
(n=4,340)	CI	87.0–89.1	3.5–4.8	1.8–2.8	0.8–1.5	1.1–1.9	0.8–1.5	1.5–2.4	10.9–13.0
Male	%	87.7	3.6	2.4	0.9	1.4	1.5	2.5	12.3
(n=2,106)	CI	86.1–89.1	2.8–4.5	1.8–3.2	0.6–1.4	1.0–2.1	1.1–2.2	1.9–3.3	10.9–13.9
Female	%	89.6	4.4	1.9	1.2	1.2	0.6	1.1	10.4
(n=2,107)	CI	88.1–90.9	3.5–5.4	1.4–2.7	0.8–1.8	0.8–1.8	0.3–1.1	0.7–1.8	9.1–11.9
White	%	88.3	4.0	2.2	1.2	1.4	1.1	1.8	11.7
(n=3,524)	CI	87.2–89.4	3.4–4.8	1.7–2.7	0.8–1.6	1.0–1.8	0.8–1.6	1.4–2.3	10.6–12.8
Black	%	78.4	8.1	2.4	0.9	4.1	2.6	3.6	21.6
(n=132)	CI	69.9–85.0	4.3–14.9	0.7–7.3	0.1–6.0	1.7–9.6	0.8–8.0	1.3–9.2	15.0–30.1
Hispanic	%	85.8	2.0	3.1	0.0	1.1	4.6	3.5	14.2
(n=105)	CI	76.9–91.6	0.5–7.9	1.0–9.4	–	0.2–7.6	1.7–11.8	1.1–10.4	8.4–23.1
8th	%	97.6	1.1	0.4	0.3	0.2	0.1	0.3	2.4
(n=1,909)	CI	96.8–98.2	0.7–1.7	0.2–0.8	0.1–0.7	0.1–0.6	0.0–0.4	0.1–0.6	1.8–3.2
10th	%	86.1	4.9	2.5	1.1	1.8	1.6	2.0	13.9
(n=1,302)	CI	84.2–87.9	3.9–6.2	1.7–3.4	0.7–1.9	1.2–2.6	1.1–2.5	1.4–2.9	12.1–15.8
12th	%	79.9	6.4	4.0	1.9	2.4	1.8	3.7	20.1
(n=1,129)	CI	77.4–82.1	5.2–8.0	3.0–5.3	1.2–2.8	1.6–3.5	1.1–2.7	2.7–5.0	17.9–22.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.0	0.4	0.1	0.1	0.1	0.0	0.3	1.0
(n=4,350)	CI	98.6–99.2	0.2–0.6	0.1–0.3	0.0–0.2	0.1–0.3	0.0–0.1	0.2–0.5	0.8–1.4
Male	%	99.0	0.4	0.1	0.1	0.2	0.0	0.3	1.0
(n=2,113)	CI	98.4–99.3	0.2–0.8	0.0–0.4	0.0–0.4	0.1–0.5	0.0–0.3	0.1–0.7	0.7–1.6
Female	%	99.7	0.2	0.0	0.0	0.0	0.0	0.1	0.3
(n=2,109)	CI	99.3–99.9	0.1–0.6	–	–	–	–	0.0–0.4	0.1–0.7
White	%	99.2	0.4	0.1	0.1	0.1	0.0	0.2	0.8
(n=3,532)	CI	98.8–99.4	0.2–0.7	0.0–0.3	0.0–0.2	0.0–0.3	0.0–0.2	0.1–0.4	0.6–1.2
Black	%	95.6	0.0	0.8	0.8	0.0	0.0	2.7	4.4
(n=133)	CI	89.8–98.2	–	0.1–5.9	0.1–5.9	–	–	0.9–8.1	1.8–10.2
Hispanic	%	96.6	1.2	1.1	0.0	0.0	0.0	1.1	3.4
(n=105)	CI	89.8–98.9	0.2–8.2	0.2–7.6	–	–	–	0.2–7.6	1.1–10.2
8th	%	99.6	0.0	0.1	0.1	0.0	0.1	0.2	0.4
(n=1,910)	CI	99.2–99.8	–	0.0–0.4	0.0–0.4	–	0.0–0.4	0.1–0.5	0.2–0.8
10th	%	99.1	0.3	0.2	0.1	0.1	0.0	0.2	0.9
(n=1,308)	CI	98.4–99.5	0.1–0.8	0.1–0.7	0.0–0.5	0.0–0.5	–	0.1–0.7	0.5–1.6
12th	%	98.2	0.8	0.1	0.1	0.4	0.0	0.5	1.8
(n=1,132)	CI	97.2–98.8	0.4–1.5	0.0–0.6	0.0–0.6	0.1–0.9	–	0.2–1.2	1.2–2.8

Note: The confidence interval (CI) is 95 percent.

Table BH5.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.0	0.4	0.1	0.0	0.1	0.1	0.3	1.0
(n=4,348)	CI	98.6–99.3	0.2–0.6	0.1–0.3	0.0–0.1	0.0–0.3	0.0–0.2	0.2–0.5	0.7–1.4
Male	%	99.1	0.4	0.1	0.0	0.1	0.0	0.3	0.9
(n=2,111)	CI	98.6–99.5	0.2–0.7	0.0–0.4	–	0.0–0.4	0.0–0.3	0.1–0.7	0.5–1.4
Female	%	99.7	0.2	0.0	0.0	0.1	0.0	0.1	0.3
(n=2,109)	CI	99.3–99.9	0.1–0.5	–	–	0.0–0.4	–	0.0–0.4	0.1–0.7
White	%	99.3	0.3	0.1	0.0	0.1	0.0	0.2	0.7
(n=3,530)	CI	98.9–99.5	0.2–0.6	0.0–0.3	0.0–0.2	0.0–0.3	0.0–0.2	0.1–0.4	0.5–1.1
Black	%	95.6	0.0	0.0	0.0	1.8	0.9	1.7	4.4
(n=133)	CI	89.7–98.2	–	–	–	0.5–7.1	0.1–6.4	0.4–6.6	1.8–10.3
Hispanic	%	94.3	3.5	1.1	0.0	0.0	0.0	1.1	5.7
(n=105)	CI	86.9–97.6	1.1–10.4	0.2–7.6	–	–	–	0.2–7.6	2.4–13.1
8th	%	99.4	0.1	0.1	0.1	0.0	0.1	0.3	0.6
(n=1,910)	CI	99.0–99.7	0.0–0.4	0.0–0.4	0.0–0.4	–	0.0–0.4	0.1–0.6	0.3–1.0
10th	%	99.0	0.5	0.2	0.0	0.1	0.0	0.3	1.0
(n=1,308)	CI	98.3–99.4	0.2–1.0	0.0–0.6	–	0.0–0.5	–	0.1–0.8	0.6–1.7
12th	%	98.5	0.6	0.2	0.0	0.3	0.2	0.3	1.5
(n=1,130)	CI	97.6–99.1	0.3–1.3	0.0–0.7	–	0.1–0.8	0.0–0.7	0.1–0.8	0.9–2.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.4	0.8	0.3	0.1	0.2	0.1	0.2	1.6
(n=4,349)	CI	97.9–98.7	0.6–1.2	0.1–0.5	0.0–0.3	0.1–0.3	0.0–0.2	0.1–0.5	1.3–2.1
Male	%	97.9	1.1	0.4	0.1	0.2	0.1	0.3	2.1
(n=2,112)	CI	97.1–98.4	0.7–1.7	0.2–0.9	0.0–0.4	0.1–0.5	0.0–0.4	0.1–0.7	1.6–2.9
Female	%	99.7	0.2	0.1	0.0	0.0	0.0	0.0	0.3
(n=2,109)	CI	99.3–99.9	0.1–0.6	0.0–0.4	–	–	–	–	0.1–0.7
White	%	98.5	0.8	0.2	0.1	0.1	0.1	0.1	1.5
(n=3,531)	CI	98.1–98.9	0.6–1.2	0.1–0.5	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.3	1.1–1.9
Black	%	94.7	0.9	0.9	0.8	0.9	0.0	1.7	5.3
(n=133)	CI	88.6–97.6	0.1–6.4	0.1–6.4	0.1–5.9	0.1–6.4	–	0.4–6.6	2.4–11.4
Hispanic	%	95.4	3.5	0.0	0.0	0.0	0.0	1.1	4.6
(n=105)	CI	88.2–98.3	1.1–10.4	–	–	–	–	0.2–7.6	1.7–11.8
8th	%	99.5	0.1	0.1	0.1	0.1	0.1	0.2	0.5
(n=1,910)	CI	99.1–99.8	0.0–0.4	0.0–0.4	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.5	0.2–0.9
10th	%	98.6	0.5	0.2	0.2	0.2	0.0	0.2	1.4
(n=1,308)	CI	97.8–99.1	0.3–1.1	0.0–0.6	0.1–0.7	0.1–0.7	–	0.1–0.7	0.9–2.2
12th	%	96.8	1.9	0.5	0.0	0.2	0.2	0.4	3.2
(n=1,131)	CI	95.6–97.7	1.3–2.9	0.2–1.2	–	0.0–0.7	0.0–0.7	0.1–0.9	2.3–4.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.2	0.1	0.0	0.1	0.0	0.3	0.8
(n=4,350)	CI	98.9–99.4	0.1–0.4	0.1–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.2–0.5	0.6–1.1
Male	%	99.2	0.4	0.1	0.0	0.1	0.0	0.3	0.8
(n=2,113)	CI	98.6–99.5	0.2–0.7	0.0–0.5	–	0.0–0.4	0.0–0.3	0.1–0.7	0.5–1.4
Female	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=2,109)	CI	–	–	–	–	–	–	–	–
White	%	99.5	0.2	0.1	0.0	0.1	0.0	0.1	0.5
(n=3,532)	CI	99.2–99.7	0.1–0.4	0.0–0.3	0.0–0.2	0.0–0.2	0.0–0.2	0.1–0.4	0.3–0.8
Black	%	95.6	0.0	0.0	0.0	0.8	0.9	2.6	4.4
(n=133)	CI	89.8–98.2	–	–	–	0.1–5.9	0.1–6.4	0.8–7.9	1.8–10.2
Hispanic	%	96.6	1.2	1.1	0.0	0.0	0.0	1.1	3.4
(n=105)	CI	89.8–98.9	0.2–8.2	0.2–7.6	–	–	–	0.2–7.6	1.1–10.2
8th	%	99.5	0.1	0.0	0.1	0.1	0.1	0.2	0.5
(n=1,910)	CI	99.1–99.8	0.0–0.4	–	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.6	0.2–0.9
10th	%	99.4	0.2	0.1	0.0	0.1	0.0	0.2	0.6
(n=1,308)	CI	98.8–99.7	0.1–0.7	0.0–0.5	–	0.0–0.5	–	0.1–0.7	0.3–1.2
12th	%	98.7	0.3	0.4	0.1	0.1	0.1	0.4	1.3
(n=1,132)	CI	97.8–99.2	0.1–0.8	0.1–0.9	0.0–0.6	0.0–0.6	0.0–0.6	0.2–1.1	0.8–2.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.2	0.0	0.2	0.0	0.0	0.4	0.8
(n=4,349)	CI	98.9–99.4	0.1–0.4	0.0–0.2	0.1–0.3	0.0–0.2	0.0–0.2	0.2–0.6	0.6–1.1
Male	%	99.2	0.2	0.0	0.2	0.1	0.0	0.3	0.8
(n=2,113)	CI	98.7–99.5	0.1–0.5	–	0.1–0.5	0.0–0.4	0.0–0.3	0.2–0.7	0.5–1.3
Female	%	99.9	0.0	0.1	0.0	0.0	0.0	0.0	0.1
(n=2,109)	CI	99.6–100.0	–	0.0–0.4	–	–	–	–	0.0–0.4
White	%	99.5	0.1	0.1	0.1	0.0	0.0	0.2	0.5
(n=3,531)	CI	99.1–99.7	0.0–0.3	0.0–0.2	0.0–0.3	0.0–0.2	0.0–0.2	0.1–0.5	0.3–0.9
Black	%	96.5	0.8	0.0	0.0	0.0	0.9	1.8	3.5
(n=133)	CI	90.9–98.7	0.1–5.9	–	–	–	0.1–6.4	0.4–6.9	1.3–9.1
Hispanic	%	95.5	1.2	0.0	1.1	0.0	0.0	2.2	4.5
(n=105)	CI	88.4–98.3	0.2–8.2	–	0.2–7.6	–	–	0.5–8.6	1.7–11.6
8th	%	99.5	0.2	0.1	0.1	0.0	0.1	0.2	0.5
(n=1,909)	CI	99.1–99.8	0.1–0.5	0.0–0.4	0.0–0.4	–	0.0–0.4	0.1–0.5	0.2–0.9
10th	%	99.2	0.2	0.1	0.2	0.0	0.0	0.3	0.8
(n=1,308)	CI	98.6–99.6	0.0–0.6	0.0–0.5	0.1–0.7	–	–	0.1–0.8	0.4–1.4
12th	%	98.8	0.3	0.0	0.2	0.1	0.1	0.6	1.2
(n=1,132)	CI	97.9–99.3	0.1–0.8	–	0.0–0.7	0.0–0.6	0.0–0.6	0.3–1.3	0.7–2.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you use ecstasy (MDMA)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	98.9	0.6	0.1	0.1	0.1	0.0	0.3	1.1
(n=4,348)	CI	98.5-99.2	0.4-0.9	0.0-0.2	0.0-0.3	0.0-0.3	0.0-0.1	0.1-0.5	0.8-1.5
Male	%	98.7	0.8	0.1	0.0	0.1	0.0	0.3	1.3
(n=2,112)	CI	98.1-99.1	0.5-1.3	0.0-0.4	0.0-0.3	0.0-0.4	0.0-0.3	0.1-0.7	0.9-1.9
Female	%	99.8	0.1	0.0	0.0	0.1	0.0	0.0	0.2
(n=2,109)	CI	99.5-99.9	0.0-0.5	-	-	0.0-0.4	-	-	0.1-0.5
White	%	99.1	0.6	0.0	0.1	0.1	0.0	0.1	0.9
(n=3,531)	CI	98.7-99.4	0.3-0.9	0.0-0.2	0.0-0.2	0.0-0.3	0.0-0.2	0.1-0.4	0.6-1.3
Black	%	95.6	0.9	0.9	0.0	0.9	0.0	1.8	4.4
(n=132)	CI	89.7-98.2	0.1-6.0	0.1-6.0	-	0.1-6.4	-	0.4-7.0	1.8-10.3
Hispanic	%	96.6	2.3	0.0	0.0	0.0	0.0	1.1	3.4
(n=105)	CI	89.8-98.9	0.6-8.9	-	-	-	-	0.2-7.6	1.1-10.2
8th	%	99.5	0.1	0.0	0.1	0.0	0.1	0.2	0.5
(n=1,910)	CI	99.1-99.8	0.0-0.4	-	0.0-0.4	-	0.0-0.4	0.1-0.6	0.2-0.9
10th	%	99.2	0.5	0.1	0.0	0.1	0.0	0.2	0.8
(n=1,308)	CI	98.5-99.5	0.3-1.1	0.0-0.5	-	0.0-0.5	-	0.0-0.6	0.5-1.5
12th	%	97.9	1.2	0.1	0.2	0.2	0.0	0.4	2.1
(n=1,130)	CI	96.9-98.6	0.7-2.1	0.0-0.6	0.0-0.7	0.0-0.7	-	0.2-1.1	1.4-3.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you use methamphetamines (crank, etc.)?							
		0 days	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	All 30 days	Any Use
Total	%	99.1	0.2	0.2	0.1	0.0	0.1	0.3	0.9
(n=4,349)	CI	98.8-99.4	0.1-0.3	0.1-0.3	0.0-0.3	0.0-0.2	0.0-0.3	0.2-0.6	0.6-1.2
Male	%	99.0	0.2	0.1	0.2	0.0	0.2	0.3	1.0
(n=2,112)	CI	98.4-99.4	0.1-0.5	0.0-0.5	0.1-0.5	-	0.0-0.5	0.2-0.7	0.6-1.6
Female	%	99.9	0.1	0.0	0.0	0.0	0.0	0.0	0.1
(n=2,109)	CI	99.6-100.0	0.0-0.4	-	-	-	-	-	0.0-0.4
White	%	99.4	0.1	0.1	0.1	0.0	0.1	0.2	0.6
(n=3,531)	CI	99.1-99.6	0.0-0.4	0.0-0.3	0.0-0.3	0.0-0.2	0.0-0.3	0.1-0.4	0.4-0.9
Black	%	95.7	0.0	0.8	0.0	0.0	0.0	3.5	4.3
(n=133)	CI	90.0-98.2	-	0.1-5.9	-	-	-	1.3-9.0	1.8-10.0
Hispanic	%	95.8	2.0	1.1	0.0	0.0	0.0	1.1	4.2
(n=105)	CI	89.2-98.5	0.5-7.9	0.2-7.6	-	-	-	0.2-7.6	1.5-10.8
8th	%	99.5	0.1	0.1	0.1	0.0	0.1	0.2	0.5
(n=1,910)	CI	99.0-99.7	0.0-0.4	0.0-0.4	0.0-0.4	-	0.0-0.4	0.1-0.6	0.3-1.0
10th	%	99.1	0.2	0.2	0.2	0.0	0.0	0.3	0.9
(n=1,307)	CI	98.4-99.5	0.0-0.6	0.1-0.7	0.1-0.7	-	-	0.1-0.8	0.5-1.6
12th	%	98.8	0.3	0.2	0.0	0.1	0.2	0.5	1.2
(n=1,132)	CI	97.9-99.3	0.1-0.8	0.0-0.7	-	0.0-0.6	0.0-0.7	0.2-1.2	0.7-2.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	25.5	28.0	19.1	27.5	46.5
(n=1,659)	CI	23.4–27.7	25.9–30.3	17.2–21.1	25.4–29.7	44.1–49.0
Male	%	24.1	27.8	18.6	29.5	48.0
(n=759)	CI	21.1–27.4	24.7–31.2	15.9–21.6	26.3–32.8	44.4–51.7
Female	%	24.9	29.5	19.4	26.2	45.6
(n=827)	CI	22.0–28.1	26.4–32.8	16.8–22.4	23.3–29.3	42.1–49.0
White	%	25.0	27.5	19.3	28.2	47.5
(n=1,357)	CI	22.7–27.4	25.1–29.9	17.3–21.6	25.8–30.7	44.8–50.2
Black	%	34.5	32.7	18.1	14.8	32.8
(n=51)	CI	22.4–49.0	20.9–47.2	9.4–31.9	7.3–27.7	21.0–47.3
Hispanic	%	27.6	33.1	21.4	17.9	39.3
(n=35)	CI	14.5–46.1	19.0–51.1	10.1–39.8	8.2–34.6	23.9–57.2
8th	%	22.3	27.4	15.0	35.3	50.3
(n=660)	CI	19.3–25.6	24.1–31.0	12.5–17.9	31.7–39.0	46.5–54.1
10th	%	27.0	27.7	20.2	25.0	45.2
(n=552)	CI	23.5–30.9	24.2–31.6	17.0–23.7	21.6–28.8	41.1–49.4
12th	%	26.7	28.9	21.6	22.9	44.4
(n=447)	CI	22.8–31.0	24.9–33.3	18.0–25.6	19.2–27.0	39.9–49.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	27.9	41.2	16.5	14.3	30.8
(n=2,482)	CI	26.2–29.8	39.3–43.2	15.1–18.1	13.0–15.7	29.0–32.7
Male	%	27.1	40.6	16.5	15.9	32.4
(n=1,155)	CI	24.5–29.8	37.7–43.5	14.5–18.8	13.9–18.1	29.7–35.2
Female	%	26.9	43.0	17.0	13.1	30.1
(n=1,223)	CI	24.5–29.5	40.2–45.8	15.0–19.2	11.3–15.1	27.6–32.7
White	%	27.3	41.5	16.8	14.4	31.2
(n=2,048)	CI	25.3–29.3	39.4–43.7	15.2–18.5	13.0–16.0	29.2–33.3
Black	%	35.3	41.5	13.0	10.1	23.2
(n=66)	CI	24.5–47.9	29.9–54.2	6.7–23.8	4.7–20.3	14.4–35.1
Hispanic	%	32.9	35.0	13.5	18.6	32.1
(n=54)	CI	21.3–47.1	23.0–49.2	6.4–26.4	10.0–32.0	20.5–46.4
8th	%	25.5	36.4	19.2	18.9	38.1
(n=898)	CI	22.8–28.5	33.3–39.6	16.7–21.9	16.5–21.6	35.0–41.3
10th	%	26.4	42.6	16.3	14.8	31.0
(n=808)	CI	23.4–29.5	39.2–46.0	13.9–19.0	12.5–17.4	27.9–34.3
12th	%	31.3	43.5	14.9	10.4	25.3
(n=776)	CI	28.1–34.6	40.0–47.0	12.5–17.6	8.4–12.7	22.3–28.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)?</u>				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	43.8	28.9	11.5	15.8	27.3
(n=2,238)	CI	41.7–45.9	27.1–30.9	10.3–12.9	14.4–17.3	25.5–29.2
Male	%	46.4	25.8	11.7	16.1	27.8
(n=1,101)	CI	43.4–49.4	23.2–28.5	10.0–13.7	14.1–18.4	25.3–30.5
Female	%	39.5	33.1	11.6	15.8	27.4
(n=1,046)	CI	36.5–42.6	30.2–36.0	9.8–13.6	13.8–18.1	24.8–30.2
White	%	44.5	29.1	11.3	15.2	26.5
(n=1,867)	CI	42.2–46.8	27.0–31.2	9.9–12.7	13.7–16.9	24.5–28.5
Black	%	37.7	39.9	9.0	13.4	22.4
(n=61)	CI	26.1–51.0	28.1–53.0	4.0–19.4	6.8–24.5	13.6–34.7
Hispanic	%	44.1	21.6	12.4	21.8	34.2
(n=51)	CI	30.5–58.7	12.0–35.8	5.8–24.8	12.3–35.5	22.2–48.6
8th	%	24.7	27.5	20.1	27.8	47.8
(n=717)	CI	21.7–28.0	24.3–30.9	17.3–23.2	24.6–31.2	44.2–51.5
10th	%	39.7	32.7	11.2	16.3	27.6
(n=762)	CI	36.3–43.3	29.4–36.1	9.2–13.7	13.9–19.1	24.5–30.9
12th	%	59.0	26.3	6.6	8.1	14.7
(n=759)	CI	55.4–62.4	23.3–29.6	5.0–8.5	6.4–10.3	12.3–17.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)?</u>				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	29.1	29.8	17.7	23.4	41.1
(n=1,770)	CI	27.0–31.3	27.7–32.0	16.0–19.6	21.5–25.3	38.8–43.4
Male	%	29.8	28.2	17.6	24.4	42.0
(n=847)	CI	26.7–33.1	25.2–31.4	15.2–20.4	21.6–27.3	38.7–45.4
Female	%	26.5	32.3	18.5	22.7	41.2
(n=843)	CI	23.6–29.7	29.1–35.6	16.0–21.3	20.0–25.6	37.9–44.6
White	%	28.5	28.7	18.7	24.1	42.8
(n=1,465)	CI	26.2–30.9	26.4–31.1	16.8–20.8	22.1–26.4	40.3–45.4
Black	%	50.5	24.6	10.8	14.1	25.0
(n=54)	CI	36.9–63.9	14.5–38.4	5.0–21.8	7.2–25.9	15.3–38.0
Hispanic	%	33.7	28.4	12.3	25.7	38.0
(n=41)	CI	20.4–50.2	16.0–45.1	4.9–27.4	14.2–41.9	23.9–54.4
8th	%	14.4	13.3	22.7	49.6	72.2
(n=472)	CI	11.5–17.9	10.6–16.7	19.1–26.7	45.1–54.1	68.0–76.1
10th	%	31.5	30.3	16.3	21.9	38.2
(n=663)	CI	28.1–35.1	26.9–33.9	13.7–19.4	18.9–25.2	34.6–42.0
12th	%	33.7	37.1	16.8	12.4	29.2
(n=635)	CI	30.1–37.5	33.5–41.0	14.1–19.9	10.0–15.2	25.7–32.8

Note: The confidence interval (CI) is 95 percent.

Table BH5.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	1.9	98.1
(n=4,342)	CI	1.5–2.4	97.6–98.5
Male	%	1.9	98.1
(n=2,109)	CI	1.4–2.6	97.4–98.6
Female	%	1.3	98.7
(n=2,107)	CI	0.8–1.9	98.1–99.2
White	%	1.7	98.3
(n=3,526)	CI	1.3–2.2	97.8–98.7
Black	%	3.6	96.4
(n=133)	CI	1.3–9.2	90.8–98.7
Hispanic	%	3.4	96.6
(n=105)	CI	1.1–10.2	89.8–98.9
8th	%	0.6	99.4
(n=1,907)	CI	0.4–1.1	98.9–99.6
10th	%	2.4	97.6
(n=1,303)	CI	1.7–3.4	96.6–98.3
12th	%	2.7	97.3
(n=1,132)	CI	1.9–3.9	96.1–98.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	15.6	84.4
(n=4,317)	CI	14.5–16.8	83.2–85.5
Male	%	15.8	84.2
(n=2,091)	CI	14.2–17.6	82.4–85.8
Female	%	14.3	85.7
(n=2,100)	CI	12.8–15.9	84.1–87.2
White	%	15.8	84.2
(n=3,504)	CI	14.5–17.1	82.9–85.5
Black	%	19.7	80.3
(n=133)	CI	13.4–28.1	71.9–86.6
Hispanic	%	13.0	87.0
(n=103)	CI	7.4–21.8	78.2–92.6
8th	%	4.5	95.5
(n=1,904)	CI	3.6–5.5	94.5–96.4
10th	%	17.5	82.5
(n=1,298)	CI	15.5–19.7	80.3–84.5
12th	%	25.7	74.3
(n=1,115)	CI	23.2–28.3	71.7–76.8

Note: The confidence interval (CI) is 95 percent.

Table BH5.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=73)	Male (n=36)	Female (n=24)	White (n=53)	Black (n=4)	Hispanic (n=3)	8th (n=12)	10th (n=31)	12th (n=30)
Got a prescription from one doctor	%	15.2	15.4	4.6	10.5	24.0	35.0	33.3	12.9	12.9
	CI	8.6–25.5	6.7–31.6	0.5–29.3	4.6–22.2	0.5–95.7	0.1–99.7	10.9–67.1	4.6–31.1	4.6–31.1
Got a prescription from more than one doctor	%	14.4	12.7	0.0	12.5	26.0	35.0	25.0	9.7	16.1
	CI	8.0–24.6	5.1–28.6	–	5.9–24.8	0.5–96.1	0.1–99.7	6.7–60.7	2.9–27.4	6.5–34.6
Bought on the Internet	%	18.7	13.6	2.9	11.4	50.0	0.0	25.0	12.9	22.6
	CI	11.2–29.6	5.5–30.0	0.3–20.7	5.0–23.8	2.5–97.5	–	6.7–60.7	4.6–31.1	10.7–41.6
Bought from a friend, relative, or someone you know	%	41.6	40.7	46.9	45.2	51.9	0.0	25.0	38.7	48.4
	CI	30.6–53.5	25.2–58.2	27.0–67.9	31.9–59.2	2.7–97.7	–	6.7–60.7	22.7–57.6	30.8–66.4
Got from a friend, relative, or someone you know for free	%	46.0	40.9	42.7	49.5	50.0	32.5	50.0	35.5	54.8
	CI	34.7–57.8	25.5–58.3	23.6–64.3	35.9–63.2	2.5–97.5	0.1–99.7	21.0–79.0	20.1–54.5	36.5–72.0
Took from a friend, relative, or someone you know without asking	%	23.1	14.1	20.1	21.3	26.0	0.0	50.0	12.9	25.8
	CI	14.7–34.2	5.7–30.9	8.4–40.9	12.1–34.6	0.5–96.1	–	21.0–79.0	4.6–31.1	12.9–44.9
Took from someone's house or workplace without asking	%	10.2	4.8	2.9	7.4	0.0	0.0	25.0	3.2	12.9
	CI	5.0–19.5	1.1–18.9	0.3–20.7	2.7–18.8	–	–	6.7–60.7	0.4–21.6	4.6–31.1
Got it at a party	%	31.3	30.6	20.9	26.9	74.0	0.0	33.3	19.4	41.9
	CI	21.5–43.2	17.2–48.3	8.3–43.5	16.3–41.0	3.9–99.5	–	10.9–67.1	8.5–38.1	25.3–60.6
Got it some other way	%	29.9	41.9	7.2	27.0	0.0	32.5	50.0	29.0	25.8
	CI	20.4–41.6	26.5–59.2	1.6–27.3	16.5–40.9	–	0.1–99.7	21.0–79.0	15.2–48.2	12.9–44.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=598)	Male (n=292)	Female (n=267)	White (n=494)	Black (n=22)	Hispanic (n=12)	8th (n=85)	10th (n=227)	12th (n=286)
I bought it in a store	%	7.2	7.7	4.5	6.0	13.7	9.3	4.7	3.1	10.8
	CI	5.4–9.7	5.1–11.6	2.5–7.8	4.2–8.5	4.1–37.4	0.9–53.3	1.7–12.1	1.5–6.3	7.7–15.0
I bought it in a gas station	%	8.8	9.0	5.6	7.5	18.1	9.3	8.2	3.5	12.9
	CI	6.8–11.5	6.1–13.0	3.4–9.2	5.4–10.2	6.4–41.7	0.9–53.3	3.9–16.5	1.8–6.9	9.5–17.3
I bought it in a restaurant	%	3.7	1.8	4.2	2.7	4.3	18.7	3.5	2.6	4.5
	CI	2.4–5.6	0.8–4.3	2.3–7.5	1.6–4.7	0.5–28.6	3.7–57.8	1.1–10.6	1.2–5.8	2.6–7.7
I bought it in a bar or club	%	4.1	1.5	5.1	3.2	4.3	18.0	3.5	2.2	5.6
	CI	2.7–6.0	0.5–3.8	2.9–8.6	1.9–5.2	0.5–28.6	3.5–56.7	1.1–10.6	0.9–5.2	3.4–8.9
I bought it at a public event	%	4.9	2.5	3.7	2.9	9.0	18.0	4.7	3.5	5.9
	CI	3.4–7.0	1.2–5.2	2.0–6.8	1.7–4.9	2.0–32.5	3.5–56.7	1.7–12.1	1.8–6.9	3.7–9.3
I gave someone money to buy it for me	%	31.3	35.4	26.9	32.4	36.1	36.0	9.4	30.3	36.2
	CI	27.7–35.2	30.0–41.1	21.8–32.7	28.4–36.8	18.0–59.2	12.0–69.9	4.7–17.9	24.6–36.6	30.8–42.0
I got it from a sibling (under 21)	%	5.7	4.4	6.1	5.7	9.0	9.3	8.2	5.7	5.2
	CI	4.1–7.9	2.5–7.4	3.7–9.7	4.0–8.1	2.0–32.5	0.9–53.3	3.9–16.5	3.3–9.6	3.2–8.5
I got it from a friend (under 21)	%	29.5	26.3	32.1	29.9	45.5	18.7	29.4	29.4	29.6
	CI	26.0–33.3	21.5–31.7	26.7–38.1	26.0–34.2	25.1–67.5	3.7–57.8	20.6–40.1	23.8–35.7	24.6–35.2
I got it at a party	%	50.7	51.9	49.1	49.5	73.3	36.0	32.9	44.7	58.5
	CI	46.7–54.8	46.1–57.6	43.1–55.2	45.1–54.0	49.7–88.4	12.0–69.9	23.6–43.8	38.4–51.3	52.7–64.1
I took it from a store or family member	%	11.8	12.2	7.9	9.5	14.1	44.6	22.4	11.8	9.8
	CI	9.5–14.6	8.9–16.4	5.2–11.8	7.2–12.4	4.2–38.0	17.4–75.5	14.6–32.6	8.2–16.8	6.8–13.8
I took it home without permission	%	17.1	14.4	17.2	16.2	18.1	35.2	30.6	23.7	9.8
	CI	14.3–20.3	10.8–18.8	13.2–22.2	13.3–19.7	6.4–41.7	12.3–68.0	21.6–41.4	18.6–29.7	6.8–13.8
I got it from my parents with permission	%	23.8	17.1	30.0	24.9	13.4	8.7	35.3	21.1	23.7
	CI	20.6–27.4	13.2–21.9	24.7–35.8	21.2–28.9	3.9–36.7	0.8–51.2	25.7–46.2	16.2–26.9	19.1–29.0
I got it from an older sibling (over 21)	%	11.8	7.6	15.3	12.0	4.7	9.3	5.9	11.4	13.2
	CI	9.4–14.7	5.0–11.3	11.4–20.3	9.4–15.2	0.6–30.3	0.9–53.3	2.4–13.6	7.9–16.3	9.8–17.7
I got it from an older friend (over 21)	%	22.5	20.4	22.9	21.4	23.1	9.3	11.8	23.7	23.7
	CI	19.3–26.1	16.1–25.5	18.2–28.5	17.9–25.3	9.2–47.0	0.9–53.3	6.4–20.7	18.6–29.7	19.1–29.0
I got it from some other adult	%	13.6	13.4	11.3	12.4	13.4	18.0	12.9	12.3	14.6
	CI	11.0–16.6	9.9–17.8	8.0–15.8	9.7–15.6	3.9–36.7	3.5–56.7	7.2–22.1	8.6–17.3	11.0–19.2
I got it some other way	%	20.7	23.4	15.1	19.3	22.7	46.1	21.2	24.1	18.1
	CI	17.6–24.2	18.9–28.7	11.3–20.0	16.1–23.1	9.1–46.5	18.0–76.8	13.6–31.4	19.0–30.1	14.1–23.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?				
		I bought or tried to buy alcohol	Yes, asked for proof of age	No, not asked for proof of age
Total	%	4.0	44.3	55.7
(n=137)	CI	3.4–4.7	36.0–52.9	47.1–64.0
Male	%	4.7	41.8	58.2
(n=73)	CI	3.7–5.8	30.7–53.8	46.2–69.3
Female	%	2.4	45.5	54.5
(n=42)	CI	1.8–3.3	30.5–61.4	38.6–69.5
White	%	3.8	47.1	52.9
(n=105)	CI	3.1–4.5	37.5–56.9	43.1–62.5
Black	%	4.4	0.0	100.0
(n=5)	CI	1.8–10.3	–	–
Hispanic	%	4.8	27.5	72.5
(n=4)	CI	1.8–12.5	0.5–96.4	3.6–99.5
8th	%	1.9	19.4	80.6
(n=31)	CI	1.3–2.7	8.5–38.1	61.9–91.5
10 th	%	3.6	33.3	66.7
(n=39)	CI	2.7–4.9	19.9–50.1	49.9–80.1
12th	%	6.6	57.4	42.6
(n=67)	CI	5.3–8.3	45.1–68.8	31.2–54.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?					
		I bought or tried to buy alcohol	No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	4.0	21.0	29.5	49.5
(n=137)	CI	3.4–4.7	15.0–28.5	22.4–37.8	41.2–57.9
Male	%	4.7	25.1	28.2	46.6
(n=73)	CI	3.7–5.8	16.5–36.3	19.0–39.8	35.4–58.2
Female	%	2.4	19.5	31.9	48.6
(n=42)	CI	1.8–3.3	10.0–34.6	19.0–48.3	33.3–64.2
White	%	3.8	22.1	29.4	48.4
(n=105)	CI	3.1–4.5	15.2–31.1	21.4–39.0	38.9–58.1
Black	%	4.4	36.0	21.3	42.6
(n=5)	CI	1.8–10.3	3.1–90.8	0.8–89.8	4.1–92.8
Hispanic	%	4.8	0.0	27.5	72.5
(n=4)	CI	1.8–12.5	–	0.5–96.4	3.6–99.5
8th	%	1.9	31.3	12.5	56.3
(n=31)	CI	1.3–2.7	17.1–50.0	4.5–30.2	38.1–72.9
10 th	%	3.6	29.3	41.5	29.3
(n=39)	CI	2.7–4.9	17.0–45.5	27.0–57.5	17.0–45.5
12th	%	6.6	13.2	27.9	58.8
(n=67)	CI	5.3–8.3	6.9–23.9	18.4–40.0	46.5–70.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.39. Ever Ridden with a Prescription Drugs Impaired Driver

		Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		
		No	Yes	Not sure
Total	%	82.1	7.4	10.5
(n=4,123)	CI	80.9–83.3	6.6–8.2	9.6–11.5
Male	%	83.3	6.1	10.5
(n=1,984)	CI	81.6–85.0	5.1–7.3	9.2–12.0
Female	%	81.9	7.9	10.2
(n=2,015)	CI	80.1–83.5	6.8–9.2	8.9–11.6
White	%	82.4	7.2	10.3
(n=3,345)	CI	81.1–83.7	6.4–8.2	9.3–11.4
Black	%	83.8	8.3	8.0
(n=123)	CI	75.9–89.4	4.4–14.9	4.3–14.5
Hispanic	%	85.0	5.2	9.9
(n=100)	CI	76.5–90.8	2.1–12.1	5.4–17.3
8th	%	82.4	5.7	11.9
(n=1,800)	CI	80.6–84.1	4.7–6.8	10.5–13.5
10th	%	81.0	7.9	11.1
(n=1,232)	CI	78.7–83.1	6.5–9.6	9.4–13.0
12th	%	82.9	8.6	8.5
(n=1,091)	CI	80.6–85.0	7.1–10.4	7.0–10.3

Note: The confidence interval (CI) is 95 percent.

Table BH5.40. Frequency Riding with a Prescription Drugs Impaired Driver

		During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?					
		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	96.0	1.1	1.3	0.2	1.4	4.0
(n=4,271)	CI	95.3–96.6	0.9–1.5	0.9–1.7	0.1–0.4	1.1–1.8	3.4–4.7
Male	%	96.2	1.1	1.0	0.2	1.5	3.8
(n=2,073)	CI	95.3–97.0	0.7–1.7	0.6–1.6	0.1–0.6	1.0–2.2	3.0–4.7
Female	%	96.3	1.2	1.4	0.1	0.9	3.7
(n=2,073)	CI	95.4–97.1	0.8–1.8	0.9–2.0	0.0–0.5	0.6–1.4	2.9–4.6
White	%	96.2	1.1	1.2	0.2	1.3	3.8
(n=3,463)	CI	95.5–96.8	0.8–1.5	0.9–1.7	0.1–0.4	1.0–1.7	3.2–4.5
Black	%	94.5	0.9	0.9	0.0	3.6	5.5
(n=130)	CI	88.2–97.5	0.1–6.5	0.1–6.5	–	1.4–9.4	2.5–11.8
Hispanic	%	97.0	1.9	1.1	0.0	0.0	3.0
(n=103)	CI	90.7–99.1	0.5–7.6	0.2–7.8	–	–	0.9–9.3
8th	%	97.7	1.0	0.5	0.1	0.8	2.3
(n=1,880)	CI	96.9–98.3	0.6–1.5	0.2–0.9	0.0–0.4	0.5–1.3	1.7–3.1
10th	%	94.6	1.6	2.0	0.3	1.6	5.4
(n=1,270)	CI	93.2–95.7	1.0–2.4	1.3–2.9	0.1–0.8	1.0–2.4	4.3–6.8
12th	%	95.7	0.9	1.3	0.2	1.9	4.3
(n=1,121)	CI	94.4–96.8	0.5–1.6	0.8–2.2	0.0–0.7	1.2–2.8	3.2–5.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	68.7	20.3	11.0
(n=4,278)	CI	67.3–70.1	19.1–21.6	10.1–12.0
Male	%	70.4	18.4	11.2
(n=2,074)	CI	68.4–72.4	16.7–20.2	9.9–12.7
Female	%	68.1	21.2	10.7
(n=2,078)	CI	66.0–70.1	19.5–23.1	9.4–12.1
White	%	68.9	20.3	10.8
(n=3,466)	CI	67.3–70.4	19.0–21.7	9.8–11.9
Black	%	65.5	23.8	10.7
(n=131)	CI	56.7–73.4	17.0–32.2	6.4–17.2
Hispanic	%	69.1	19.0	11.9
(n=104)	CI	59.3–77.5	12.4–28.0	6.9–19.7
8th	%	73.1	13.5	13.4
(n=1,880)	CI	71.0–75.0	12.0–15.1	11.9–15.0
10th	%	68.1	20.6	11.3
(n=1,279)	CI	65.5–70.6	18.5–22.9	9.7–13.1
12th	%	64.6	27.2	8.1
(n=1,119)	CI	61.8–67.4	24.7–29.9	6.6–9.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	88.6	4.3	3.5	1.0	2.6	11.4
(n=4,279)	CI	87.5–89.5	3.7–5.0	3.0–4.2	0.7–1.4	2.1–3.1	10.5–12.5
Male	%	89.2	3.3	3.3	1.0	3.1	10.8
(n=2,075)	CI	87.7–90.5	2.6–4.2	2.6–4.2	0.7–1.6	2.4–4.1	9.5–12.3
Female	%	88.7	5.1	3.7	1.0	1.4	11.3
(n=2,079)	CI	87.2–90.0	4.2–6.2	3.0–4.7	0.6–1.6	1.0–2.1	10.0–12.8
White	%	88.5	4.3	3.7	1.1	2.5	11.5
(n=3,469)	CI	87.4–89.6	3.6–5.0	3.1–4.4	0.8–1.5	2.0–3.1	10.4–12.6
Black	%	86.9	3.9	2.1	2.7	4.5	13.1
(n=131)	CI	79.5–91.9	1.6–9.1	0.6–6.4	0.9–8.2	1.8–10.4	8.1–20.5
Hispanic	%	87.5	7.0	5.5	0.0	0.0	12.5
(n=104)	CI	79.7–92.7	3.4–13.6	2.4–12.1	–	–	7.3–20.3
8th	%	92.6	3.6	2.1	0.5	1.2	7.4
(n=1,887)	CI	91.4–93.7	2.9–4.5	1.5–2.8	0.2–0.9	0.8–1.8	6.3–8.6
10th	%	87.8	5.0	3.7	1.1	2.4	12.2
(n=1,274)	CI	85.9–89.5	3.9–6.3	2.8–4.9	0.6–1.8	1.7–3.4	10.5–14.1
12th	%	85.0	4.4	5.0	1.4	4.2	15.0
(n=1,118)	CI	82.8–87.0	3.3–5.7	3.9–6.4	0.9–2.3	3.2–5.5	13.0–17.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.8	1.6	5.6	91.0	96.6
(n=4,246)	CI	1.4–2.3	1.2–2.0	5.0–6.4	90.0–91.8	96.0–97.1
Male	%	1.7	1.2	5.7	91.4	97.1
(n=2,058)	CI	1.2–2.4	0.8–1.8	4.7–6.8	90.1–92.6	96.2–97.8
Female	%	1.3	1.8	5.3	91.6	97.0
(n=2,063)	CI	0.9–1.9	1.3–2.4	4.4–6.4	90.3–92.8	96.1–97.6
White	%	1.4	1.4	5.8	91.3	97.2
(n=3,441)	CI	1.0–1.9	1.1–1.9	5.1–6.7	90.3–92.3	96.5–97.7
Black	%	5.5	3.0	5.5	86.0	91.5
(n=129)	CI	2.5–11.7	1.1–8.1	2.7–10.9	78.6–91.1	84.7–95.4
Hispanic	%	6.2	0.0	3.5	90.3	93.8
(n=103)	CI	2.8–13.5	–	1.3–9.1	82.5–94.8	86.5–97.2
8th	%	1.0	1.2	4.6	93.1	97.8
(n=1,883)	CI	0.6–1.6	0.8–1.8	3.8–5.7	91.9–94.2	97.0–98.3
10th	%	1.7	1.9	5.3	91.1	96.4
(n=1,262)	CI	1.1–2.6	1.3–2.8	4.2–6.7	89.4–92.5	95.2–97.3
12th	%	2.7	1.6	7.1	88.5	95.7
(n=1,101)	CI	1.9–3.9	1.0–2.6	5.8–8.8	86.5–90.3	94.3–96.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.8	5.3	11.8	80.2	91.9
(n=4,261)	CI	2.3–3.4	4.6–6.0	10.8–12.8	78.9–81.4	91.0–92.7
Male	%	3.4	5.9	12.7	78.0	90.7
(n=2,062)	CI	2.6–4.3	4.9–7.1	11.2–14.2	76.1–79.8	89.3–92.0
Female	%	1.6	4.5	10.6	83.3	93.9
(n=2,073)	CI	1.1–2.3	3.6–5.5	9.3–12.1	81.6–84.9	92.8–94.9
White	%	2.5	5.0	12.3	80.1	92.4
(n=3,450)	CI	2.0–3.1	4.3–5.9	11.2–13.5	78.7–81.5	91.5–93.3
Black	%	6.0	4.2	8.1	81.7	89.9
(n=131)	CI	2.8–12.1	1.7–9.8	4.5–14.4	73.7–87.7	82.8–94.2
Hispanic	%	6.7	4.2	8.9	80.1	89.1
(n=103)	CI	3.0–14.3	1.7–10.1	4.6–16.6	70.9–87.0	81.0–94.0
8th	%	1.5	2.7	8.5	87.3	95.8
(n=1,889)	CI	1.0–2.1	2.1–3.5	7.3–9.9	85.7–88.7	94.8–96.6
10th	%	2.4	5.1	11.7	80.8	92.5
(n=1,266)	CI	1.7–3.4	4.0–6.5	10.1–13.6	78.5–82.9	90.9–93.9
12th	%	4.7	8.3	15.3	71.7	87.0
(n=1,106)	CI	3.6–6.1	6.8–10.1	13.3–17.6	69.0–74.3	84.9–88.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.45. Parents Feel Regularly Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.7	9.6	13.6	70.1	83.7
(n=4,259)	CI	6.0–7.6	8.7–10.6	12.6–14.7	68.7–71.5	82.5–84.8
Male	%	7.1	9.4	14.4	69.2	83.6
(n=2,063)	CI	6.0–8.3	8.1–10.8	12.9–16.0	67.1–71.2	81.8–85.2
Female	%	5.5	9.6	12.7	72.2	84.9
(n=2,071)	CI	4.6–6.7	8.3–11.0	11.3–14.3	70.1–74.1	83.2–86.4
White	%	6.4	9.3	14.4	70.0	84.3
(n=3,450)	CI	5.6–7.3	8.3–10.4	13.2–15.6	68.3–71.5	83.0–85.5
Black	%	6.0	10.2	10.3	73.6	83.8
(n=131)	CI	2.8–12.1	5.8–17.3	6.1–16.9	64.9–80.7	75.9–89.5
Hispanic	%	14.7	7.8	7.5	70.0	77.5
(n=103)	CI	8.8–23.6	3.8–15.2	3.7–14.6	60.0–78.4	67.8–84.8
8th	%	3.2	4.4	11.6	80.7	92.3
(n=1,889)	CI	2.5–4.1	3.6–5.5	10.3–13.2	78.8–82.4	91.0–93.4
10th	%	5.7	10.4	13.8	70.1	83.9
(n=1,266)	CI	4.5–7.1	8.8–12.2	12.1–15.9	67.5–72.6	81.8–85.9
12th	%	11.6	14.4	15.4	58.6	74.0
(n=1,104)	CI	9.9–13.7	12.4–16.5	13.4–17.7	55.6–61.4	71.3–76.5

Note: The confidence interval (CI) is 95 percent.

Table BH5.46. Parents Feel Smoking Tobacco is Wrong

		How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.9	4.3	10.9	81.9	92.8
(n=4,254)	CI	2.4–3.5	3.7–5.0	10.0–11.9	80.7–83.1	92.0–93.6
Male	%	3.9	5.0	11.8	79.2	91.0
(n=2,059)	CI	3.1–4.9	4.1–6.2	10.5–13.4	77.3–81.0	89.6–92.3
Female	%	1.4	3.2	9.6	85.7	95.4
(n=2,069)	CI	1.0–2.1	2.5–4.1	8.4–11.1	84.1–87.2	94.3–96.2
White	%	2.6	4.3	11.5	81.7	93.1
(n=3,446)	CI	2.1–3.2	3.6–5.1	10.4–12.6	80.3–83.0	92.2–94.0
Black	%	6.2	1.9	10.0	82.0	92.0
(n=131)	CI	2.9–12.5	0.5–7.2	5.9–16.5	74.1–87.9	85.2–95.8
Hispanic	%	7.5	7.8	4.7	80.0	84.7
(n=103)	CI	3.5–15.1	3.7–15.7	1.9–11.1	70.6–87.0	75.6–90.8
8th	%	1.3	1.4	6.1	91.2	97.3
(n=1,887)	CI	0.9–1.9	1.0–2.1	5.1–7.3	89.8–92.4	96.5–97.9
10th	%	2.4	3.9	11.0	82.8	93.8
(n=1,263)	CI	1.7–3.4	2.9–5.1	9.4–12.8	80.6–84.8	92.3–95.0
12th	%	5.2	7.8	16.2	70.8	87.0
(n=1,104)	CI	4.1–6.7	6.3–9.5	14.1–18.4	68.1–73.5	84.9–88.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	1.9	1.6	5.8	90.7	96.4
(n=4,254)	CI	1.5–2.4	1.3–2.1	5.1–6.5	89.7–91.6	95.8–97.0
Male	%	2.4	1.8	6.4	89.4	95.8
(n=2,058)	CI	1.8–3.2	1.3–2.5	5.4–7.7	87.9–90.7	94.8–96.7
Female	%	1.0	1.3	4.8	92.9	97.7
(n=2,071)	CI	0.6–1.5	0.9–1.9	3.9–5.9	91.6–94.0	96.9–98.3
White	%	1.6	1.5	6.0	90.9	96.9
(n=3,446)	CI	1.2–2.1	1.1–2.0	5.2–6.9	89.9–91.9	96.2–97.5
Black	%	4.4	0.9	6.4	88.3	94.7
(n=131)	CI	1.8–10.2	0.1–6.4	3.3–12.0	81.3–93.0	88.6–97.6
Hispanic	%	6.7	0.8	1.1	91.3	92.5
(n=102)	CI	3.0–14.4	0.1–5.5	0.2–7.9	83.4–95.7	84.8–96.5
8th	%	0.8	0.8	3.1	95.3	98.4
(n=1,886)	CI	0.5–1.4	0.5–1.3	2.4–4.0	94.2–96.2	97.7–98.8
10th	%	2.0	1.4	4.6	92.0	96.6
(n=1,264)	CI	1.3–2.9	0.9–2.2	3.6–6.0	90.3–93.3	95.5–97.5
12th	%	3.1	2.8	9.8	84.3	94.1
(n=1,104)	CI	2.2–4.3	2.0–4.0	8.2–11.7	82.0–86.3	92.6–95.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.48. Parents Feel Using Electronic Cigarettes is Wrong

		How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.3	8.1	14.0	72.6	86.6
(n=4,249)	CI	4.6–6.1	7.2–9.0	13.0–15.1	71.2–74.0	85.5–87.7
Male	%	6.3	8.6	14.8	70.3	85.1
(n=2,056)	CI	5.2–7.5	7.4–9.9	13.3–16.5	68.2–72.3	83.4–86.7
Female	%	3.6	7.6	12.6	76.1	88.8
(n=2068)	CI	2.9–4.5	6.5–8.9	11.2–14.2	74.2–78.0	87.3–90.1
White	%	5.0	8.5	14.5	72.0	86.5
(n=3,444)	CI	4.3–5.9	7.6–9.5	13.3–15.7	70.4–73.5	85.2–87.6
Black	%	9.1	4.8	13.2	72.8	86.1
(n=131)	CI	5.0–15.8	2.1–10.6	8.4–20.4	64.3–80.0	78.6–91.3
Hispanic	%	10.4	4.3	10.7	74.7	85.4
(n=102)	CI	5.4–18.9	1.7–10.2	5.9–18.7	64.8–82.5	76.5–91.3
8th	%	2.7	3.8	9.4	84.1	93.5
(n=1,883)	CI	2.1–3.5	3.0–4.8	8.2–10.8	82.3–85.7	92.3–94.5
10th	%	4.7	8.3	14.3	72.6	87.0
(n=1,260)	CI	3.7–6.1	6.9–10.0	12.5–16.4	70.1–75.0	85.0–88.7
12th	%	8.7	12.4	18.7	60.1	78.8
(n=1,106)	CI	7.2–10.6	10.6–14.5	16.5–21.1	57.2–62.9	76.3–81.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.9	6.0	9.3	79.8	89.1
(n=4,258)	CI	4.2–5.6	5.3–6.8	8.5–10.3	78.5–81.0	88.1–90.1
Male	%	5.3	5.6	10.1	79.0	89.1
(n=2,061)	CI	4.3–6.4	4.6–6.8	8.8–11.5	77.1–80.8	87.6–90.5
Female	%	3.6	6.1	8.5	81.7	90.3
(n=2,072)	CI	2.8–4.6	5.1–7.3	7.3–9.9	79.9–83.4	88.8–91.5
White	%	4.4	6.1	9.4	80.0	89.4
(n=3,449)	CI	3.8–5.2	5.3–7.0	8.5–10.5	78.5–81.3	88.3–90.5
Black	%	8.6	4.5	7.9	79.0	86.9
(n=131)	CI	4.7–15.4	1.9–10.5	4.2–14.3	70.6–85.5	79.3–92.0
Hispanic	%	7.5	3.1	7.8	81.6	89.4
(n=103)	CI	3.5–15.1	1.0–9.6	3.7–15.7	72.2–88.4	81.2–94.3
8th	%	1.9	2.0	5.0	91.0	96.1
(n=1,888)	CI	1.4–2.6	1.5–2.8	4.1–6.1	89.7–92.3	95.1–96.9
10th	%	5.4	6.8	9.2	78.7	87.9
(n=1,266)	CI	4.2–6.7	5.5–8.3	7.7–10.9	76.3–80.8	86.0–89.6
12th	%	7.6	9.5	14.2	68.8	82.9
(n=1,104)	CI	6.2–9.3	7.9–11.4	12.2–16.4	66.0–71.4	80.6–85.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	4.3	4.0	7.9	83.9	91.7
(n=4,249)	CI	3.7–5.0	3.4–4.6	7.0–8.8	82.7–85.0	90.8–92.6
Male	%	4.5	3.8	8.5	83.2	91.6
(n=2,057)	CI	3.7–5.6	3.0–4.8	7.3–9.8	81.4–84.8	90.3–92.8
Female	%	3.2	4.0	7.0	85.8	92.8
(n=2,067)	CI	2.5–4.1	3.2–5.0	5.9–8.2	84.2–87.3	91.5–93.9
White	%	3.9	4.0	8.1	84.0	92.1
(n=3,440)	CI	3.3–4.6	3.3–4.7	7.2–9.1	82.7–85.3	91.1–93.0
Black	%	8.6	3.7	5.4	82.3	87.7
(n=131)	CI	4.7–15.4	1.4–9.4	2.5–11.2	74.2–88.2	80.3–92.6
Hispanic	%	6.7	3.9	3.1	86.3	89.4
(n=103)	CI	3.0–14.3	1.4–10.3	1.0–9.6	77.5–92.0	81.2–94.3
8th	%	1.7	1.2	4.1	93.0	97.1
(n=1,884)	CI	1.2–2.4	0.8–1.8	3.3–5.1	91.7–94.1	96.2–97.8
10th	%	4.3	4.9	7.9	82.9	90.8
(n=1,259)	CI	3.3–5.5	3.8–6.2	6.5–9.5	80.7–84.9	89.1–92.3
12th	%	7.2	5.9	11.9	75.0	86.8
(n=1,106)	CI	5.8–8.9	4.7–7.5	10.1–13.9	72.3–77.4	84.7–88.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.51. Friends Feel Misusing Prescription Drugs is Wrong

		How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.0	7.2	0.0	69.1	85.8
(n=4,235)	CI	6.3–7.9	6.5–8.1	–	67.7–70.6	84.6–86.8
Male	%	7.6	7.1	0.0	66.6	85.3
(n=2,046)	CI	6.5–8.9	6.1–8.4	–	64.5–68.7	83.6–86.8
Female	%	5.3	7.3	0.0	72.4	87.4
(n=2,065)	CI	4.4–6.4	6.2–8.6	–	70.4–74.4	85.8–88.8
White	%	6.4	7.2	17.1	69.3	86.3
(n=3,433)	CI	5.6–7.3	6.4–8.2	15.8–18.4	67.7–70.8	85.1–87.5
Black	%	13.9	6.7	19.3	60.1	79.5
(n=130)	CI	8.7–21.4	3.4–12.6	13.2–27.4	51.2–68.5	71.3–85.8
Hispanic	%	13.5	9.0	10.1	67.3	77.5
(n=103)	CI	7.9–22.2	4.8–16.3	5.6–17.8	57.4–75.9	68.0–84.8
8th	%	4.6	5.2	0.0	77.4	90.1
(n=1,877)	CI	3.8–5.7	4.3–6.3	–	75.4–79.2	88.7–91.4
10th	%	8.3	7.1	0.0	66.9	84.6
(n=1,258)	CI	6.9–10.0	5.8–8.7	–	64.3–69.4	82.5–86.5
12th	%	8.2	9.5	0.0	62.6	82.2
(n=1,100)	CI	6.8–10.0	7.9–11.4	–	59.7–65.4	79.9–84.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.52. Friends Feel Drinking Alcohol is Wrong

		How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.0	14.8	0.0	50.8	71.2
(n=4,251)	CI	12.9–15.1	13.7–15.9	–	49.3–52.4	69.8–72.6
Male	%	16.3	15.0	0.0	48.1	68.7
(n=2,054)	CI	14.7–18.1	13.4–16.7	–	45.9–50.3	66.6–70.7
Female	%	10.7	14.5	0.0	54.3	74.7
(n=2,072)	CI	9.4–12.2	13.0–16.2	–	52.1–56.5	72.7–76.6
White	%	13.8	15.0	20.9	50.3	71.2
(n=3,443)	CI	12.7–15.1	13.8–16.2	19.6–22.3	48.6–52.0	69.6–72.7
Black	%	18.6	14.1	18.5	48.7	67.3
(n=131)	CI	12.5–26.8	9.0–21.5	12.6–26.4	40.0–57.5	58.4–75.1
Hispanic	%	13.2	15.0	23.9	47.9	71.8
(n=103)	CI	7.7–21.7	9.1–23.7	16.4–33.4	38.1–57.9	61.9–80.0
8th	%	6.2	9.1	0.0	65.9	84.7
(n=1,884)	CI	5.2–7.4	7.9–10.5	–	63.8–68.0	83.0–86.3
10th	%	15.2	17.1	0.0	46.7	67.6
(n=1,262)	CI	13.4–17.3	15.1–19.3	–	44.0–49.5	65.0–70.2
12th	%	21.2	18.5	0.0	38.8	60.3
(n=1,105)	CI	18.9–23.7	16.3–20.9	–	35.9–41.7	57.4–63.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	13.7	12.3	0.0	55.1	74.0
(n=4,241)	CI	12.6–14.8	11.3–13.4	–	53.6–56.6	72.6–75.4
Male	%	16.7	12.2	0.0	51.1	71.1
(n=2,047)	CI	15.1–18.5	10.8–13.7	–	48.9–53.3	69.0–73.1
Female	%	9.6	12.3	0.0	59.8	78.0
(n=2,069)	CI	8.4–11.1	10.9–13.9	–	57.6–61.9	76.1–79.9
White	%	13.6	12.4	19.5	54.5	74.1
(n=3,434)	CI	12.4–14.8	11.3–13.5	18.2–20.9	52.8–56.2	72.5–75.6
Black	%	20.2	11.7	16.2	52.0	68.1
(n=131)	CI	13.8–28.4	7.1–18.8	10.6–23.8	43.2–60.6	59.3–75.8
Hispanic	%	14.0	10.5	16.8	58.7	75.5
(n=103)	CI	8.2–22.8	5.8–18.3	10.6–25.5	48.6–68.2	65.8–83.2
8th	%	5.5	7.5	0.0	70.9	86.9
(n=1,882)	CI	4.6–6.7	6.4–8.8	–	68.8–72.9	85.3–88.4
10th	%	14.0	13.5	0.0	52.0	72.5
(n=1,257)	CI	12.2–16.1	11.7–15.5	–	49.2–54.7	70.0–74.9
12th	%	22.2	16.3	0.0	41.2	61.6
(n=1,102)	CI	19.8–24.7	14.2–18.6	–	38.4–44.2	58.7–64.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	27.6	14.5	14.3	43.5	57.9
(n=4,244)	CI	26.2–29.0	13.5–15.6	13.3–15.5	42.0–45.0	56.3–59.4
Male	%	29.5	14.3	14.3	41.9	56.2
(n=2,050)	CI	27.4–31.6	12.9–16.0	12.8–15.9	39.8–44.1	54.0–58.4
Female	%	24.8	14.7	14.9	45.6	60.5
(n=2,070)	CI	22.9–26.8	13.2–16.4	13.4–16.5	43.4–47.7	58.3–62.6
White	%	28.3	14.8	14.3	42.7	57.0
(n=3,438)	CI	26.7–29.9	13.6–16.0	13.1–15.5	41.0–44.4	55.3–58.7
Black	%	33.3	13.1	14.7	38.9	53.6
(n=131)	CI	25.4–42.2	8.3–20.1	9.5–22.0	30.8–47.7	44.8–62.2
Hispanic	%	19.3	11.1	19.7	49.9	69.6
(n=103)	CI	12.5–28.6	6.1–19.3	13.0–28.7	40.0–59.8	59.5–78.0
8th	%	12.7	12.7	14.9	59.8	74.6
(n=1,881)	CI	11.2–14.2	11.3–14.3	13.3–16.6	57.5–62.0	72.6–76.6
10th	%	31.1	15.2	14.5	39.1	53.7
(n=1,260)	CI	28.6–33.8	13.3–17.3	12.7–16.6	36.5–41.9	50.9–56.4
12th	%	40.1	15.8	13.6	30.5	44.1
(n=1,103)	CI	37.3–43.0	13.8–18.1	11.7–15.7	27.9–33.3	41.2–47.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	22.0	12.2	0.0	51.5	65.9
(n=4,234)	CI	20.7–23.3	11.2–13.2	–	49.9–53.0	64.4–67.4
Male	%	23.4	11.7	0.0	49.4	64.9
(n=2,049)	CI	21.5–25.4	10.3–13.3	–	47.2–51.6	62.7–67.0
Female	%	19.3	12.6	0.0	54.3	68.1
(n=2,061)	CI	17.6–21.2	11.2–14.2	–	52.1–56.5	65.9–70.1
White	%	21.9	12.3	14.4	51.5	65.8
(n=3,431)	CI	20.5–23.4	11.2–13.5	13.2–15.6	49.8–53.2	64.2–67.5
Black	%	28.4	12.4	14.0	45.3	59.3
(n=129)	CI	20.9–37.2	7.5–19.7	9.0–21.3	36.7–54.1	50.2–67.8
Hispanic	%	17.0	11.4	17.6	54.1	71.6
(n=103)	CI	10.6–26.2	6.4–19.3	11.3–26.4	44.1–63.8	61.7–79.8
8th	%	7.4	6.5	0.0	71.8	86.1
(n=1,878)	CI	6.3–8.7	5.5–7.7	–	69.7–73.8	84.5–87.6
10th	%	24.3	12.9	0.0	46.6	62.8
(n=1,258)	CI	22.0–26.8	11.2–14.9	–	43.9–49.4	60.1–65.4
12th	%	35.3	17.5	0.0	34.5	47.2
(n=1,098)	CI	32.5–38.2	15.4–19.9	–	31.7–37.3	44.2–50.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	87.2	6.0	2.8	1.0	2.9	12.8
(n=4,260)	CI	86.1–88.3	5.3–6.8	2.3–3.4	0.7–1.3	2.5–3.5	11.7–13.9
Male	%	88.4	5.4	2.4	0.9	3.0	11.6
(n=2,064)	CI	86.9–89.8	4.4–6.5	1.8–3.2	0.5–1.4	2.3–3.9	10.2–13.1
Female	%	86.7	6.8	3.1	1.0	2.4	13.3
(n=2,073)	CI	85.1–88.2	5.7–8.0	2.4–4.0	0.6–1.6	1.8–3.2	11.8–14.9
White	%	87.2	6.3	2.6	1.1	2.8	12.8
(n=3,458)	CI	86.0–88.3	5.5–7.2	2.1–3.2	0.7–1.5	2.3–3.5	11.7–14.0
Black	%	85.5	0.9	4.7	0.9	8.0	14.5
(n=131)	CI	77.8–90.9	0.1–6.5	2.1–10.3	0.1–6.0	4.2–14.7	9.1–22.2
Hispanic	%	87.3	7.1	1.9	1.1	2.7	12.7
(n=104)	CI	79.0–92.6	3.3–14.4	0.5–7.6	0.2–7.7	0.8–8.2	7.4–21.0
8th	%	93.9	3.5	1.3	0.2	1.1	6.1
(n=1,876)	CI	92.7–94.9	2.7–4.4	0.9–1.9	0.1–0.6	0.7–1.7	5.1–7.3
10th	%	84.5	6.4	3.6	1.9	3.5	15.5
(n=1,267)	CI	82.4–86.4	5.2–7.9	2.7–4.8	1.3–2.8	2.7–4.7	13.6–17.6
12th	%	82.9	8.4	3.6	0.8	4.3	17.1
(n=1,117)	CI	80.6–85.0	6.9–10.2	2.6–4.8	0.4–1.5	3.2–5.6	15.0–19.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	54.3	12.2	8.9	5.4	19.2	45.7
(n=4,286)	CI	52.8–55.8	11.2–13.2	8.1–9.9	4.8–6.2	17.9–20.4	44.2–47.2
Male	%	58.2	10.3	7.0	4.7	19.8	41.8
(n=2,073)	CI	56.0–60.3	9.1–11.8	5.9–8.2	3.8–5.7	18.1–21.7	39.7–44.0
Female	%	51.2	13.9	10.8	6.2	17.8	48.8
(n=2,086)	CI	49.0–53.4	12.5–15.5	9.5–12.3	5.2–7.4	16.2–19.6	46.6–51.0
White	%	53.5	12.0	9.2	5.5	19.7	46.5
(n=3,477)	CI	51.8–55.2	11.0–13.2	8.3–10.3	4.8–6.4	18.3–21.1	44.8–48.2
Black	%	55.3	10.3	6.7	3.0	24.6	44.7
(n=129)	CI	46.4–64.0	6.0–17.2	3.4–13.1	1.1–7.8	17.6–33.4	36.0–53.6
Hispanic	%	63.7	15.2	5.4	3.8	11.9	36.3
(n=105)	CI	53.7–72.7	9.4–23.7	2.2–12.5	1.4–10.1	6.8–20.1	27.3–46.3
8th	%	76.8	10.1	5.5	2.1	5.6	23.2
(n=1,885)	CI	74.8–78.6	8.8–11.5	4.5–6.6	1.6–2.9	4.6–6.7	21.4–25.2
10th	%	49.1	14.5	9.7	5.9	20.8	50.9
(n=1,279)	CI	46.3–51.8	12.7–16.5	8.2–11.5	4.8–7.4	18.7–23.1	48.2–53.7
12th	%	35.8	12.0	11.8	8.4	32.0	64.2
(n=1,122)	CI	33.0–38.6	10.2–14.0	10.1–13.8	6.9–10.2	29.3–34.8	61.4–67.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	67.6	9.5	5.5	4.1	13.3	32.4
(n=4,281)	CI	66.1–69.1	8.6–10.5	4.8–6.3	3.5–4.8	12.2–14.4	30.9–33.9
Male	%	69.3	8.1	4.1	3.9	14.6	30.7
(n=2,070)	CI	67.1–71.3	7.0–9.5	3.3–5.1	3.1–4.9	13.1–16.3	28.7–32.9
Female	%	66.7	10.7	7.1	4.1	11.4	33.3
(n=2,085)	CI	64.6–68.8	9.4–12.2	6.0–8.4	3.3–5.1	10.0–12.9	31.2–35.4
White	%	66.8	9.6	5.9	4.1	13.6	33.2
(n=3,472)	CI	65.2–68.5	8.6–10.7	5.2–6.8	3.4–4.8	12.4–14.8	31.5–34.8
Black	%	66.5	9.0	4.2	5.3	15.0	33.5
(n=131)	CI	57.4–74.5	4.9–16.0	1.7–9.8	2.4–11.4	9.5–22.9	25.5–42.6
Hispanic	%	77.8	10.6	0.8	2.0	8.8	22.2
(n=105)	CI	68.3–85.1	5.7–18.9	0.1–5.4	0.5–7.9	4.6–16.5	14.9–31.7
8th	%	89.6	4.7	1.7	1.0	2.9	10.4
(n=1,883)	CI	88.2–90.9	3.9–5.8	1.2–2.4	0.6–1.6	2.2–3.8	9.1–11.8
10th	%	64.6	11.2	6.2	4.2	13.7	35.4
(n=1,277)	CI	61.9–67.2	9.6–13.1	5.0–7.7	3.2–5.5	11.9–15.7	32.8–38.1
12th	%	47.3	12.8	8.9	7.2	23.8	52.7
(n=1,121)	CI	44.4–50.2	11.0–14.9	7.4–10.7	5.8–8.9	21.4–26.4	49.8–55.6

Note: The confidence interval (CI) is 95 percent.

Table BH5.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have smoked part, or all of a cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	74.5	9.7	5.3	3.0	7.5	25.5
(n=4,261)	CI	73.1–75.8	8.9–10.7	4.7–6.1	2.5–3.6	6.7–8.4	24.2–26.9
Male	%	75.0	9.6	4.6	2.8	8.1	25.0
(n=2,068)	CI	73.0–76.9	8.3–11.0	3.7–5.6	2.1–3.6	7.0–9.5	23.1–27.0
Female	%	74.8	10.0	6.0	3.0	6.2	25.2
(n=2,070)	CI	72.8–76.7	8.7–11.4	5.0–7.2	2.3–3.9	5.2–7.4	23.3–27.2
White	%	73.7	9.9	5.7	2.9	7.8	26.3
(n=3,459)	CI	72.2–75.2	8.9–11.0	4.9–6.6	2.4–3.6	6.9–8.7	24.8–27.8
Black	%	79.7	6.2	3.6	1.5	8.9	20.3
(n=130)	CI	71.3–86.1	3.1–12.2	1.4–9.4	0.4–6.3	4.8–15.9	13.9–28.7
Hispanic	%	81.2	9.3	1.9	2.0	5.7	18.8
(n=104)	CI	72.2–87.7	4.9–16.8	0.5–7.6	0.5–8.0	2.5–12.3	12.3–27.8
8th	%	87.1	6.7	2.4	1.1	2.8	12.9
(n=1,879)	CI	85.5–88.6	5.6–7.9	1.8–3.2	0.7–1.6	2.1–3.6	11.4–14.5
10th	%	71.9	9.8	5.9	3.4	9.1	28.1
(n=1,264)	CI	69.3–74.3	8.3–11.5	4.7–7.4	2.5–4.5	7.6–10.8	25.7–30.7
12th	%	63.6	13.0	7.8	4.6	10.9	36.4
(n=1,118)	CI	60.8–66.4	11.2–15.1	6.4–9.6	3.5–6.0	9.2–12.8	33.6–39.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used an e-cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	55.6	10.2	6.9	5.1	22.2	44.4
(n=4,264)	CI	54.0–57.1	9.3–11.2	6.1–7.7	4.4–5.8	21.0–23.6	42.9–46.0
Male	%	55.7	9.3	5.7	5.2	24.1	44.3
(n=2,063)	CI	53.4–57.8	8.1–10.7	4.7–6.8	4.2–6.3	22.3–26.1	42.2–46.6
Female	%	55.8	11.3	8.1	5.2	19.6	44.2
(n=2,076)	CI	53.6–57.9	10.0–12.8	7.0–9.4	4.3–6.3	17.8–21.4	42.1–46.4
White	%	53.9	10.6	7.1	5.3	23.1	46.1
(n=3,459)	CI	52.2–55.6	9.6–11.7	6.3–8.0	4.5–6.1	21.6–24.6	44.4–47.8
Black	%	58.3	7.7	7.3	3.9	22.9	41.7
(n=131)	CI	49.3–66.7	4.1–14.0	3.8–13.7	1.6–9.1	16.1–31.4	33.3–50.7
Hispanic	%	72.1	10.1	1.5	2.8	13.5	27.9
(n=104)	CI	62.3–80.2	5.4–18.0	0.4–6.1	0.8–8.6	7.9–22.2	19.8–37.7
8th	%	73.6	8.7	5.3	2.7	9.7	26.4
(n=1,877)	CI	71.5–75.5	7.5–10.1	4.3–6.4	2.0–3.5	8.5–11.2	24.5–28.5
10th	%	50.5	11.0	6.6	6.0	25.8	49.5
(n=1,269)	CI	47.8–53.3	9.4–12.8	5.4–8.1	4.9–7.5	23.5–28.3	46.7–52.2
12th	%	41.6	11.1	8.9	6.6	31.8	58.4
(n=1,118)	CI	38.7–44.5	9.4–13.0	7.4–10.7	5.3–8.2	29.2–34.6	55.5–61.3

Note: The confidence interval (CI) is 95 percent.

Table BH5.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	66.8	9.2	5.8	4.0	14.3	33.2
(n=4,276)	CI	65.3–68.2	8.3–10.1	5.1–6.6	3.4–4.6	13.2–15.5	31.8–34.7
Male	%	68.4	8.6	4.7	3.7	14.5	31.6
(n=2,068)	CI	66.3–70.5	7.5–10.0	3.8–5.8	2.9–4.7	13.0–16.2	29.5–33.7
Female	%	65.8	9.9	7.0	4.2	13.1	34.2
(n=2,082)	CI	63.7–67.9	8.6–11.3	5.9–8.3	3.4–5.2	11.6–14.7	32.1–36.3
White	%	66.4	9.3	6.0	4.0	14.3	33.6
(n=3,472)	CI	64.8–68.1	8.3–10.4	5.2–6.9	3.3–4.7	13.1–15.5	31.9–35.2
Black	%	62.2	8.7	4.3	4.3	20.6	37.8
(n=131)	CI	53.2–70.4	4.8–15.3	1.7–10.0	1.7–10.0	14.2–29.0	29.6–46.8
Hispanic	%	78.4	5.8	1.5	1.2	13.1	21.6
(n=104)	CI	68.9–85.6	2.5–12.6	0.4–6.1	0.2–8.3	7.5–21.8	14.4–31.1
8th	%	88.1	5.9	2.0	1.0	3.0	11.9
(n=1,881)	CI	86.5–89.5	4.9–7.1	1.5–2.8	0.6–1.6	2.3–3.8	10.5–13.5
10th	%	62.3	10.5	5.1	5.1	17.0	37.7
(n=1,277)	CI	59.6–64.9	9.0–12.3	4.0–6.4	4.0–6.4	15.0–19.2	35.1–40.4
12th	%	48.6	11.2	10.6	6.0	23.6	51.4
(n=1,118)	CI	45.7–51.5	9.5–13.2	8.9–12.6	4.7–7.5	21.2–26.2	48.5–54.3

Note: The confidence interval (CI) is 95 percent.

Table BH5.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	7.2	5.4	14.8	72.6	87.4
(n=4,227)	CI	6.4–8.0	4.8–6.2	13.8–15.9	71.2–73.9	86.3–88.3
Male	%	9.3	5.1	15.7	70.0	85.6
(n=2,047)	CI	8.1–10.7	4.2–6.1	14.1–17.3	67.9–71.9	84.0–87.1
Female	%	4.2	5.7	13.8	76.3	90.1
(n=2,054)	CI	3.4–5.2	4.7–6.8	12.4–15.4	74.3–78.1	88.7–91.3
White	%	7.0	5.4	14.4	73.2	87.6
(n=3,427)	CI	6.1–7.9	4.7–6.2	13.3–15.7	71.6–74.7	86.5–88.7
Black	%	11.5	8.7	17.7	62.1	79.8
(n=131)	CI	6.9–18.4	4.8–15.3	12.0–25.4	53.2–70.2	71.7–86.0
Hispanic	%	13.2	5.0	17.3	64.5	81.8
(n=103)	CI	7.8–21.4	2.0–11.8	11.0–26.2	54.5–73.4	72.8–88.3
8th	%	7.0	5.6	15.3	72.1	87.4
(n=1,867)	CI	5.9–8.3	4.7–6.8	13.7–17.0	70.0–74.1	85.8–88.8
10th	%	7.5	5.9	15.3	71.3	86.6
(n=1,256)	CI	6.2–9.1	4.7–7.3	13.4–17.4	68.7–73.7	84.6–88.4
12th	%	7.0	4.8	13.8	74.4	88.2
(n=1,104)	CI	5.7–8.7	3.7–6.2	11.9–16.0	71.7–76.9	86.1–90.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.63. Perceived Risk of Regularly Drinking Alcohol

		How much do you think people risk harming themselves physically and in other ways when they <u>take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	11.8	14.8	24.7	48.7	73.4
(n=4,236)	CI	10.9–12.8	13.7–15.9	23.4–26.1	47.2–50.2	72.1–74.8
Male	%	15.4	17.4	25.1	42.1	67.2
(n=2,051)	CI	13.9–17.1	15.7–19.1	23.2–27.1	39.9–44.3	65.1–69.3
Female	%	7.2	12.2	24.6	56.0	80.6
(n=2,060)	CI	6.1–8.4	10.8–13.7	22.8–26.6	53.8–58.2	78.8–82.3
White	%	11.8	15.4	25.3	47.5	72.8
(n=3,434)	CI	10.8–13.0	14.2–16.7	23.8–26.8	45.8–49.2	71.2–74.3
Black	%	12.4	14.0	23.5	50.1	73.6
(n=131)	CI	7.6–19.5	8.9–21.2	16.8–31.7	41.4–58.9	65.2–80.6
Hispanic	%	18.8	7.8	23.5	49.9	73.4
(n=102)	CI	12.2–28.0	4.0–14.6	16.0–33.0	40.0–59.8	63.7–81.2
8th	%	10.4	12.7	22.0	55.0	77.0
(n=1,872)	CI	9.1–11.8	11.2–14.2	20.1–23.9	52.8–57.3	75.0–78.8
10th	%	13.2	14.7	25.1	47.0	72.1
(n=1,259)	CI	11.5–15.2	12.9–16.8	22.8–27.5	44.3–49.8	69.5–74.5
12th	%	11.9	17.0	27.3	43.7	71.1
(n=1,105)	CI	10.1–13.9	14.9–19.4	24.8–30.0	40.8–46.7	68.3–73.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.64. Perceived Risk of Binge Drinking

		How much do you think people risk harming themselves physically and in other ways when they <u>take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	9.5	10.2	21.7	58.5	80.3
(n=4,244)	CI	8.6–10.4	9.3–11.2	20.5–23.0	57.0–60.1	79.0–81.5
Male	%	12.5	11.8	24.0	51.7	75.7
(n=2,058)	CI	11.1–14.0	10.4–13.3	22.2–26.0	49.5–53.9	73.8–77.6
Female	%	5.4	8.7	19.6	66.3	85.9
(n=2,061)	CI	4.5–6.5	7.5–10.0	17.9–21.4	64.2–68.4	84.3–87.4
White	%	9.3	10.7	22.3	57.6	79.9
(n=3,442)	CI	8.4–10.4	9.7–11.8	20.9–23.8	55.9–59.3	78.5–81.3
Black	%	11.5	10.2	18.2	60.1	78.4
(n=131)	CI	6.9–18.4	5.8–17.2	12.3–26.2	51.2–68.4	70.1–84.8
Hispanic	%	16.1	8.9	17.6	57.3	74.9
(n=102)	CI	10.0–25.1	4.8–16.2	11.2–26.6	47.2–66.8	65.3–82.6
8th	%	8.1	8.1	17.4	66.3	83.8
(n=1,878)	CI	6.9–9.4	7.0–9.5	15.8–19.2	64.2–68.5	82.0–85.4
10th	%	10.1	10.8	22.7	56.4	79.1
(n=1,262)	CI	8.6–11.9	9.2–12.6	20.4–25.0	53.7–59.1	76.8–81.2
12th	%	10.3	11.9	25.5	52.3	77.8
(n=1,104)	CI	8.6–12.2	10.1–14.0	23.0–28.1	49.4–55.3	75.3–80.2

Note: The confidence interval (CI) is 95 percent.

Table BH5.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	7.5	5.0	15.6	71.8	87.5
(n=4,246)	CI	6.7–8.4	4.4–5.8	14.5–16.8	70.4–73.2	86.4–88.4
Male	%	9.6	5.3	17.1	68.1	85.2
(n=2,059)	CI	8.3–10.9	4.4–6.4	15.5–18.8	66.0–70.1	83.5–86.7
Female	%	4.5	4.8	14.2	76.6	90.7
(n=2,060)	CI	3.7–5.5	3.9–5.8	12.7–15.8	74.6–78.4	89.3–91.9
White	%	7.5	4.9	15.6	72.1	87.6
(n=3,441)	CI	6.6–8.4	4.2–5.7	14.4–16.9	70.5–73.6	86.5–88.7
Black	%	9.0	8.1	14.2	68.7	82.9
(n=132)	CI	5.1–15.5	4.3–14.6	9.1–21.4	60.0–76.3	75.1–88.6
Hispanic	%	13.2	9.3	20.5	57.0	77.6
(n=103)	CI	7.8–21.4	4.8–17.2	13.7–29.6	47.0–66.5	68.1–84.8
8th	%	7.1	4.0	15.2	73.7	88.9
(n=1,881)	CI	6.0–8.4	3.2–5.0	13.7–16.9	71.6–75.6	87.4–90.2
10th	%	7.9	5.5	15.6	71.0	86.6
(n=1,262)	CI	6.5–9.5	4.4–6.9	13.7–17.7	68.4–73.4	84.6–88.4
12th	%	7.5	5.7	16.1	70.7	86.8
(n=1,103)	CI	6.1–9.2	4.5–7.2	14.0–18.4	68.0–73.3	84.7–88.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.3	20.0	29.3	40.4	69.7
(n=4245)	CI	9.4–11.3	18.8–21.3	27.9–30.7	38.9–41.9	68.2–71.1
Male	%	13.4	22.9	27.8	35.9	63.7
(n=2,058)	CI	12.0–15.0	21.0–24.8	25.9–29.9	33.8–38.0	61.6–65.8
Female	%	6.4	17.2	31.0	45.5	76.5
(n=2,062)	CI	5.4–7.5	15.6–18.9	28.9–33.0	43.4–47.7	74.6–78.3
White	%	10.3	20.4	30.2	39.1	69.3
(n=3,445)	CI	9.3–11.4	19.1–21.9	28.7–31.8	37.4–40.7	67.7–70.8
Black	%	10.4	23.6	29.0	37.0	66.0
(n=130)	CI	6.0–17.3	17.0–31.9	21.6–37.7	29.0–45.8	57.1–73.8
Hispanic	%	18.8	16.4	23.8	41.0	64.8
(n=102)	CI	12.1–27.9	10.3–25.2	16.2–33.4	31.6–51.2	54.7–73.7
8th	%	9.6	16.3	25.1	49.0	74.1
(n=1,880)	CI	8.3–11.0	14.7–18.1	23.1–27.1	46.8–51.3	72.1–76.0
10th	%	10.8	20.9	30.3	38.0	68.3
(n=1,261)	CI	9.2–12.7	18.7–23.2	27.9–32.9	35.4–40.7	65.7–70.8
12th	%	10.6	23.2	32.8	33.5	66.2
(n=1,104)	CI	8.9–12.5	20.8–25.8	30.1–35.6	30.8–36.3	63.4–69.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	30.3	21.4	15.6	32.7	48.3
(n=4,228)	CI	28.9–31.8	20.2–22.7	14.5–16.7	31.3–34.1	46.7–49.8
Male	%	34.5	21.1	15.7	28.7	44.4
(n=2,046)	CI	32.4–36.6	19.4–23.0	14.2–17.4	26.7–30.7	42.2–46.6
Female	%	24.8	22.4	15.7	37.0	52.7
(n=2,056)	CI	22.9–26.9	20.6–24.3	14.2–17.4	34.9–39.1	50.5–54.9
White	%	30.6	21.8	15.2	32.3	47.5
(n=3,428)	CI	29.1–32.3	20.5–23.3	14.0–16.4	30.8–33.9	45.8–49.2
Black	%	31.0	22.4	19.2	27.3	46.5
(n=132)	CI	23.4–39.9	16.0–30.6	13.2–27.0	20.4–35.6	38.0–55.3
Hispanic	%	27.8	11.0	16.6	44.5	61.1
(n=101)	CI	19.6–37.9	6.2–18.8	10.4–25.5	34.8–54.7	50.9–70.4
8th	%	15.9	18.8	18.8	46.5	65.3
(n=1,870)	CI	14.3–17.6	17.1–20.7	17.1–20.6	44.3–48.8	63.1–67.4
10th	%	31.8	23.1	15.8	29.3	45.1
(n=1,255)	CI	29.3–34.5	20.8–25.5	13.9–17.9	26.8–31.9	42.3–47.8
12th	%	44.3	22.5	11.9	21.2	33.2
(n=1,103)	CI	41.4–47.3	20.1–25.1	10.1–14.0	18.9–23.8	30.5–36.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	20.0	16.1	20.6	43.4	64.0
(n=4,232)	CI	18.7–21.2	14.9–17.3	19.4–21.9	41.9–44.9	62.5–65.5
Male	%	23.9	17.1	20.3	38.8	59.1
(n=2,053)	CI	22.0–25.8	15.4–18.8	18.5–22.1	36.7–41.0	56.9–61.2
Female	%	14.6	15.3	21.4	48.7	70.1
(n=2,054)	CI	13.1–16.3	13.7–17.0	19.6–23.2	46.5–50.9	68.0–72.1
White	%	19.9	16.5	20.8	42.7	63.6
(n=3,431)	CI	18.6–21.4	15.2–17.8	19.5–22.3	41.1–44.4	61.9–65.2
Black	%	24.2	16.1	21.2	38.5	59.6
(n=132)	CI	17.4–32.7	10.6–23.9	14.9–29.2	30.4–47.2	50.7–68.0
Hispanic	%	22.8	10.6	22.3	44.4	66.6
(n=102)	CI	15.3–32.5	5.8–18.4	15.1–31.5	34.7–54.5	56.5–75.4
8th	%	11.5	9.4	18.4	60.6	79.1
(n=1,872)	CI	10.2–13.1	8.2–10.8	16.7–20.3	58.4–62.8	77.2–80.8
10th	%	21.8	17.8	20.9	39.5	60.4
(n=1,260)	CI	19.6–24.2	15.8–20.0	18.7–23.2	36.9–42.3	57.7–63.1
12th	%	27.1	21.5	22.7	28.7	51.4
(n=1,100)	CI	24.5–29.8	19.1–24.0	20.4–25.3	26.1–31.5	48.5–54.4

Note: The confidence interval (CI) is 95 percent.

Table BH5.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.4	4.5	12.1	80.0	92.1
(n=4,248)	CI	2.9–4.0	3.9–5.2	11.1–13.2	78.7–81.2	91.2–92.9
Male	%	4.0	4.5	12.5	79.0	91.5
(n=2,060)	CI	3.2–4.9	3.7–5.6	11.1–14.1	77.1–80.8	90.2–92.7
Female	%	2.0	4.5	11.6	81.9	93.5
(n=2,063)	CI	1.4–2.7	3.7–5.5	10.2–13.1	80.2–83.6	92.3–94.5
White	%	3.0	4.2	12.2	80.6	92.8
(n=3,449)	CI	2.4–3.6	3.6–5.0	11.1–13.3	79.2–82.0	91.9–93.6
Black	%	7.4	8.7	11.4	72.5	83.9
(n=131)	CI	3.8–13.8	4.9–14.9	6.9–18.3	64.0–79.7	76.3–89.4
Hispanic	%	8.8	6.4	9.8	74.9	84.7
(n=100)	CI	4.4–17.0	2.8–13.9	5.2–17.6	65.0–82.8	75.6–90.9
8th	%	2.2	2.9	8.7	86.3	94.9
(n=1,881)	CI	1.6–2.9	2.2–3.7	7.5–10.0	84.7–87.8	93.9–95.9
10th	%	3.5	5.6	12.9	77.9	90.9
(n=1,264)	CI	2.7–4.7	4.5–7.0	11.2–14.9	75.6–80.1	89.1–92.3
12th	%	4.5	5.1	15.0	75.3	90.3
(n=1,103)	CI	3.4–5.9	4.0–6.6	13.0–17.2	72.7–77.8	88.4–91.9

Note: The confidence interval (CI) is 95 percent.

Table BH5.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.7	9.6	19.9	63.9	83.8
(n=4,263)	CI	5.9–7.5	8.7–10.5	18.7–21.2	62.4–65.4	82.6–84.9
Male	%	8.6	9.2	19.9	62.4	82.2
(n=2,066)	CI	7.4–9.9	8.0–10.6	18.1–21.7	60.2–64.5	80.4–83.9
Female	%	3.7	9.7	20.0	66.5	86.5
(n=2,071)	CI	3.0–4.7	8.5–11.2	18.2–21.8	64.4–68.6	84.9–88.0
White	%	6.4	9.7	20.0	63.8	83.9
(n=3,456)	CI	5.6–7.3	8.8–10.8	18.7–21.5	62.2–65.5	82.6–85.1
Black	%	9.1	10.5	15.9	64.5	80.4
(n=132)	CI	5.1–15.9	6.1–17.5	10.4–23.4	55.6–72.5	72.2–86.6
Hispanic	%	8.9	11.2	16.4	63.5	79.9
(n=103)	CI	4.5–17.1	6.3–19.0	10.1–25.4	53.3–72.6	70.6–86.8
8th	%	2.5	4.9	13.8	78.7	92.5
(n=1,888)	CI	1.9–3.4	4.0–6.0	12.3–15.5	76.8–80.5	91.3–93.6
10th	%	7.0	11.8	21.7	59.6	81.2
(n=1,269)	CI	5.7–8.5	10.1–13.7	19.5–24.0	56.9–62.2	79.0–83.3
12th	%	10.8	12.3	24.6	52.3	76.9
(n=1,106)	CI	9.1–12.8	10.4–14.3	22.1–27.2	49.4–55.3	74.4–79.3

Note: The confidence interval (CI) is 95 percent.

Table BH5.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	8.1	9.9	18.8	63.2	82.0
(n=4,263)	CI	7.2–9.0	9.0–10.9	17.6–20.1	61.7–64.7	80.8–83.2
Male	%	10.4	10.0	18.2	61.3	79.6
(n=2,066)	CI	9.1–11.9	8.7–11.4	16.6–20.0	59.1–63.5	77.7–81.4
Female	%	4.8	9.5	19.5	66.2	85.8
(n=2,071)	CI	3.9–5.9	8.2–10.9	17.8–21.3	64.1–68.3	84.1–87.3
White	%	8.0	10.0	19.3	62.7	82.0
(n=3,458)	CI	7.1–9.0	9.0–11.1	18.0–20.7	61.1–64.4	80.7–83.4
Black	%	8.2	10.1	21.1	60.6	81.7
(n=132)	CI	4.4–14.7	5.8–17.1	14.8–29.1	51.7–68.9	73.6–87.7
Hispanic	%	11.5	9.4	16.2	63.0	79.2
(n=102)	CI	6.3–19.8	4.8–17.3	10.0–25.1	52.8–72.2	69.7–86.3
8th	%	2.6	4.1	12.1	81.1	93.3
(n=1,885)	CI	2.0–3.4	3.3–5.1	10.7–13.7	79.3–82.8	92.0–94.3
10th	%	7.4	11.2	21.4	60.0	81.4
(n=1,270)	CI	6.1–8.9	9.6–13.1	19.2–23.8	57.3–62.7	79.2–83.5
12th	%	14.7	14.7	23.3	47.2	70.5
(n=1,108)	CI	12.8–17.0	12.8–17.0	20.9–25.9	44.3–50.2	67.8–73.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	14.5	17.1	17.3	51.1	68.4
(n=4,262)	CI	13.4–15.6	16.0–18.3	16.1–18.5	49.6–52.6	66.9–69.8
Male	%	16.6	17.1	15.9	50.3	66.2
(n=2,064)	CI	15.0–18.4	15.5–18.9	14.3–17.6	48.1–52.5	64.1–68.3
Female	%	11.4	16.8	19.0	52.7	71.7
(n=2,073)	CI	10.1–13.0	15.2–18.6	17.3–20.8	50.5–54.9	69.7–73.7
White	%	14.9	17.5	17.6	50.0	67.6
(n=3,456)	CI	13.7–16.2	16.2–18.9	16.3–18.9	48.3–51.7	66.0–69.2
Black	%	12.3	21.0	18.6	48.1	66.7
(n=132)	CI	7.5–19.5	14.7–29.1	12.7–26.5	39.5–56.8	57.9–74.5
Hispanic	%	14.1	11.4	15.6	59.0	74.6
(n=103)	CI	8.2–22.9	6.4–19.3	9.5–24.5	48.8–68.4	64.8–82.4
8th	%	6.1	10.7	13.6	69.6	83.2
(n=1,886)	CI	5.1–7.3	9.4–12.2	12.1–15.2	67.5–71.7	81.4–84.8
10th	%	15.2	19.6	19.6	45.6	65.2
(n=1,268)	CI	13.3–17.2	17.5–21.9	17.5–21.9	42.8–48.3	62.5–67.8
12th	%	22.8	21.4	18.8	37.0	55.8
(n=1,108)	CI	20.5–25.4	19.1–23.9	16.6–21.2	34.2–39.8	52.8–58.7

Note: The confidence interval (CI) is 95 percent.

Table BH5.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	13.8	12.5	14.5	59.2	73.7
(n=4,254)	CI	12.7–14.9	11.5–13.6	13.5–15.7	57.6–60.7	72.3–75.1
Male	%	15.1	11.8	15.4	57.7	73.1
(n=2,062)	CI	13.5–16.8	10.4–13.4	13.8–17.1	55.5–59.9	71.0–75.1
Female	%	11.5	12.8	14.0	61.7	75.7
(n=2,066)	CI	10.1–13.0	11.4–14.4	12.5–15.6	59.5–63.9	73.7–77.6
White	%	13.5	12.7	14.6	59.1	73.7
(n=3,450)	CI	12.4–14.8	11.6–14.0	13.4–15.9	57.4–60.8	72.1–75.2
Black	%	20.4	14.2	14.2	51.2	65.3
(n=132)	CI	14.0–28.7	9.0–21.7	9.1–21.4	42.4–59.8	56.4–73.3
Hispanic	%	14.2	4.6	16.5	64.7	81.1
(n=102)	CI	8.3–23.2	1.9–11.0	10.2–25.5	54.4–73.7	71.8–87.9
8th	%	3.7	5.1	10.0	81.1	91.1
(n=1,885)	CI	2.9–4.7	4.2–6.2	8.7–11.5	79.3–82.8	89.8–92.3
10th	%	16.1	14.3	15.5	54.1	69.6
(n=1,266)	CI	14.2–18.3	12.5–16.4	13.6–17.6	51.3–56.8	67.0–72.0
12th	%	22.3	18.5	18.4	40.7	59.2
(n=1,103)	CI	20.0–24.9	16.3–20.9	16.2–20.8	37.9–43.7	56.2–62.0

Note: The confidence interval (CI) is 95 percent.

Table BH5.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	64.5	24.7	10.8	35.5
(n=4,247)	CI	63.0–66.0	23.4–26.0	9.9–11.8	34.0–37.0
Male	%	64.7	24.2	11.2	35.3
(n=2,056)	CI	62.5–66.8	22.3–26.1	9.8–12.6	33.2–37.5
Female	%	64.1	26.0	9.9	35.9
(n=2,065)	CI	62.0–66.2	24.1–27.9	8.7–11.3	33.8–38.0
White	%	64.5	24.9	10.5	35.5
(n=3,444)	CI	62.9–66.2	23.5–26.4	9.5–11.6	33.8–37.1
Black	%	66.0	18.1	15.9	34.0
(n=130)	CI	57.2–73.9	12.3–25.9	10.3–23.5	26.1–42.8
Hispanic	%	64.5	22.5	13.0	35.5
(n=103)	CI	54.4–73.5	15.1–32.1	7.6–21.4	26.5–45.6
8th	%	62.4	26.7	10.9	37.6
(n=1,886)	CI	60.2–64.6	24.8–28.8	9.5–12.4	35.4–39.8
10th	%	63.4	24.6	12.1	36.6
(n=1,264)	CI	60.7–66.0	22.3–27.0	10.4–14.0	34.0–39.3
12th	%	68.0	22.5	9.4	32.0
(n=1,097)	CI	65.2–70.7	20.1–25.1	7.9–11.3	29.3–34.8

Note: The confidence interval (CI) is 95 percent.

Table BH5.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	43.0	36.4	20.6	57.0
(n=4,238)	CI	41.5–44.5	35.0–37.9	19.4–21.8	55.5–58.5
Male	%	44.0	35.9	20.1	56.0
(n=2,054)	CI	41.8–46.2	33.8–38.0	18.4–22.0	53.8–58.2
Female	%	41.5	37.8	20.6	58.5
(n=2,058)	CI	39.4–43.7	35.7–40.0	18.9–22.5	56.3–60.6
White	%	42.6	37.0	20.4	57.4
(n=3,438)	CI	40.9–44.3	35.4–38.7	19.1–21.8	55.7–59.1
Black	%	48.4	25.7	25.9	51.6
(n=130)	CI	39.7–57.2	18.7–34.2	18.9–34.4	42.8–60.3
Hispanic	%	45.9	40.7	13.4	54.1
(n=102)	CI	36.2–56.0	31.3–50.9	7.9–21.7	44.0–63.8
8th	%	42.3	37.4	20.3	57.7
(n=1,880)	CI	40.1–44.5	35.2–39.6	18.6–22.2	55.5–59.9
10th	%	42.1	36.1	21.8	57.9
(n=1,262)	CI	39.4–44.8	33.5–38.8	19.6–24.1	55.2–60.6
12th	%	44.8	35.6	19.5	55.2
(n=1,096)	CI	41.9–47.8	32.9–38.5	17.3–22.0	52.2–58.1

Note: The confidence interval (CI) is 95 percent.

Table BH5.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	37.4	32.4	30.2	62.6
(n=4,243)	CI	35.9–38.9	31.0–33.9	28.8–31.6	61.1–64.1
Male	%	37.9	31.1	31.0	62.1
(n=2,052)	CI	35.8–40.1	29.1–33.2	29.0–33.0	59.9–64.2
Female	%	36.5	34.3	29.3	63.5
(n=2,066)	CI	34.4–38.6	32.2–36.4	27.3–31.3	61.4–65.6
White	%	37.3	33.0	29.7	62.7
(n=3,441)	CI	35.7–39.0	31.4–34.6	28.2–31.3	61.0–64.3
Black	%	37.1	28.5	34.4	62.9
(n=129)	CI	29.0–46.0	21.3–37.1	26.5–43.2	54.0–71.0
Hispanic	%	43.0	31.9	25.1	57.0
(n=103)	CI	33.5–53.0	23.3–41.9	17.5–34.7	47.0–66.5
8th	%	33.6	32.1	34.3	66.4
(n=1,884)	CI	31.5–35.8	30.0–34.2	32.2–36.5	64.2–68.5
10th	%	37.0	33.3	29.7	63.0
(n=1,264)	CI	34.4–39.7	30.7–35.9	27.3–32.3	60.3–65.6
12th	%	41.9	32.0	26.1	58.1
(n=1,095)	CI	39.0–44.8	29.3–34.8	23.6–28.8	55.2–61.0

Note: The confidence interval (CI) is 95 percent.

Behavioral Health Planning Region 6 Results Summary

Table BH6.1. Lifetime Prescription Drug Misuse

		On how many occasions (if any) have you ever used prescription drugs not prescribed to you?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	86.7	6.7	2.5	1.0	1.2	0.7	1.2	13.3
(n=3,424)	CI	85.5–87.8	5.9–7.6	2.0–3.1	0.7–1.3	0.9–1.7	0.5–1.1	0.9–1.7	12.2–14.5
Male	%	90.0	4.2	1.7	0.6	1.2	0.9	1.4	10.0
(n=1,668)	CI	88.4–91.4	3.3–5.3	1.1–2.4	0.4–1.2	0.8–1.9	0.5–1.5	0.9–2.1	8.6–11.6
Female	%	84.3	8.6	3.4	1.3	1.3	0.5	0.8	15.7
(n=1,649)	CI	82.4–86.0	7.3–10.1	2.6–4.4	0.8–1.9	0.8–2.0	0.2–0.9	0.4–1.4	14.0–17.6
White	%	86.6	7.1	2.4	0.9	1.4	0.8	0.9	13.4
(n=2,341)	CI	85.1–87.9	6.1–8.3	1.8–3.1	0.6–1.4	0.9–2.0	0.5–1.3	0.5–1.3	12.1–14.9
Black	%	90.5	4.6	2.1	0.4	0.3	0.5	1.6	9.5
(n=574)	CI	87.8–92.7	3.1–6.7	1.1–3.7	0.1–1.5	0.1–1.3	0.2–1.7	0.8–3.0	7.3–12.2
Hispanic	%	93.7	3.0	1.8	0.0	1.5	0.0	0.0	6.3
(n=66)	CI	84.0–97.7	0.7–11.6	0.2–12.3	–	0.2–10.4	–	–	2.3–16.0
8th	%	91.8	5.0	1.5	0.8	0.4	0.2	0.5	8.2
(n=1,302)	CI	90.2–93.2	3.9–6.3	0.9–2.3	0.4–1.4	0.2–0.9	0.0–0.6	0.2–1.0	6.8–9.8
10th	%	84.5	8.0	3.0	0.9	1.0	1.0	1.6	15.5
(n=1,189)	CI	82.3–86.4	6.6–9.7	2.2–4.2	0.5–1.7	0.6–1.8	0.6–1.8	1.0–2.5	13.6–17.7
12th	%	83.9	7.0	3.0	1.2	2.3	1.0	1.6	16.1
(n=933)	CI	81.4–86.1	5.6–8.9	2.1–4.3	0.7–2.1	1.5–3.5	0.5–1.8	1.0–2.6	13.9–18.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.2. Lifetime Alcohol Use

		On how many occasions (if any) have you ever had one or more drinks of an alcoholic beverage?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	54.1	20.9	8.6	4.9	4.5	2.7	4.3	45.9
(n=3,459)	CI	52.5–55.8	19.6–22.3	7.7–9.6	4.2–5.7	3.9–5.3	2.2–3.3	3.6–5.0	44.2–47.5
Male	%	57.3	18.9	6.6	4.6	4.3	2.6	5.6	42.7
(n=1,689)	CI	54.9–59.7	17.1–20.9	5.5–7.9	3.7–5.7	3.4–5.4	2.0–3.6	4.5–6.9	40.3–45.1
Female	%	52.1	22.5	10.0	5.2	4.8	2.9	2.6	47.9
(n=1,661)	CI	49.6–54.5	20.6–24.6	8.6–11.6	4.2–6.4	3.9–6.0	2.1–3.8	1.9–3.5	45.5–50.4
White	%	51.9	21.8	8.2	5.2	4.8	3.3	4.7	48.1
(n=2,361)	CI	49.9–54.0	20.2–23.5	7.1–9.4	4.4–6.2	4.0–5.8	2.7–4.2	3.9–5.7	46.0–50.1
Black	%	68.7	16.9	6.5	2.7	2.6	0.7	1.9	31.3
(n=582)	CI	64.8–72.4	14.1–20.2	4.7–8.8	1.6–4.4	1.5–4.3	0.3–2.0	1.0–3.5	27.6–35.2
Hispanic	%	61.3	17.2	8.6	4.7	3.1	1.8	3.3	38.7
(n=67)	CI	48.8–72.6	9.6–28.8	3.8–18.2	1.5–14.1	0.7–12.1	0.2–12.1	0.8–12.6	27.4–51.2
8th	%	68.5	20.5	6.1	2.8	0.9	0.5	0.6	31.5
(n=1,314)	CI	65.9–71.0	18.4–22.8	4.9–7.5	2.0–3.9	0.5–1.6	0.3–1.1	0.3–1.2	29.0–34.1
10th	%	48.3	24.8	10.1	5.7	5.6	2.8	2.7	51.7
(n=1,202)	CI	45.5–51.1	22.5–27.3	8.5–12.0	4.5–7.2	4.4–7.0	2.0–3.9	1.9–3.7	48.9–54.5
12th	%	45.5	17.1	9.5	6.1	7.2	4.9	9.7	54.5
(n=943)	CI	42.4–48.7	14.8–19.6	7.8–11.5	4.8–7.8	5.7–9.0	3.7–6.4	8.0–11.8	51.3–57.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.3. Lifetime Binge Drinking

		On how many occasions (if any) have you ever <u>had five or more drinks on the same occasion?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.1	7.8	4.4	2.8	2.3	1.5	2.0	20.9
(n=3,449)	CI	77.7–80.5	7.0–8.8	3.8–5.2	2.3–3.4	1.8–2.9	1.1–2.0	1.5–2.5	19.5–22.3
Male	%	78.6	7.1	4.2	3.0	2.5	1.9	2.7	21.4
(n=1,681)	CI	76.5–80.6	5.9–8.4	3.3–5.3	2.2–3.9	1.9–3.5	1.3–2.8	2.0–3.6	19.4–23.5
Female	%	80.1	8.5	4.7	2.7	2.0	1.1	1.0	19.9
(n=1,659)	CI	78.0–82.0	7.2–9.9	3.7–5.8	2.0–3.6	1.4–2.8	0.7–1.8	0.6–1.7	18.0–22.0
White	%	76.9	8.2	5.1	3.1	2.8	1.8	2.2	23.1
(n=2,355)	CI	75.1–78.6	7.1–9.4	4.2–6.1	2.5–3.9	2.2–3.6	1.3–2.4	1.7–3.0	21.4–24.9
Black	%	89.5	5.2	1.9	1.6	0.6	0.3	1.0	10.5
(n=580)	CI	86.6–91.8	3.6–7.3	1.0–3.5	0.8–3.0	0.2–1.8	0.1–1.3	0.4–2.3	8.2–13.4
Hispanic	%	82.4	8.0	2.8	1.8	0.0	3.3	1.8	17.6
(n=67)	CI	70.6–90.1	3.3–18.2	0.7–10.9	0.2–12.1	–	0.8–12.6	0.2–12.1	9.9–29.4
8th	%	92.8	4.4	1.3	0.8	0.3	0.2	0.2	7.2
(n=1,307)	CI	91.3–94.1	3.4–5.7	0.8–2.1	0.4–1.4	0.1–0.8	0.1–0.7	0.0–0.6	5.9–8.7
10th	%	79.0	8.6	5.1	3.6	1.7	1.0	1.0	21.0
(n=1,199)	CI	76.6–81.2	7.1–10.3	4.0–6.5	2.7–4.8	1.1–2.7	0.6–1.8	0.6–1.8	18.8–23.4
12th	%	65.2	10.6	7.0	4.1	5.0	3.4	4.9	34.8
(n=943)	CI	62.1–68.1	8.8–12.7	5.5–8.8	3.0–5.6	3.7–6.5	2.4–4.7	3.7–6.4	31.9–37.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.4. Lifetime Cigarette Use

		On how many occasions (if any) have you ever <u>smoked all or part of a cigarette?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	79.5	9.3	3.2	1.8	1.5	1.0	3.7	20.5
(n=3,452)	CI	78.1–80.8	8.4–10.3	2.6–3.8	1.4–2.4	1.2–2.0	0.7–1.4	3.1–4.4	19.2–21.9
Male	%	78.8	8.7	3.4	1.7	1.8	0.7	4.9	21.2
(n=1,685)	CI	76.8–80.7	7.4–10.2	2.6–4.4	1.2–2.5	1.2–2.5	0.4–1.3	3.9–6.0	19.3–23.2
Female	%	80.8	9.5	3.1	1.9	1.3	1.1	2.3	19.2
(n=1,658)	CI	78.8–82.6	8.2–11.0	2.4–4.1	1.3–2.7	0.9–2.0	0.7–1.7	1.6–3.2	17.4–21.2
White	%	77.3	9.4	3.7	2.0	1.9	1.1	4.6	22.7
(n=2,358)	CI	75.5–79.0	8.3–10.6	3.0–4.6	1.5–2.7	1.4–2.5	0.8–1.7	3.8–5.6	21.0–24.5
Black	%	91.5	5.3	1.0	0.8	0.4	0.2	0.8	8.5
(n=578)	CI	88.9–93.5	3.8–7.5	0.5–2.3	0.3–2.0	0.1–1.6	0.0–1.1	0.3–2.1	6.5–11.1
Hispanic	%	77.4	14.8	0.0	2.8	1.8	0.0	3.3	22.6
(n=67)	CI	65.4–86.1	8.0–25.8	–	0.7–10.9	0.2–12.1	–	0.8–12.6	13.9–34.6
8th	%	85.6	8.9	2.2	1.3	0.6	0.8	0.6	14.4
(n=1,313)	CI	83.6–87.4	7.5–10.6	1.5–3.2	0.8–2.1	0.3–1.2	0.4–1.4	0.3–1.2	12.6–16.4
10th	%	79.3	10.0	3.3	1.8	1.7	0.2	3.7	20.7
(n=1,202)	CI	76.9–81.5	8.4–11.8	2.4–4.5	1.2–2.8	1.1–2.7	0.1–0.8	2.7–4.9	18.5–23.1
12th	%	73.3	9.0	4.0	2.4	2.2	1.9	7.0	26.7
(n=937)	CI	70.4–76.1	7.4–11.0	3.0–5.5	1.6–3.7	1.5–3.4	1.2–3.0	5.5–8.8	23.9–29.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.5. Lifetime Electronic Cigarette Use

		On how many occasions (if any) have you ever <u>used an e-cigarette</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	68.4	9.8	4.9	2.4	3.3	2.5	8.7	31.6
(n=3,446)	CI	66.8–70.0	8.9–10.9	4.2–5.7	1.9–3.0	2.7–4.0	2.0–3.1	7.8–9.7	30.0–33.2
Male	%	67.2	10.4	4.5	2.0	3.1	2.7	10.2	32.8
(n=1,684)	CI	64.8–69.4	9.0–11.9	3.6–5.6	1.5–2.9	2.3–4.0	2.0–3.7	8.8–11.8	30.6–35.2
Female	%	70.1	9.1	5.5	2.6	3.7	2.2	6.9	29.9
(n=1,654)	CI	67.8–72.3	7.8–10.6	4.4–6.7	1.9–3.5	2.8–4.7	1.6–3.1	5.7–8.3	27.7–32.2
White	%	64.0	10.8	5.8	2.6	3.8	2.9	10.2	36.0
(n=2,355)	CI	62.0–65.9	9.6–12.1	4.9–6.8	2.0–3.3	3.0–4.6	2.3–3.7	9.0–11.6	34.1–38.0
Black	%	87.0	5.4	1.6	1.2	1.2	0.5	3.1	13.0
(n=576)	CI	84.0–89.6	3.8–7.5	0.8–3.1	0.6–2.4	0.6–2.6	0.2–1.7	1.9–4.9	10.4–16.0
Hispanic	%	77.8	4.7	3.0	3.2	1.8	4.5	5.1	22.2
(n=66)	CI	65.5–86.6	1.5–13.9	0.7–11.6	0.8–12.3	0.2–12.3	1.4–13.5	1.6–15.1	13.4–34.5
8th	%	80.9	9.0	2.9	1.5	2.1	1.3	2.4	19.1
(n=1,307)	CI	78.6–82.9	7.5–10.6	2.1–4.0	1.0–2.4	1.4–3.0	0.8–2.1	1.7–3.4	17.1–21.4
10th	%	64.2	11.8	6.7	3.3	3.3	2.4	8.3	35.8
(n=1,196)	CI	61.5–66.9	10.1–13.7	5.4–8.2	2.5–4.5	2.4–4.4	1.7–3.5	6.9–10.0	33.1–38.5
12th	%	60.0	8.7	5.1	2.3	4.6	3.7	15.6	40.0
(n=943)	CI	56.8–63.1	7.0–10.6	3.8–6.7	1.5–3.5	3.5–6.2	2.7–5.1	13.4–18.1	36.9–43.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.6. Lifetime Smokeless Tobacco Use

		On how many occasions (if any) have you ever <u>used smokeless tobacco</u> ?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	87.7	4.6	1.4	1.2	1.0	0.7	3.5	12.3
(n=3,444)	CI	86.5–88.7	3.9–5.3	1.1–1.9	0.9–1.6	0.7–1.4	0.5–1.1	2.9–4.2	11.3–13.5
Male	%	81.7	5.9	2.0	1.7	1.6	1.1	6.0	18.3
(n=1,684)	CI	79.7–83.5	4.9–7.2	1.5–2.9	1.1–2.4	1.0–2.3	0.7–1.7	5.0–7.3	16.5–20.3
Female	%	94.3	2.9	0.9	0.7	0.3	0.4	0.6	5.7
(n=1,652)	CI	93.1–95.4	2.1–3.8	0.5–1.5	0.4–1.2	0.1–0.7	0.2–0.9	0.3–1.2	4.6–6.9
White	%	85.6	5.2	1.5	1.4	1.0	0.8	4.5	14.4
(n=2,349)	CI	84.1–87.0	4.4–6.2	1.1–2.1	1.0–2.0	0.6–1.5	0.5–1.3	3.7–5.4	13.0–15.9
Black	%	95.8	1.9	0.7	0.4	0.6	0.0	0.6	4.2
(n=579)	CI	93.8–97.2	1.1–3.4	0.3–1.8	0.1–1.6	0.2–1.9	–	0.2–1.8	2.8–6.2
Hispanic	%	96.7	0.0	0.0	0.0	0.0	0.0	3.3	3.3
(n=67)	CI	87.4–99.2	–	–	–	–	–	0.8–12.6	0.8–12.6
8th	%	93.0	3.3	1.0	0.9	0.2	0.5	1.1	7.0
(n=1,307)	CI	91.5–94.3	2.4–4.4	0.6–1.7	0.5–1.6	0.1–0.7	0.2–1.0	0.6–1.8	5.7–8.5
10th	%	87.6	4.5	1.7	1.1	1.3	0.9	2.8	12.4
(n=1,197)	CI	85.6–89.3	3.5–5.8	1.1–2.7	0.6–1.9	0.8–2.2	0.5–1.6	2.0–3.9	10.7–14.4
12th	%	82.2	5.9	1.6	1.6	1.4	0.7	6.6	17.8
(n=940)	CI	79.6–84.5	4.6–7.6	1.0–2.6	1.0–2.6	0.8–2.4	0.4–1.5	5.2–8.3	15.5–20.4

Note: The confidence interval (CI) is 95 percent.

Table BH6.7. Lifetime Marijuana Use

		On how many occasions (if any) have you ever <u>used marijuana or hashish?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	75.3	7.1	4.1	2.5	2.9	1.5	6.6	24.7
(n=3,453)	CI	73.8–76.7	6.3–8.0	3.5–4.8	2.0–3.1	2.4–3.6	1.2–2.0	5.8–7.5	23.3–26.2
Male	%	75.4	6.2	3.2	2.8	3.1	1.4	7.9	24.6
(n=1,692)	CI	73.3–77.5	5.1–7.5	2.4–4.1	2.1–3.8	2.3–4.0	0.9–2.1	6.7–9.4	22.5–26.7
Female	%	75.8	8.0	4.8	2.2	2.7	1.5	4.9	24.2
(n=1,653)	CI	73.7–77.9	6.8–9.4	3.9–6.0	1.6–3.1	2.0–3.7	1.0–2.2	3.9–6.1	22.1–26.3
White	%	76.6	7.2	3.6	2.3	2.9	1.2	6.2	23.4
(n=2,356)	CI	74.8–78.3	6.2–8.3	2.9–4.5	1.7–3.0	2.2–3.6	0.9–1.8	5.3–7.4	21.7–25.2
Black	%	76.3	5.7	4.2	2.7	2.6	1.5	7.0	23.7
(n=582)	CI	72.6–79.7	4.1–7.9	2.8–6.2	1.6–4.4	1.5–4.4	0.8–2.8	5.1–9.5	20.3–27.4
Hispanic	%	76.7	7.9	5.8	3.1	1.5	3.6	1.5	23.3
(n=67)	CI	64.6–85.7	3.2–17.9	2.1–14.8	0.7–12.1	0.2–10.2	0.9–13.5	0.2–10.2	14.3–35.4
8th	%	89.1	5.0	2.1	0.5	1.0	0.7	1.6	10.9
(n=1,310)	CI	87.3–90.7	4.0–6.4	1.4–3.0	0.3–1.1	0.6–1.7	0.4–1.3	1.0–2.4	9.3–12.7
10th	%	72.5	7.6	5.5	3.4	3.1	2.0	6.0	27.5
(n=1,201)	CI	69.9–75.0	6.2–9.2	4.3–6.9	2.5–4.6	2.2–4.2	1.3–3.0	4.8–7.5	25.0–30.1
12th	%	64.1	8.7	4.7	3.5	4.8	1.9	12.5	35.9
(n=942)	CI	60.9–67.1	7.0–10.6	3.5–6.2	2.5–4.9	3.6–6.3	1.2–3.0	10.5–14.7	32.9–39.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.8. Lifetime Methamphetamines Use

		On how many occasions (if any) have you ever <u>used methamphetamines (crank, etc.)?</u>							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	98.5	0.5	0.2	0.3	0.2	0.0	0.3	1.5
(n=3,450)	CI	98.0–98.9	0.3–0.9	0.1–0.4	0.1–0.5	0.1–0.4	–	0.2–0.6	1.1–2.0
Male	%	98.4	0.7	0.1	0.3	0.1	0.0	0.3	1.6
(n=1,687)	CI	97.6–98.9	0.4–1.3	0.0–0.5	0.1–0.8	0.0–0.5	–	0.1–0.8	1.1–2.4
Female	%	99.0	0.3	0.3	0.2	0.1	0.0	0.2	1.0
(n=1,654)	CI	98.4–99.4	0.1–0.7	0.1–0.7	0.1–0.6	0.0–0.4	–	0.1–0.6	0.6–1.6
White	%	98.9	0.4	0.1	0.3	0.1	0.0	0.2	1.1
(n=2,359)	CI	98.4–99.3	0.2–0.8	0.0–0.3	0.1–0.6	0.0–0.4	–	0.1–0.5	0.7–1.6
Black	%	98.2	0.4	0.5	0.5	0.0	0.0	0.4	1.8
(n=576)	CI	96.6–99.0	0.1–1.5	0.2–1.7	0.2–1.7	–	–	0.1–1.5	1.0–3.4
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=66)	CI	–	–	–	–	–	–	–	–
8th	%	99.2	0.3	0.2	0.2	0.0	0.0	0.2	0.8
(n=1,310)	CI	98.5–99.5	0.1–0.8	0.1–0.7	0.0–0.6	–	–	0.0–0.6	0.5–1.5
10th	%	98.1	0.7	0.2	0.3	0.3	0.0	0.3	1.9
(n=1,202)	CI	97.1–98.7	0.3–1.3	0.1–0.8	0.1–0.9	0.1–0.9	–	0.1–0.9	1.3–2.9
12th	%	98.3	0.6	0.1	0.3	0.2	0.0	0.4	1.7
(n=938)	CI	97.2–99.0	0.3–1.4	0.0–0.8	0.1–1.0	0.1–0.8	–	0.2–1.1	1.0–2.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.9. Lifetime Other Illegal Drug Use

		On how many occasions (if any) have you ever used any other illegal drugs (cocaine, heroin, inhalants, etc.)?							
		Never	1–2 times	3–5 times	6–9 times	10–19 times	20–39 times	40+ times	Any Use
Total	%	96.6	1.8	0.3	0.3	0.3	0.2	0.5	3.4
(n=3,453)	CI	96.0–97.2	1.4–2.3	0.2–0.6	0.1–0.5	0.1–0.5	0.1–0.4	0.3–0.9	2.8–4.0
Male	%	96.7	1.5	0.5	0.3	0.3	0.2	0.6	3.3
(n=1,686)	CI	95.7–97.5	1.0–2.2	0.2–1.0	0.1–0.7	0.1–0.7	0.1–0.5	0.3–1.2	2.5–4.3
Female	%	97.0	1.9	0.1	0.3	0.3	0.1	0.3	3.0
(n=1,658)	CI	96.0–97.7	1.4–2.7	0.0–0.5	0.1–0.8	0.1–0.7	0.0–0.5	0.1–0.7	2.3–4.0
White	%	96.7	2.0	0.3	0.2	0.3	0.0	0.4	3.3
(n=2,357)	CI	95.9–97.4	1.5–2.6	0.2–0.7	0.1–0.6	0.1–0.6	0.0–0.3	0.2–0.8	2.6–4.1
Black	%	97.7	0.7	0.2	0.4	0.2	0.0	0.9	2.3
(n=578)	CI	96.0–98.6	0.3–1.8	0.0–1.4	0.1–1.5	0.0–1.4	–	0.4–2.1	1.4–4.0
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	98.2	1.4	0.1	0.1	0.0	0.0	0.2	1.8
(n=1,313)	CI	97.3–98.8	0.9–2.3	0.0–0.5	0.0–0.5	–	–	0.1–0.7	1.2–2.7
10th	%	96.1	2.2	0.2	0.2	0.3	0.2	0.7	3.9
(n=1,203)	CI	94.8–97.1	1.5–3.2	0.0–0.7	0.1–0.8	0.1–0.9	0.1–0.8	0.3–1.3	2.9–5.2
12th	%	95.6	1.7	0.7	0.5	0.4	0.2	0.7	4.4
(n=937)	CI	94.1–96.8	1.0–2.8	0.4–1.6	0.2–1.3	0.2–1.1	0.1–0.8	0.4–1.6	3.2–5.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.10. Age of First Prescription Drug Misuse

How old were you when you first used prescription drugs not prescribed to you?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	88.0	1.6	0.7	1.3	1.7	1.9	2.2	1.5	0.9	13.58
(n=3,459)	CI	86.9–89.1	1.3–2.1	0.5–1.0	1.0–1.8	1.3–2.2	1.5–2.5	1.8–2.8	1.2–2.1	0.6–1.3	13.4–13.8
Male	%	91.5	1.4	0.6	1.0	0.8	1.3	1.7	1.2	0.5	13.45
(n=1,688)	CI	90.1–92.8	1.0–2.1	0.3–1.0	0.6–1.6	0.5–1.3	0.8–2.0	1.1–2.4	0.8–2.0	0.2–1.0	13.1–13.8
Female	%	85.3	1.7	0.9	1.4	2.5	2.4	2.7	1.8	1.4	13.72
(n=1,662)	CI	83.5–86.9	1.2–2.4	0.5–1.4	0.9–2.1	1.8–3.3	1.7–3.2	2.0–3.7	1.3–2.7	0.9–2.1	13.5–14.0
White	%	87.9	1.7	0.8	1.0	1.7	1.9	2.4	1.6	0.9	13.64
(n=2,363)	CI	86.5–89.2	1.3–2.3	0.5–1.2	0.7–1.5	1.2–2.3	1.4–2.5	1.8–3.2	1.2–2.3	0.6–1.5	13.4–13.9
Black	%	91.5	1.3	0.5	1.5	0.7	0.7	1.6	0.9	1.2	13.61
(n=580)	CI	88.9–93.6	0.6–2.6	0.2–1.6	0.8–3.0	0.2–1.7	0.3–2.0	0.8–3.0	0.4–2.3	0.5–2.7	12.9–14.3
Hispanic	%	97.0	0.0	0.0	0.0	0.0	0.0	3.0	0.0	0.0	15.00
(n=66)	CI	88.4–99.3	–	–	–	–	–	0.7–11.6	–	–	15.0–15.0
8th	%	93.0	1.9	0.9	1.7	2.0	0.5	0.0	0.0	0.0	11.76
(n=1,313)	CI	91.5–94.3	1.3–2.8	0.5–1.6	1.1–2.5	1.4–2.9	0.3–1.1	–	–	–	11.5–12.0
10th	%	85.9	2.1	1.0	2.0	2.1	3.2	3.3	0.3	0.1	13.07
(n=1,203)	CI	83.8–87.8	1.4–3.1	0.6–1.7	1.3–3.0	1.4–3.1	2.4–4.4	2.4–4.5	0.1–0.9	0.0–0.6	12.8–13.3
12th	%	85.2	0.8	0.2	0.3	1.0	2.0	3.4	4.4	2.6	14.99
(n=943)	CI	82.8–87.3	0.4–1.7	0.1–0.8	0.1–1.0	0.5–1.8	1.3–3.1	2.4–4.7	3.3–6.0	1.8–3.9	14.7–15.3

Note: The confidence interval (CI) is 95 percent.

Table BH6.11. Age of First Alcohol Use

How old were you when you first had more than a sip or two of beer, wine, or hard liquor (vodka, whiskey, or gin)?											
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	58.1	5.5	2.5	4.0	6.5	6.3	8.2	4.6	4.2	13.67
(n=3,448)	CI	56.4–59.8	4.8–6.4	2.0–3.1	3.4–4.7	5.7–7.4	5.5–7.2	7.3–9.2	4.0–5.5	3.6–5.0	13.6–13.8
Male	%	61.0	5.3	2.8	4.0	6.0	5.3	7.3	4.1	4.2	13.60
(n=1,687)	CI	58.7–63.4	4.3–6.5	2.1–3.7	3.2–5.0	5.0–7.3	4.3–6.5	6.1–8.7	3.2–5.2	3.3–5.3	13.4–13.8
Female	%	56.2	5.1	2.1	3.7	6.8	7.2	9.3	5.3	4.2	13.81
(n=1,654)	CI	53.8–58.6	4.2–6.3	1.5–3.0	2.9–4.7	5.7–8.1	6.1–8.6	8.0–10.9	4.3–6.6	3.3–5.4	13.7–14.0
White	%	56.0	5.4	2.5	3.8	7.0	6.8	8.6	5.1	4.8	13.75
(n=2,355)	CI	54.0–58.0	4.6–6.4	2.0–3.3	3.1–4.6	6.1–8.1	5.8–7.9	7.5–9.8	4.2–6.1	3.9–5.8	13.6–13.9
Black	%	72.4	3.0	1.7	2.7	3.6	3.6	5.9	3.8	3.4	13.90
(n=581)	CI	68.5–75.9	1.9–4.8	0.9–3.2	1.6–4.3	2.4–5.4	2.3–5.4	4.2–8.1	2.4–5.8	2.1–5.4	13.6–14.3
Hispanic	%	62.5	1.3	0.0	10.0	7.1	4.4	10.0	3.0	1.8	13.71
(n=66)	CI	49.8–73.6	0.2–9.4	–	4.5–20.8	2.9–16.3	1.4–13.1	4.5–20.8	0.7–11.7	0.2–12.3	13.0–14.4
8th	%	72.7	6.5	3.1	4.8	9.7	2.9	0.2	0.0	0.0	12.00
(n=1,313)	CI	70.3–75.1	5.3–8.0	2.3–4.2	3.8–6.1	8.2–11.4	2.1–4.0	0.1–0.7	–	–	11.9–12.1
10th	%	52.9	6.1	2.1	4.5	6.2	10.8	14.0	3.2	0.3	13.48
(n=1,196)	CI	50.0–55.7	4.9–7.6	1.4–3.1	3.5–5.8	4.9–7.7	9.1–12.6	12.2–16.1	2.3–4.3	0.1–0.9	13.3–13.6
12th	%	48.5	3.9	2.2	2.5	3.6	5.1	10.4	11.0	12.7	14.77
(n=939)	CI	45.3–51.7	2.9–5.4	1.5–3.4	1.7–3.8	2.6–5.0	3.9–6.7	8.6–12.5	9.2–13.2	10.7–15.0	14.6–15.0

Note: The confidence interval (CI) is 95 percent.

Table BH6.12. Age of First Cigarette Use

		How old were you when you first smoked a cigarette, even just a puff?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	80.1	3.4	2.0	2.3	2.7	2.6	3.0	1.9	2.0	13.28
(n=3,458)	CI	78.7–81.5	2.9–4.1	1.6–2.5	1.9–2.9	2.2–3.3	2.1–3.2	2.5–3.7	1.5–2.4	1.5–2.5	13.1–13.5
Male	%	79.5	3.8	2.4	2.4	2.6	2.6	2.3	1.9	2.5	13.21
(n=1,688)	CI	77.4–81.4	3.0–4.9	1.8–3.3	1.7–3.2	1.9–3.4	2.0–3.5	1.7–3.2	1.3–2.8	1.8–3.5	12.9–13.5
Female	%	81.4	2.7	1.7	2.1	2.8	2.4	3.7	1.9	1.3	13.39
(n=1,661)	CI	79.4–83.2	2.0–3.6	1.2–2.4	1.5–2.9	2.2–3.8	1.8–3.3	2.8–4.7	1.3–2.8	0.9–2.1	13.1–13.6
White	%	77.9	3.5	2.1	2.4	3.0	3.0	3.3	2.3	2.4	13.41
(n=2,360)	CI	76.1–79.6	2.8–4.3	1.6–2.8	1.9–3.2	2.4–3.7	2.4–3.8	2.6–4.1	1.8–3.1	1.8–3.2	13.2–13.6
Black	%	91.8	2.0	0.3	1.3	1.6	0.3	1.5	0.6	0.6	12.94
(n=581)	CI	89.2–93.7	1.1–3.5	0.1–1.2	0.7–2.7	0.8–2.9	0.1–1.3	0.8–3.0	0.2–1.8	0.2–1.9	12.3–13.6
Hispanic	%	77.4	2.8	4.6	2.8	2.8	4.6	1.5	0.0	3.6	13.06
(n=67)	CI	65.4–86.1	0.7–10.9	1.4–13.7	0.7–10.9	0.7–10.9	1.4–13.7	0.2–10.2	–	0.9–13.5	11.7–14.4
8th	%	86.7	3.4	2.8	2.9	3.2	0.9	0.0	0.0	0.0	11.65
(n=1,311)	CI	84.8–88.5	2.6–4.6	2.1–3.9	2.1–4.0	2.4–4.3	0.5–1.6	–	–	–	11.5–11.8
10th	%	79.8	4.1	1.7	2.2	3.1	4.1	4.5	0.4	0.2	12.86
(n=1,205)	CI	77.4–82.0	3.1–5.3	1.1–2.7	1.5–3.1	2.2–4.2	3.1–5.4	3.4–5.8	0.2–1.0	0.0–0.7	12.6–13.1
12th	%	73.7	2.7	1.4	1.9	1.8	2.6	4.5	5.4	5.9	14.47
(n=942)	CI	70.8–76.4	1.9–4.0	0.8–2.4	1.2–3.0	1.1–2.9	1.8–3.9	3.4–6.1	4.1–7.0	4.6–7.6	14.2–14.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.13. Age of First Electronic Cigarette Use

		How old were you when you first used an electronic cigarette?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	69.4	1.1	1.1	2.0	4.8	4.4	8.1	4.5	4.6	14.48
(n=3,437)	CI	67.8–71.0	0.8–1.5	0.8–1.5	1.6–2.5	4.2–5.5	3.7–5.1	7.2–9.1	3.8–5.3	3.9–5.5	14.4–14.6
Male	%	68.4	1.5	1.2	1.9	5.2	4.6	7.7	3.9	5.6	14.43
(n=1,679)	CI	66.1–70.6	1.0–2.2	0.8–1.9	1.3–2.6	4.3–6.4	3.7–5.7	6.5–9.1	3.0–5.0	4.5–6.9	14.3–14.6
Female	%	71.0	0.5	1.0	1.9	4.3	3.8	8.5	5.3	3.7	14.59
(n=1,650)	CI	68.7–73.1	0.3–0.9	0.6–1.6	1.4–2.7	3.5–5.4	3.0–4.8	7.3–10.0	4.2–6.6	2.8–4.8	14.4–14.7
White	%	64.9	1.2	1.3	2.1	5.3	5.0	9.2	5.4	5.6	14.52
(n=2,343)	CI	62.9–66.8	0.9–1.8	0.9–1.9	1.6–2.7	4.5–6.3	4.2–6.0	8.1–10.5	4.5–6.5	4.7–6.7	14.4–14.7
Black	%	88.6	0.5	0.0	1.1	1.7	0.5	4.1	2.6	0.8	14.51
(n=580)	CI	85.8–91.0	0.2–1.5	–	0.5–2.3	1.0–3.1	0.2–1.7	2.7–6.0	1.6–4.4	0.3–2.1	14.1–14.9
Hispanic	%	76.6	0.0	0.0	0.0	5.5	6.1	1.5	5.0	5.3	14.95
(n=67)	CI	64.4–85.6	–	–	–	2.0–14.0	2.2–15.5	0.2–10.2	1.6–14.9	1.7–15.6	14.1–15.8
8th	%	81.6	1.6	2.1	3.0	9.4	2.3	0.0	0.0	0.0	12.48
(n=1,304)	CI	79.4–83.6	1.1–2.5	1.4–3.0	2.2–4.1	8.0–11.1	1.6–3.3	–	–	–	12.3–12.6
10th	%	65.2	0.9	1.0	1.7	2.9	7.0	18.0	3.1	0.1	14.33
(n=1,197)	CI	62.5–67.9	0.5–1.6	0.6–1.8	1.1–2.7	2.1–4.0	5.7–8.6	15.9–20.3	2.2–4.2	0.0–0.6	14.2–14.5
12th	%	61.3	0.6	0.3	1.2	2.0	3.7	6.0	10.6	14.3	15.60
(n=936)	CI	58.1–64.3	0.3–1.4	0.1–1.0	0.6–2.1	1.3–3.1	2.7–5.1	4.6–7.7	8.8–12.8	12.2–16.6	15.4–15.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.14. Age of First Marijuana Use

		How old were you when you first <u>smoked marijuana</u> ?									
		Never	Age 10 or younger	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17 or older	Average age of first use
Total	%	75.5	1.2	1.1	2.0	4.1	4.5	5.8	3.6	2.3	14.15
(n=3,465)	CI	74.0–76.9	0.9–1.7	0.8–1.5	1.6–2.5	3.5–4.8	3.8–5.2	5.1–6.7	3.0–4.3	1.8–2.9	14.0–14.3
Male	%	75.6	1.6	1.4	2.3	4.4	3.4	5.6	3.3	2.4	13.98
(n=1,694)	CI	73.4–77.6	1.1–2.4	1.0–2.1	1.7–3.1	3.5–5.5	2.6–4.4	4.5–6.8	2.5–4.4	1.7–3.3	13.8–14.2
Female	%	76.0	0.7	0.5	1.7	3.5	5.5	6.1	3.9	2.3	14.40
(n=1,663)	CI	73.8–78.0	0.4–1.2	0.3–0.9	1.1–2.4	2.7–4.5	4.4–6.7	5.0–7.4	3.0–5.1	1.6–3.2	14.2–14.6
White	%	76.5	0.9	0.8	1.7	4.5	4.3	5.6	3.7	2.0	14.23
(n=2,368)	CI	74.8–78.3	0.6–1.3	0.5–1.3	1.2–2.3	3.7–5.4	3.5–5.2	4.7–6.7	2.9–4.6	1.5–2.8	14.1–14.4
Black	%	76.6	1.7	1.0	2.0	3.0	3.4	6.6	2.9	2.8	14.18
(n=581)	CI	73.0–80.0	0.9–3.1	0.5–2.3	1.1–3.5	1.9–4.8	2.2–5.3	4.8–9.0	1.7–4.7	1.7–4.7	13.8–14.5
Hispanic	%	76.7	1.5	1.3	0.0	1.3	5.8	4.4	3.6	5.3	14.70
(n=67)	CI	64.6–85.7	0.2–10.2	0.2–9.3	–	0.2–9.3	2.1–14.8	1.4–13.2	0.9–13.5	1.7–15.6	13.6–15.8
8th	%	89.4	1.2	1.4	1.9	4.6	1.6	0.0	0.0	0.0	12.37
(n=1,318)	CI	87.6–90.9	0.7–2.0	0.9–2.2	1.3–2.8	3.5–5.8	1.0–2.4	–	–	–	12.2–12.6
10th	%	72.7	1.7	1.2	2.4	4.0	7.0	9.3	1.8	0.0	13.78
(n=1,204)	CI	70.1–75.1	1.1–2.6	0.7–2.0	1.7–3.4	3.0–5.2	5.7–8.6	7.8–11.1	1.2–2.8	–	13.6–14.0
12th	%	64.1	0.8	0.6	1.6	3.7	4.8	8.2	9.2	7.0	14.99
(n=943)	CI	61.0–67.1	0.4–1.7	0.3–1.4	1.0–2.6	2.7–5.1	3.6–6.3	6.6–10.2	7.5–11.2	5.5–8.8	14.8–15.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.15. Past Month Prescription Drug Misuse

		During the past 30 days, on how many days did you <u>use prescription drugs that were not prescribed to you?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	93.9	3.1	1.3	0.5	0.5	0.3	0.4	6.1
(n=3,476)	CI	93.1–94.7	2.6–3.8	1.0–1.7	0.3–0.8	0.3–0.8	0.2–0.6	0.2–0.7	5.3–6.9
Male	%	96.2	1.7	0.8	0.3	0.5	0.2	0.3	3.8
(n=1,699)	CI	95.2–97.1	1.2–2.4	0.5–1.4	0.1–0.7	0.2–1.0	0.1–0.5	0.1–0.7	2.9–4.8
Female	%	92.2	4.2	1.7	0.5	0.5	0.4	0.4	7.8
(n=1,664)	CI	90.8–93.5	3.3–5.3	1.2–2.5	0.2–1.0	0.2–0.9	0.2–0.9	0.2–0.9	6.5–9.2
White	%	94.1	3.5	1.2	0.4	0.4	0.2	0.3	5.9
(n=2,371)	CI	93.0–95.0	2.8–4.3	0.8–1.8	0.2–0.7	0.2–0.7	0.1–0.4	0.1–0.6	5.0–7.0
Black	%	94.7	1.6	1.6	0.5	0.7	0.7	0.1	5.3
(n=585)	CI	92.5–96.3	0.8–3.1	0.8–3.0	0.2–1.6	0.3–1.9	0.3–1.9	0.0–1.1	3.7–7.5
Hispanic	%	98.5	0.0	1.5	0.0	0.0	0.0	0.0	1.5
(n=67)	CI	89.8–99.8	–	0.2–10.2	–	–	–	–	0.2–10.2
8th	%	95.9	2.0	0.8	0.1	0.5	0.2	0.5	4.1
(n=1,319)	CI	94.7–96.9	1.4–3.0	0.4–1.4	0.0–0.5	0.3–1.1	0.1–0.7	0.2–1.0	3.1–5.3
10th	%	92.6	3.6	1.7	0.9	0.4	0.5	0.4	7.4
(n=1,207)	CI	90.9–93.9	2.6–4.8	1.1–2.5	0.5–1.6	0.2–1.0	0.2–1.1	0.2–1.0	6.1–9.1
12th	%	93.3	3.8	1.5	0.4	0.5	0.2	0.3	6.7
(n=950)	CI	91.5–94.7	2.7–5.2	0.9–2.5	0.2–1.1	0.2–1.3	0.1–0.8	0.1–1.0	5.3–8.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.16. Past Month Alcohol Use

		During the past 30 days, on how many days did you <u>drink one or more drinks of an alcoholic beverage?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	79.6	12.1	4.7	1.2	1.5	0.5	0.4	20.4
(n=3,472)	CI	78.1–80.9	11.0–13.2	4.0–5.4	0.9–1.6	1.2–2.0	0.3–0.8	0.3–0.7	19.1–21.9
Male	%	82.1	9.8	4.5	1.3	1.5	0.5	0.3	17.9
(n=1,700)	CI	80.1–83.9	8.5–11.4	3.6–5.7	0.8–2.0	1.0–2.2	0.3–1.0	0.1–0.7	16.1–19.9
Female	%	77.6	14.2	4.8	0.9	1.6	0.3	0.5	22.4
(n=1,660)	CI	75.5–79.6	12.6–16.0	3.9–6.0	0.6–1.6	1.1–2.4	0.1–0.7	0.2–1.0	20.4–24.5
White	%	78.6	12.6	5.0	1.3	1.8	0.5	0.3	21.4
(n=2,368)	CI	76.8–80.2	11.3–14.0	4.2–6.0	0.9–1.8	1.3–2.4	0.3–0.9	0.1–0.6	19.8–23.2
Black	%	85.2	9.4	3.0	0.9	0.5	0.4	0.5	14.8
(n=583)	CI	82.0–88.0	7.3–12.1	1.8–4.9	0.4–2.2	0.2–1.7	0.1–1.6	0.2–1.6	12.0–18.0
Hispanic	%	84.6	9.3	2.8	1.8	1.5	0.0	0.0	15.4
(n=67)	CI	73.3–91.7	4.2–19.7	0.7–10.9	0.2–12.1	0.2–10.2	–	–	8.3–26.7
8th	%	89.0	7.4	1.9	0.4	0.5	0.4	0.4	11.0
(n=1,319)	CI	87.2–90.6	6.1–9.0	1.3–2.8	0.2–0.9	0.3–1.1	0.2–0.9	0.2–0.9	9.4–12.8
10th	%	78.3	14.0	4.7	1.2	1.2	0.2	0.4	21.7
(n=1,203)	CI	75.9–80.5	12.2–16.1	3.7–6.1	0.7–2.0	0.7–2.0	0.1–0.8	0.2–1.0	19.5–24.1
12th	%	71.2	14.9	7.4	2.1	2.9	0.9	0.5	28.8
(n=950)	CI	68.2–74.0	12.8–17.3	5.9–9.3	1.4–3.2	2.0–4.2	0.5–1.8	0.2–1.3	26.0–31.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.17. Past Month Binge Drinking

		During the past 30 days, on how many days did you <u>have five or more drinks on the same occasion?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	90.2	4.9	2.3	0.9	1.1	0.3	0.3	9.8
(n=3,469)	CI	89.1–91.2	4.3–5.8	1.8–2.8	0.6–1.3	0.8–1.5	0.2–0.6	0.2–0.6	8.8–10.9
Male	%	90.2	4.5	2.2	1.2	1.2	0.5	0.2	9.8
(n=1,698)	CI	88.6–91.6	3.5–5.6	1.6–3.1	0.7–1.8	0.8–1.9	0.3–1.0	0.1–0.6	8.4–11.4
Female	%	90.8	5.5	2.2	0.5	0.8	0.0	0.2	9.2
(n=1,658)	CI	89.3–92.2	4.4–6.7	1.5–3.0	0.3–1.0	0.5–1.4	–	0.1–0.7	7.8–10.7
White	%	89.3	5.7	2.4	1.1	1.1	0.3	0.2	10.7
(n=2,368)	CI	87.9–90.5	4.8–6.8	1.8–3.1	0.7–1.6	0.7–1.7	0.1–0.6	0.1–0.5	9.5–12.1
Black	%	94.4	1.5	1.5	0.5	1.1	0.4	0.5	5.6
(n=584)	CI	92.2–96.1	0.8–3.0	0.7–3.0	0.2–1.7	0.5–2.4	0.1–1.6	0.2–1.5	3.9–7.8
Hispanic	%	93.9	2.8	1.5	1.8	0.0	0.0	0.0	6.1
(n=67)	CI	84.5–97.8	0.7–10.9	0.2–10.2	0.2–12.1	–	–	–	2.2–15.5
8th	%	96.7	2.0	0.5	0.2	0.2	0.2	0.2	3.3
(n=1,319)	CI	95.6–97.6	1.3–2.9	0.2–1.0	0.1–0.7	0.1–0.7	0.1–0.7	0.0–0.6	2.4–4.4
10th	%	90.4	5.0	2.5	0.7	1.0	0.1	0.3	9.6
(n=1,203)	CI	88.6–91.9	3.9–6.4	1.7–3.5	0.4–1.4	0.6–1.7	0.0–0.6	0.1–0.9	8.1–11.4
12th	%	83.3	8.0	3.9	1.7	2.0	0.7	0.4	16.7
(n=947)	CI	80.8–85.5	6.4–9.9	2.8–5.3	1.0–2.7	1.3–3.1	0.4–1.5	0.2–1.1	14.5–19.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.18. Past Month Cigarette Use

		During the past 30 days, on how many days did you <u>smoke all or part of a cigarette?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	89.9	4.1	1.8	1.0	1.1	0.6	1.6	10.1
(n=3,472)	CI	88.9–90.9	3.5–4.8	1.4–2.3	0.7–1.4	0.8–1.5	0.4–0.9	1.2–2.1	9.1–11.1
Male	%	89.7	3.6	2.3	0.9	1.0	0.6	1.8	10.3
(n=1,698)	CI	88.2–91.1	2.8–4.7	1.7–3.2	0.5–1.5	0.6–1.6	0.3–1.2	1.2–2.5	8.9–11.8
Female	%	90.7	4.2	1.3	0.9	1.0	0.5	1.3	9.3
(n=1,662)	CI	89.1–92.0	3.4–5.4	0.9–2.0	0.6–1.6	0.6–1.7	0.2–1.0	0.8–2.0	8.0–10.9
White	%	88.3	4.7	2.2	0.9	1.4	0.6	2.0	11.7
(n=2,367)	CI	86.9–89.6	3.9–5.6	1.6–2.9	0.6–1.4	1.0–2.0	0.3–1.0	1.5–2.6	10.4–13.1
Black	%	96.0	1.9	0.3	0.7	0.2	0.3	0.5	4.0
(n=586)	CI	94.0–97.3	1.0–3.4	0.1–1.2	0.3–1.9	0.0–1.4	0.1–1.4	0.2–1.7	2.7–6.0
Hispanic	%	89.3	4.5	0.0	6.2	0.0	0.0	0.0	10.7
(n=66)	CI	78.8–95.0	1.4–13.5	–	2.3–15.7	–	–	–	5.0–21.2
8th	%	92.9	3.5	1.3	0.6	0.6	0.5	0.6	7.1
(n=1,318)	CI	91.4–94.2	2.6–4.6	0.8–2.1	0.3–1.2	0.3–1.2	0.2–1.0	0.3–1.2	5.8–8.6
10th	%	91.2	3.6	1.7	0.9	0.8	0.3	1.5	8.8
(n=1,205)	CI	89.5–92.7	2.6–4.8	1.1–2.6	0.5–1.6	0.4–1.5	0.1–0.9	0.9–2.4	7.3–10.5
12th	%	85.5	5.2	2.4	1.4	1.8	0.9	2.7	14.5
(n=949)	CI	83.1–87.6	4.0–6.9	1.6–3.6	0.8–2.3	1.1–2.9	0.5–1.8	1.9–4.0	12.4–16.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.19. Past Month Electronic Cigarette Use

		During the past 30 days, on how many days did you <u>use an e-cigarette, e-cigar, or e-hookah?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	77.9	6.4	4.5	2.5	3.1	1.6	4.0	22.1
(n=3,465)	CI	76.4–79.3	5.6–7.3	3.9–5.3	2.0–3.0	2.6–3.8	1.2–2.1	3.4–4.7	20.7–23.6
Male	%	76.6	7.2	4.6	2.1	3.2	1.6	4.7	23.4
(n=1,693)	CI	74.5–78.6	6.0–8.5	3.7–5.7	1.5–2.9	2.4–4.2	1.1–2.4	3.8–5.9	21.4–25.5
Female	%	79.6	5.6	4.5	2.6	3.1	1.4	3.2	20.4
(n=1,661)	CI	77.6–81.5	4.6–6.8	3.6–5.6	1.9–3.6	2.4–4.1	0.9–2.1	2.4–4.2	18.5–22.4
White	%	74.6	7.0	5.1	2.9	3.5	1.9	4.8	25.4
(n=2,359)	CI	72.8–76.4	6.0–8.1	4.3–6.1	2.3–3.7	2.9–4.4	1.4–2.6	4.0–5.8	23.6–27.2
Black	%	91.4	4.1	1.9	0.7	0.7	0.2	1.0	8.6
(n=586)	CI	88.8–93.4	2.7–6.0	1.0–3.4	0.3–1.9	0.3–1.9	0.0–1.2	0.5–2.3	6.6–11.2
Hispanic	%	81.6	6.1	1.8	0.0	5.6	1.3	3.6	18.4
(n=67)	CI	70.0–89.5	2.2–15.5	0.2–12.1	–	2.1–14.4	0.2–9.3	0.9–13.5	10.5–30.0
8th	%	87.2	4.9	2.8	1.1	2.0	1.0	1.1	12.8
(n=1,316)	CI	85.2–88.9	3.9–6.3	2.0–3.9	0.6–1.8	1.3–2.9	0.6–1.7	0.6–1.8	11.1–14.8
10th	%	75.3	7.5	6.5	2.7	2.9	1.6	3.6	24.7
(n=1,204)	CI	72.8–77.7	6.1–9.1	5.2–8.0	1.9–3.8	2.1–4.0	1.0–2.5	2.7–4.8	22.3–27.2
12th	%	71.0	6.8	4.3	3.6	4.5	2.2	7.5	29.0
(n=945)	CI	68.0–73.8	5.4–8.6	3.2–5.8	2.6–5.0	3.4–6.1	1.4–3.4	6.0–9.3	26.2–32.0

Note: The confidence interval (CI) is 95 percent.

Table BH6.20. Past Month Smokeless Tobacco Use

		During the past 30 days, on how many days did you <u>use smokeless tobacco?</u>							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	93.1	2.1	1.1	0.8	0.8	0.5	1.6	6.9
(n=3,472)	CI	92.2–93.9	1.7–2.6	0.8–1.5	0.5–1.1	0.5–1.1	0.3–0.8	1.2–2.1	6.1–7.8
Male	%	89.4	2.6	1.9	1.0	1.2	1.0	2.9	10.6
(n=1,695)	CI	87.9–90.9	1.9–3.5	1.4–2.7	0.6–1.6	0.8–1.9	0.6–1.6	2.2–3.9	9.1–12.1
Female	%	97.4	1.5	0.3	0.4	0.2	0.1	0.2	2.6
(n=1,665)	CI	96.5–98.1	1.0–2.2	0.1–0.8	0.2–0.8	0.1–0.6	0.0–0.4	0.1–0.6	1.9–3.5
White	%	91.8	2.4	1.3	0.9	0.9	0.7	2.1	8.2
(n=2,367)	CI	90.6–92.9	1.8–3.1	0.9–1.8	0.6–1.4	0.6–1.3	0.4–1.1	1.6–2.8	7.1–9.4
Black	%	98.0	1.0	0.0	0.6	0.4	0.0	0.0	2.0
(n=586)	CI	96.4–98.9	0.5–2.3	–	0.2–1.7	0.1–1.6	–	–	1.1–3.6
Hispanic	%	98.2	0.0	0.0	0.0	0.0	0.0	1.8	1.8
(n=67)	CI	87.9–99.8	–	–	–	–	–	0.2–12.1	0.2–12.1
8th	%	95.7	1.6	1.0	0.5	0.5	0.2	0.6	4.3
(n=1,318)	CI	94.4–96.7	1.0–2.4	0.6–1.7	0.2–1.0	0.3–1.1	0.0–0.6	0.3–1.2	3.3–5.6
10th	%	92.8	2.8	1.4	0.6	0.8	0.5	1.1	7.2
(n=1,206)	CI	91.2–94.1	2.0–3.9	0.9–2.3	0.3–1.2	0.4–1.5	0.2–1.1	0.6–1.8	5.9–8.8
12th	%	90.9	1.9	0.9	1.3	0.9	0.8	3.3	9.1
(n=948)	CI	88.9–92.5	1.2–3.0	0.5–1.8	0.7–2.2	0.5–1.8	0.4–1.7	2.3–4.6	7.5–11.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.21. Past Month Marijuana Use

		During the past 30 days, on how many days did you <u>use marijuana or hashish</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	86.3	4.3	2.7	1.4	1.9	1.2	2.3	13.7
(n=3,465)	CI	85.1–87.4	3.6–5.0	2.2–3.3	1.0–1.9	1.5–2.4	0.9–1.7	1.8–2.8	12.6–14.9
Male	%	86.6	3.3	2.1	1.6	1.9	1.4	3.0	13.4
(n=1,694)	CI	84.8–88.2	2.6–4.3	1.5–2.9	1.1–2.3	1.4–2.7	1.0–2.2	2.3–4.0	11.8–15.2
Female	%	86.4	5.2	3.2	1.1	1.6	1.0	1.4	13.6
(n=1,660)	CI	84.7–88.0	4.2–6.4	2.5–4.2	0.7–1.8	1.1–2.3	0.6–1.6	0.9–2.2	12.0–15.3
White	%	87.7	4.0	2.4	1.2	1.5	1.1	2.1	12.3
(n=2,366)	CI	86.2–89.0	3.3–4.9	1.8–3.1	0.8–1.8	1.1–2.2	0.7–1.6	1.6–2.8	11.0–13.8
Black	%	85.7	3.8	2.4	1.2	1.8	2.0	3.0	14.3
(n=580)	CI	82.5–88.4	2.5–5.8	1.4–4.1	0.6–2.6	0.9–3.3	1.1–3.6	1.8–4.8	11.6–17.5
Hispanic	%	91.0	4.7	2.8	1.5	0.0	0.0	0.0	9.0
(n=66)	CI	80.9–96.0	1.5–13.9	0.7–11.1	0.2–10.4	–	–	–	4.0–19.1
8th	%	93.4	2.6	1.4	0.5	0.9	0.5	0.8	6.6
(n=1,317)	CI	91.9–94.6	1.8–3.6	0.9–2.2	0.2–1.0	0.5–1.6	0.2–1.0	0.5–1.5	5.4–8.1
10th	%	83.6	5.2	3.7	1.7	2.2	1.7	2.0	16.4
(n=1,202)	CI	81.4–85.6	4.1–6.6	2.7–4.9	1.1–2.7	1.5–3.2	1.1–2.6	1.3–3.0	14.4–18.6
12th	%	81.9	5.1	2.9	2.0	2.5	1.6	4.0	18.1
(n=946)	CI	79.3–84.2	3.8–6.6	2.0–4.2	1.3–3.1	1.7–3.7	1.0–2.6	2.9–5.5	15.8–20.7

Note: The confidence interval (CI) is 95 percent.

Table BH6.22. Past Month Cocaine Use

		During the past 30 days, on how many days did you <u>use cocaine (crack, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.0	0.4	0.1	0.2	0.1	0.1	0.1	1.0
(n=3,478)	CI	98.6–99.3	0.2–0.7	0.0–0.3	0.1–0.5	0.0–0.3	0.0–0.2	0.0–0.3	0.7–1.4
Male	%	99.1	0.3	0.1	0.2	0.1	0.1	0.2	0.9
(n=1,701)	CI	98.5–99.4	0.1–0.7	0.0–0.5	0.1–0.6	0.0–0.4	0.0–0.5	0.1–0.6	0.6–1.5
Female	%	99.2	0.6	0.1	0.0	0.1	0.0	0.1	0.8
(n=1,665)	CI	98.6–99.5	0.3–1.1	0.0–0.4	–	0.0–0.4	–	0.0–0.5	0.5–1.4
White	%	99.2	0.4	0.0	0.2	0.0	0.0	0.1	0.8
(n=2,371)	CI	98.7–99.5	0.2–0.8	0.0–0.3	0.1–0.5	0.0–0.3	–	0.0–0.4	0.5–1.3
Black	%	98.9	0.4	0.2	0.4	0.0	0.2	0.0	1.1
(n=586)	CI	97.5–99.5	0.1–1.5	0.0–1.4	0.1–1.5	–	0.0–1.4	–	0.5–2.5
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.4	0.2	0.1	0.2	0.1	0.1	0.1	0.6
(n=1,319)	CI	98.8–99.7	0.0–0.6	0.0–0.5	0.0–0.6	0.0–0.5	0.0–0.5	0.0–0.5	0.3–1.2
10th	%	98.9	0.6	0.1	0.2	0.2	0.0	0.1	1.1
(n=1,209)	CI	98.2–99.4	0.3–1.2	0.0–0.6	0.0–0.7	0.0–0.7	–	0.0–0.6	0.6–1.8
12th	%	98.6	0.5	0.1	0.4	0.0	0.1	0.2	1.4
(n=950)	CI	97.7–99.2	0.2–1.3	0.0–0.7	0.2–1.1	–	0.0–0.7	0.1–0.8	0.8–2.3

Note: The confidence interval (CI) is 95 percent.

Table BH6.23. Past Month Inhalants Use

		During the past 30 days, on how many days did you <u>use inhalants (glue, gas, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.1	0.3	0.1	0.1	0.1	0.1	0.2	0.9
(n=3,478)	CI	98.7–99.3	0.1–0.5	0.0–0.3	0.1–0.3	0.0–0.2	0.1–0.3	0.1–0.4	0.7–1.3
Male	%	99.2	0.2	0.2	0.2	0.0	0.1	0.2	0.8
(n=1,701)	CI	98.6–99.5	0.1–0.6	0.1–0.6	0.1–0.5	–	0.0–0.5	0.1–0.6	0.5–1.4
Female	%	99.2	0.3	0.1	0.1	0.1	0.2	0.1	0.8
(n=1,665)	CI	98.7–99.6	0.1–0.7	0.0–0.4	0.0–0.4	0.0–0.4	0.1–0.5	0.0–0.5	0.4–1.3
White	%	99.4	0.3	0.0	0.1	0.0	0.1	0.2	0.6
(n=2,371)	CI	99.0–99.6	0.1–0.6	0.0–0.3	0.0–0.3	–	0.0–0.3	0.1–0.4	0.4–1.0
Black	%	98.4	0.4	0.4	0.4	0.1	0.4	0.0	1.6
(n=586)	CI	96.9–99.2	0.1–1.5	0.1–1.5	0.1–1.5	0.0–1.1	0.1–1.5	–	0.8–3.1
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.2	0.1	0.0	0.2	0.2	0.2	0.2	0.8
(n=1,319)	CI	98.6–99.6	0.0–0.5	–	0.0–0.6	0.0–0.6	0.1–0.7	0.0–0.6	0.4–1.4
10th	%	99.0	0.4	0.2	0.2	0.0	0.1	0.2	1.0
(n=1,209)	CI	98.3–99.4	0.2–1.0	0.0–0.7	0.0–0.7	–	0.0–0.6	0.0–0.7	0.6–1.7
12th	%	99.0	0.3	0.2	0.1	0.0	0.1	0.3	1.0
(n=950)	CI	98.1–99.4	0.1–1.0	0.1–0.8	0.0–0.7	–	0.0–0.7	0.1–1.0	0.6–1.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.24. Past Month Hallucinogens Use

		During the past 30 days, on how many days did you <u>use hallucinogens (PCP, LSD, etc.)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	98.9	0.4	0.2	0.1	0.1	0.1	0.1	1.1
(n=3,478)	CI	98.5–99.2	0.2–0.7	0.1–0.5	0.1–0.3	0.0–0.3	0.0–0.3	0.0–0.3	0.8–1.5
Male	%	98.9	0.3	0.2	0.2	0.1	0.1	0.2	1.1
(n=1,701)	CI	98.2–99.3	0.1–0.8	0.1–0.6	0.1–0.5	0.0–0.5	0.0–0.5	0.1–0.6	0.7–1.8
Female	%	99.3	0.3	0.2	0.1	0.1	0.0	0.1	0.7
(n=1,665)	CI	98.7–99.6	0.1–0.8	0.1–0.6	0.0–0.4	0.0–0.4	–	0.0–0.5	0.4–1.3
White	%	99.1	0.4	0.2	0.0	0.1	0.0	0.1	0.9
(n=2,371)	CI	98.6–99.4	0.2–0.8	0.1–0.5	0.0–0.3	0.0–0.3	0.0–0.3	0.0–0.4	0.6–1.4
Black	%	99.0	0.2	0.2	0.4	0.0	0.2	0.0	1.0
(n=586)	CI	97.7–99.6	0.0–1.4	0.0–1.4	0.1–1.5	–	0.0–1.4	–	0.4–2.3
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.5	0.0	0.2	0.2	0.0	0.0	0.1	0.5
(n=1,319)	CI	99.0–99.8	–	0.0–0.6	0.1–0.7	–	–	0.0–0.5	0.2–1.0
10th	%	98.8	0.5	0.2	0.1	0.2	0.1	0.1	1.2
(n=1,209)	CI	98.1–99.3	0.2–1.1	0.0–0.7	0.0–0.6	0.1–0.8	0.0–0.6	0.0–0.6	0.7–1.9
12th	%	98.4	0.6	0.4	0.1	0.1	0.1	0.2	1.6
(n=950)	CI	97.4–99.1	0.3–1.4	0.2–1.1	0.0–0.7	0.0–0.7	0.0–0.7	0.1–0.8	0.9–2.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.25. Past Month Heroin Use

		During the past 30 days, on how many days did you <u>use heroin (opiates)</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.4	0.1	0.1	0.2	0.0	0.0	0.2	0.6
(n=3,477)	CI	99.1–99.6	0.0–0.3	0.0–0.2	0.1–0.4	0.0–0.2	–	0.1–0.4	0.4–0.9
Male	%	99.5	0.2	0.0	0.1	0.1	0.0	0.2	0.5
(n=1,700)	CI	99.0–99.8	0.1–0.6	–	0.0–0.4	0.0–0.4	–	0.1–0.6	0.2–1.0
Female	%	99.7	0.0	0.1	0.1	0.0	0.0	0.1	0.3
(n=1,665)	CI	99.2–99.9	–	0.0–0.5	0.0–0.5	–	–	0.0–0.5	0.1–0.8
White	%	99.6	0.1	0.1	0.1	0.0	0.0	0.1	0.4
(n=2,371)	CI	99.2–99.8	0.0–0.4	0.0–0.3	0.0–0.4	0.0–0.3	–	0.0–0.4	0.2–0.8
Black	%	99.4	0.2	0.0	0.4	0.0	0.0	0.0	0.6
(n=585)	CI	98.2–99.8	0.0–1.4	–	0.1–1.5	–	–	–	0.2–1.8
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.6	0.0	0.0	0.2	0.1	0.0	0.1	0.4
(n=1,319)	CI	99.1–99.8	–	–	0.1–0.7	0.0–0.5	–	0.0–0.5	0.2–0.9
10th	%	99.6	0.1	0.2	0.1	0.0	0.0	0.1	0.4
(n=1,209)	CI	99.0–99.8	0.0–0.6	0.0–0.7	0.0–0.6	–	–	0.0–0.6	0.2–1.0
12th	%	99.1	0.3	0.0	0.3	0.0	0.0	0.3	0.9
(n=949)	CI	98.2–99.5	0.1–1.0	–	0.1–1.0	–	–	0.1–1.0	0.5–1.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.26. Past Month Steroids Use

		During the past 30 days, on how many days did you <u>use steroids</u> ?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.6	0.0	0.1	0.1	0.0	0.0	0.1	0.4
(n=3,476)	CI	99.3–99.7	0.0–0.2	0.0–0.3	0.1–0.4	0.0–0.2	–	0.1–0.4	0.3–0.7
Male	%	99.6	0.1	0.0	0.1	0.1	0.0	0.2	0.4
(n=1,700)	CI	99.2–99.8	0.0–0.4	–	0.0–0.5	0.0–0.4	–	0.1–0.6	0.2–0.8
Female	%	99.7	0.0	0.1	0.1	0.0	0.0	0.1	0.3
(n=1,665)	CI	99.2–99.9	–	0.0–0.5	0.0–0.5	–	–	0.0–0.5	0.1–0.8
White	%	99.6	0.0	0.1	0.1	0.0	0.0	0.1	0.4
(n=2,371)	CI	99.3–99.8	0.0–0.3	0.0–0.3	0.0–0.4	0.0–0.3	–	0.0–0.4	0.2–0.7
Black	%	99.6	0.0	0.0	0.4	0.0	0.0	0.0	0.4
(n=585)	CI	98.5–99.9	–	–	0.1–1.5	–	–	–	0.1–1.5
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.6	0.1	0.0	0.1	0.1	0.0	0.2	0.4
(n=1,318)	CI	99.1–99.8	0.0–0.5	–	0.0–0.5	0.0–0.5	–	0.0–0.6	0.2–0.9
10th	%	99.6	0.0	0.2	0.2	0.0	0.0	0.1	0.4
(n=1,209)	CI	99.0–99.8	–	0.0–0.7	0.0–0.7	–	–	0.0–0.6	0.2–1.0
12th	%	99.5	0.0	0.1	0.2	0.0	0.0	0.2	0.5
(n=949)	CI	98.7–99.8	–	0.0–0.7	0.1–0.8	–	–	0.1–0.8	0.2–1.3

Note: The confidence interval (CI) is 95 percent.

Table BH6.27. Past Month Ecstasy Use

		During the past 30 days, on how many days did you use ecstasy (MDMA)?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.2	0.4	0.1	0.1	0.1	0.0	0.1	0.8
(n=3,475)	CI	98.9–99.5	0.3–0.7	0.0–0.3	0.0–0.3	0.0–0.2	0.0–0.2	0.0–0.3	0.5–1.1
Male	%	99.1	0.4	0.1	0.1	0.1	0.1	0.1	0.9
(n=1,700)	CI	98.5–99.5	0.2–0.9	0.0–0.5	0.0–0.5	0.0–0.4	0.0–0.4	0.0–0.5	0.5–1.5
Female	%	99.5	0.3	0.1	0.0	0.1	0.0	0.1	0.5
(n=1,665)	CI	99.0–99.8	0.1–0.7	0.0–0.4	–	0.0–0.4	–	0.0–0.5	0.2–1.0
White	%	99.3	0.5	0.0	0.0	0.1	0.0	0.1	0.7
(n=2,370)	CI	98.8–99.5	0.3–1.0	0.0–0.3	–	0.0–0.3	–	0.0–0.4	0.5–1.2
Black	%	99.1	0.2	0.2	0.5	0.0	0.0	0.0	0.9
(n=585)	CI	97.8–99.6	0.0–1.4	0.0–1.4	0.2–1.6	–	–	–	0.4–2.2
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.5	0.2	0.0	0.0	0.2	0.0	0.1	0.5
(n=1,317)	CI	99.0–99.8	0.1–0.7	–	–	0.0–0.6	–	0.0–0.5	0.2–1.0
10th	%	99.3	0.3	0.1	0.2	0.0	0.1	0.0	0.7
(n=1,209)	CI	98.7–99.7	0.1–0.9	0.0–0.6	0.0–0.7	–	0.0–0.6	–	0.3–1.3
12th	%	98.8	0.7	0.1	0.1	0.0	0.0	0.2	1.2
(n=949)	CI	97.9–99.4	0.4–1.5	0.0–0.7	0.0–0.7	–	–	0.1–0.8	0.6–2.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.28. Past Month Methamphetamines Use

		During the past 30 days, on how many days did you use methamphetamines (crank, etc.)?							
		0 days	1–2 days	3–5 days	6–9 days	10–19 days	20–29 days	All 30 days	Any Use
Total	%	99.3	0.1	0.2	0.1	0.1	0.0	0.2	0.7
(n=3,474)	CI	99.0–99.6	0.0–0.3	0.1–0.5	0.0–0.3	0.0–0.3	0.0–0.2	0.1–0.4	0.4–1.0
Male	%	99.4	0.1	0.2	0.1	0.1	0.0	0.1	0.6
(n=1,700)	CI	98.9–99.7	0.0–0.5	0.1–0.6	0.0–0.4	0.0–0.5	–	0.0–0.4	0.3–1.1
Female	%	99.5	0.1	0.2	0.0	0.0	0.1	0.2	0.5
(n=1,664)	CI	99.1–99.8	0.0–0.4	0.1–0.5	–	–	0.0–0.4	0.1–0.6	0.2–0.9
White	%	99.5	0.1	0.2	0.0	0.0	0.0	0.0	0.5
(n=2,370)	CI	99.1–99.7	0.0–0.4	0.1–0.6	–	0.0–0.3	0.0–0.3	0.0–0.3	0.3–0.9
Black	%	99.4	0.0	0.0	0.4	0.2	0.0	0.0	0.6
(n=584)	CI	98.2–99.8	–	–	0.1–1.5	0.0–1.4	–	–	0.2–1.8
Hispanic	%	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
(n=67)	CI	–	–	–	–	–	–	–	–
8th	%	99.8	0.1	0.2	0.0	0.0	0.0	0.0	0.2
(n=1,316)	CI	99.3–99.9	0.0–0.5	0.0–0.6	–	–	–	–	0.1–0.7
10th	%	99.3	0.1	0.2	0.1	0.1	0.1	0.2	0.7
(n=1,208)	CI	98.6–99.6	0.0–0.6	0.0–0.7	0.0–0.6	0.0–0.6	0.0–0.6	0.1–0.8	0.4–1.4
12th	%	99.0	0.2	0.4	0.1	0.1	0.0	0.2	1.0
(n=950)	CI	98.1–99.4	0.1–0.8	0.2–1.1	0.0–0.7	0.0–0.7	–	0.1–0.8	0.6–1.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.29. Availability of Prescription Drugs

		How easy is it to get <u>prescription drugs not prescribed to you</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	26.2	25.4	16.3	32.2	48.4
(n=1,412)	CI	23.9–28.6	23.2–27.8	14.4–18.3	29.8–34.6	45.8–51.0
Male	%	24.7	24.1	15.7	35.4	51.2
(n=620)	CI	21.4–28.3	20.9–27.7	13.0–18.8	31.8–39.3	47.2–55.1
Female	%	25.7	26.9	16.6	30.7	47.4
(n=727)	CI	22.7–29.1	23.7–30.2	14.1–19.6	27.5–34.2	43.7–51.1
White	%	22.7	26.6	18.2	32.4	50.6
(n=1,000)	CI	20.2–25.5	24.0–29.5	15.9–20.7	29.6–35.4	47.5–53.7
Black	%	30.0	16.5	10.6	42.9	53.5
(n=179)	CI	23.6–37.2	11.7–22.8	6.8–16.1	35.7–50.3	46.0–60.8
Hispanic	%	32.4	19.0	22.3	26.2	48.6
(n=18)	CI	13.9–58.7	5.6–48.3	7.7–49.8	10.2–52.8	25.2–72.5
8th	%	16.2	22.7	13.6	47.5	61.1
(n=542)	CI	13.3–19.6	19.3–26.4	11.0–16.8	43.3–51.7	57.0–65.2
10th	%	31.2	26.4	17.7	24.7	42.4
(n=472)	CI	27.2–35.6	22.6–30.5	14.5–21.4	21.0–28.8	38.0–46.9
12th	%	31.3	27.3	17.5	24.0	41.5
(n=398)	CI	26.9–36.0	23.1–31.8	14.1–21.6	20.1–28.5	36.7–46.4

Note: The confidence interval (CI) is 95 percent.

Table BH6.30. Availability of Alcohol

		How easy is it to get <u>alcohol (beer, coolers, liquor, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	27.5	35.4	19.0	18.1	37.1
(n=1,990)	CI	25.6–29.6	33.3–37.5	17.3–20.7	16.5–19.9	35.0–39.2
Male	%	26.4	33.8	20.4	19.4	39.8
(n=928)	CI	23.6–29.4	30.8–36.9	17.9–23.1	17.0–22.1	36.7–43.0
Female	%	27.0	37.9	17.4	17.7	35.1
(n=989)	CI	24.3–29.9	34.9–41.0	15.2–19.9	15.5–20.2	32.2–38.1
White	%	24.7	38.4	19.2	17.7	36.9
(n=1,431)	CI	22.5–27.0	35.9–41.0	17.2–21.3	15.8–19.7	34.4–39.4
Black	%	32.5	20.4	18.8	28.3	47.1
(n=247)	CI	26.8–38.7	15.7–26.0	14.4–24.2	23.0–34.3	40.9–53.5
Hispanic	%	30.1	45.8	14.0	10.1	24.1
(n=35)	CI	16.5–48.3	29.5–63.1	5.7–30.8	3.6–25.1	12.5–41.5
8th	%	19.0	28.6	23.3	29.1	52.5
(n=668)	CI	16.2–22.1	25.2–32.1	20.3–26.7	25.8–32.7	48.7–56.2
10th	%	26.9	38.2	20.7	14.2	34.9
(n=696)	CI	23.7–30.3	34.7–41.9	17.9–23.9	11.8–17.0	31.5–38.5
12th	%	34.9	38.2	13.9	13.1	26.9
(n=626)	CI	31.2–38.7	34.5–42.1	11.4–16.8	10.6–15.9	23.6–30.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.31. Availability of Tobacco

		How easy is it to get <u>tobacco (cigarettes, cigars, dip, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	39.0	30.5	11.2	19.3	30.5
(n=1,915)	CI	36.8–41.2	28.5–32.6	9.9–12.7	17.6–21.1	28.5–32.6
Male	%	39.1	30.3	11.2	19.4	30.6
(n=921)	CI	35.9–42.3	27.4–33.3	9.4–13.4	17.0–22.1	27.8–33.7
Female	%	37.4	31.3	11.0	20.2	31.3
(n=921)	CI	34.3–40.7	28.3–34.4	9.2–13.2	17.8–22.9	28.4–34.4
White	%	39.7	31.9	11.8	16.6	28.4
(n=1,423)	CI	37.2–42.4	29.5–34.3	10.3–13.6	14.8–18.5	26.1–30.8
Black	%	30.8	21.9	7.4	39.9	47.3
(n=209)	CI	24.8–37.5	16.7–28.2	4.6–11.9	33.4–46.7	40.5–54.2
Hispanic	%	47.8	32.1	6.8	13.3	20.2
(n=27)	CI	28.8–67.4	16.6–52.8	1.5–25.6	4.7–32.4	8.7–40.0
8th	%	19.4	31.3	13.9	35.4	49.3
(n=638)	CI	16.5–22.7	27.8–35.0	11.4–16.8	31.7–39.2	45.4–53.2
10th	%	38.3	33.8	14.1	13.7	27.8
(n=662)	CI	34.7–42.1	30.3–37.5	11.7–17.0	11.3–16.5	24.5–31.4
12th	%	54.6	27.0	6.5	12.0	18.4
(n=615)	CI	50.7–58.5	23.6–30.6	4.8–8.7	9.6–14.8	15.5–21.7

Note: The confidence interval (CI) is 95 percent.

Table BH6.32. Availability of Marijuana

		How easy is it to get <u>marijuana (pot, hash, etc.)</u> ?				
		Very easy	Fairly easy	Fairly difficult	Very difficult	Combined: (Very difficult, Fairly difficult)
Total	%	31.8	25.2	13.1	29.8	43.0
(n=1,640)	CI	29.6–34.2	23.1–27.4	11.6–14.9	27.7–32.0	40.6–45.4
Male	%	32.0	25.6	13.8	28.6	42.4
(n=784)	CI	28.8–35.5	22.6–28.8	11.5–16.4	25.6–31.9	39.0–45.9
Female	%	29.8	25.7	12.7	31.8	44.5
(n=793)	CI	26.6–33.2	22.7–28.9	10.5–15.2	28.7–35.1	41.0–48.0
White	%	27.8	26.4	13.6	32.1	45.8
(n=1,120)	CI	25.2–30.5	23.9–29.2	11.7–15.8	29.5–34.9	42.9–48.7
Black	%	41.5	22.0	6.7	29.7	36.4
(n=244)	CI	35.4–48.0	17.2–27.8	4.2–10.6	24.3–35.7	30.6–42.7
Hispanic	%	30.6	33.7	28.5	7.2	35.7
(n=24)	CI	14.5–53.3	16.9–56.1	13.3–50.9	1.6–27.0	18.4–57.6
8th	%	9.6	15.4	13.5	61.6	75.1
(n=533)	CI	7.3–12.4	12.5–18.7	10.8–16.7	57.4–65.7	71.2–78.6
10th	%	36.6	28.3	15.5	19.6	35.1
(n=574)	CI	32.8–40.7	24.8–32.1	12.7–18.6	16.6–23.1	31.3–39.1
12th	%	44.1	29.7	10.8	15.3	26.2
(n=533)	CI	39.9–48.4	26.0–33.7	8.5–13.8	12.5–18.6	22.6–30.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.33. Availability of Prescription Drugs

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)			
		I did get Rx drugs	I did <u>not</u> get Rx drugs
Total	%	3.3	96.7
(n=3,469)	CI	2.7–4.0	96.0–97.3
Male	%	2.2	97.8
(n=1,697)	CI	1.6–3.0	97.0–98.4
Female	%	3.8	96.2
(n=1,658)	CI	3.0–4.9	95.1–97.0
White	%	3.1	96.9
(n=2,367)	CI	2.5–3.9	96.1–97.5
Black	%	2.8	97.2
(n=581)	CI	1.7–4.5	95.5–98.3
Hispanic	%	0.0	100.0
(n=67)	CI	–	–
8th	%	1.4	98.6
(n=1,316)	CI	0.9–2.3	97.7–99.1
10th	%	4.1	95.9
(n=1,205)	CI	3.1–5.4	94.6–96.9
12th	%	4.3	95.7
(n=948)	CI	3.2–5.8	94.2–96.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.34. Availability of Alcohol

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)			
		I did get alcohol	I did <u>not</u> get alcohol
Total	%	18.2	81.8
(n=3,456)	CI	16.9–19.5	80.5–83.1
Male	%	16.1	83.9
(n=1,694)	CI	14.4–18.0	82.0–85.6
Female	%	19.8	80.2
(n=1,651)	CI	17.9–21.8	78.2–82.1
White	%	6.3	93.7
(n=2,358)	CI	5.7–7.0	93.0–94.3
Black	%	15.6	84.4
(n=582)	CI	14.2–17.1	82.9–85.8
Hispanic	%	21.1	78.9
(n=66)	CI	19.6–22.8	77.2–80.4
8th	%	8.8	91.2
(n=1,314)	CI	7.4–10.5	89.5–92.6
10th	%	19.3	80.7
(n=1,202)	CI	17.2–21.7	78.3–82.8
12th	%	26.6	73.4
(n=940)	CI	23.9–29.6	70.4–76.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.35. Availability of Prescription Drugs – Sources

During the past 30 days, if you used prescription drugs to get high, how did you get the drugs? (Mark all that apply)										
		Total (n=108)	Male (n=36)	Female (n=61)	White (n=71)	Black (n=16)	Hispanic (n=0)	8th (n=19)	10th (n=50)	12th (n=39)
Got a prescription from one doctor	%	14.6	13.8	16.5	15.8	17.9	0.0	21.1	10.0	17.1
	CI	9.0–22.7	5.6–30.2	8.9–28.5	8.8–26.7	5.0–47.5	–	7.3–47.3	4.1–22.4	8.1–32.5
Got a prescription from more than one doctor	%	2.4	2.6	2.8	1.2	6.0	0.0	10.5	2.0	0.0
	CI	0.8–7.3	0.3–18.0	0.7–10.9	0.2–8.3	0.7–38.3	–	2.3–37.1	0.3–13.7	–
Bought on the Internet	%	6.3	5.3	1.6	2.9	0.0	0.0	5.3	8.0	4.9
	CI	3.0–12.8	1.2–20.0	0.2–10.8	0.7–11.3	–	–	0.6–33.8	2.9–20.0	1.1–18.4
Bought from a friend, relative, or someone you know	%	37.7	43.4	38.4	36.8	36.5	0.0	57.9	44.0	24.4
	CI	29.1–47.2	27.7–60.5	26.8–51.5	26.3–48.9	15.5–64.3	–	33.5–78.9	30.6–58.4	13.3–40.4
Got from a friend, relative, or someone you know for free	%	63.2	55.3	68.7	62.6	56.3	0.0	68.4	66.0	58.5
	CI	53.6–71.9	38.2–71.2	55.6–79.4	50.4–73.3	29.7–79.8	–	42.7–86.3	51.4–78.1	42.5–73.0
Took from a friend, relative, or someone you know without asking	%	9.4	0.0	14.3	10.9	0.0	0.0	21.1	10.0	4.9
	CI	5.2–16.5	–	7.4–25.8	5.4–20.7	–	–	7.3–47.3	4.1–22.4	1.1–18.4
Took from someone's house or workplace without asking	%	9.1	3.2	8.4	4.2	14.4	0.0	10.5	8.0	9.8
	CI	4.9–16.3	0.4–21.0	3.4–19.1	1.3–12.7	3.1–47.1	–	2.3–37.1	2.9–20.0	3.6–24.1
Got it at a party	%	29.6	32.8	26.6	25.6	30.5	0.0	47.4	36.0	17.1
	CI	21.8–38.9	19.0–50.4	16.9–39.2	16.7–37.1	11.8–59.2	–	25.0–70.8	23.6–50.6	8.1–32.5
Got it some other way	%	22.9	16.1	18.5	21.5	19.8	0.0	42.1	22.0	17.1
	CI	15.9–31.7	7.1–32.7	10.6–30.3	13.3–32.7	5.6–50.5	–	21.1–66.5	12.4–36.1	8.1–32.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.36. Availability of Alcohol – Sources

During the past 30 days, if you used alcohol, in which ways (if any) did you get alcohol? (Mark all that apply)										
		Total (n=597)	Male (n=258)	Female (n=312)	White (n=434)	Black (n=73)	Hispanic (n=9)	8th (n=116)	10th (n=231)	12th (n=250)
I bought it in a store	%	4.3	4.9	2.6	3.1	5.7	12.7	2.6	3.9	5.2
	CI	2.9–6.3	2.8–8.4	1.3–5.1	1.8–5.3	2.1–14.5	1.1–66.0	0.8–7.8	2.0–7.3	3.0–8.7
I bought it in a gas station	%	5.1	5.0	4.8	4.3	7.2	0.0	3.4	4.7	6.0
	CI	3.6–7.2	2.9–8.5	2.8–7.9	2.7–6.7	3.0–16.5	–	1.3–8.9	2.6–8.4	3.6–9.7
I bought it in a restaurant	%	1.5	1.6	0.7	0.9	3.1	12.7	1.7	0.9	2.0
	CI	0.8–2.9	0.6–4.3	0.2–2.7	0.3–2.5	0.7–11.8	1.1–66.0	0.4–6.8	0.2–3.4	0.8–4.7
I bought it in a bar or club	%	1.2	0.9	1.7	0.7	4.6	0.0	0.9	0.4	2.0
	CI	0.6–2.6	0.2–3.4	0.7–4.0	0.2–2.3	1.5–13.7	–	0.1–6.0	0.1–3.0	0.8–4.7
I bought it at a public event	%	3.2	4.2	1.6	2.5	5.9	0.0	2.6	3.4	3.2
	CI	2.0–4.9	2.3–7.6	0.7–3.9	1.4–4.5	2.2–15.0	–	0.8–7.8	1.7–6.7	1.6–6.2
I gave someone money to buy it for me	%	27.2	28.7	25.8	30.1	17.0	12.7	14.7	20.2	36.9
	CI	23.7–31.0	23.4–34.6	21.1–31.1	25.8–34.6	9.8–27.9	1.1–66.0	9.2–22.5	15.5–25.9	31.1–43.1
I got it from a sibling (under 21)	%	5.4	4.7	5.6	4.6	11.4	0.0	8.6	6.4	3.6
	CI	3.9–7.5	2.8–8.0	3.5–8.8	3.0–7.0	5.9–20.9	–	4.7–15.4	3.9–10.4	1.9–6.7
I got it from a friend (under 21)	%	22.9	18.5	26.2	23.1	18.1	21.1	29.3	23.6	20.2
	CI	19.7–26.5	14.2–23.7	21.6–31.4	19.3–27.3	10.6–29.1	3.6–65.5	21.7–38.4	18.6–29.5	15.7–25.7
I got it at a party	%	42.3	41.9	43.1	41.7	44.8	43.1	31.0	36.9	50.4
	CI	38.4–46.4	35.9–48.1	37.6–48.7	37.1–46.4	33.5–56.7	12.7–79.8	23.2–40.1	30.9–43.3	44.2–56.6
I took it from a store or family member	%	13.8	13.5	12.6	12.2	16.3	19.9	19.8	17.6	8.7
	CI	11.2–16.7	9.8–18.2	9.4–16.7	9.4–15.6	9.3–26.9	3.4–64.0	13.5–28.2	13.2–23.1	5.8–12.9
I took it home without permission	%	15.0	13.5	15.3	14.3	11.9	30.5	21.6	20.2	8.7
	CI	12.4–18.0	9.8–18.2	11.7–19.6	11.3–17.9	6.2–21.8	7.2–71.3	14.9–30.1	15.5–25.9	5.8–12.9
I got it from my parents with permission	%	23.1	19.5	25.6	24.1	18.5	19.9	31.9	27.0	17.1
	CI	19.9–26.6	15.1–24.8	21.0–30.7	20.3–28.3	11.1–29.2	3.4–64.0	24.0–41.0	21.7–33.1	12.9–22.3
I got it from an older sibling (over 21)	%	11.3	10.8	11.8	10.2	14.5	12.7	10.3	11.6	11.5
	CI	9.0–14.2	7.5–15.3	8.7–16.0	7.7–13.5	8.0–24.7	1.1–66.0	5.9–17.5	8.0–16.4	8.1–16.1
I got it from an older friend (over 21)	%	23.0	23.4	23.5	25.3	12.9	33.7	17.2	21.9	25.8
	CI	19.7–26.6	18.6–29.1	19.0–28.6	21.3–29.7	7.0–22.8	8.2–74.3	11.3–25.4	17.0–27.7	20.7–31.6
I got it from some other adult	%	17.4	14.2	19.7	17.2	19.6	21.1	20.7	18.0	15.9
	CI	14.6–20.7	10.4–19.1	15.6–24.5	13.9–21.0	12.0–30.5	3.6–65.5	14.2–29.2	13.6–23.5	11.8–21.0
I got it some other way	%	23.2	27.8	17.5	20.5	28.9	46.4	25.9	23.2	22.2
	CI	19.9–26.7	22.6–33.6	13.7–22.1	17.0–24.6	19.4–40.6	14.3–81.8	18.6–34.7	18.2–29.1	17.5–27.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.37. Asked to Show Proof of Age to Buy Alcohol

If you bought or tried to buy alcohol yourself during the past 30 days, were you ever asked to show proof of age?				
		I bought or tried to buy alcohol	Yes, asked for proof of age	No, not asked for proof of age
Total	%	4.4	23.3	76.7
(n=122)	CI	3.7–5.2	16.6–31.8	68.2–83.4
Male	%	4.9	28.4	71.6
(n=67)	CI	3.8–6.1	18.7–40.8	59.2–81.3
Female	%	3.2	19.2	80.8
(n=42)	CI	2.4–4.2	9.6–34.8	65.2–90.4
White	%	3.4	27.4	72.6
(n=67)	CI	2.7–4.3	17.7–39.7	60.3–82.3
Black	%	5.4	21.8	78.2
(n=23)	CI	3.6–7.9	8.7–45.1	54.9–91.3
Hispanic	%	3.9	45.4	54.6
(n=2)	CI	0.9–14.8	0.0–100.0	0.0–100.0
8th	%	2.4	25.0	75.0
(n=27)	CI	1.6–3.4	11.8–45.3	54.7–88.2
10 th	%	4.1	19.0	81.0
(n=42)	CI	3.0–5.4	9.5–34.4	65.6–90.5
12th	%	6.8	25.5	74.5
(n=53)	CI	5.3–8.7	15.4–39.0	61.0–84.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.38. Ability to Purchase Alcohol

If you tried to buy alcohol yourself during the past 30 days, were you able to purchase it?					
		I bought or tried to buy alcohol	No, not able to purchase	Yes, able to purchase sometimes	Yes, able to purchase every time
Total	%	4.4	32.7	35.7	31.6
(n=122)	CI	3.7–5.2	24.9–41.5	27.7–44.5	24.0–40.4
Male	%	4.9	33.4	44.6	22.1
(n=67)	CI	3.8–6.1	22.8–45.9	32.8–56.9	13.5–33.9
Female	%	3.2	33.5	32.9	33.5
(n=42)	CI	2.4–4.2	20.9–49.1	20.4–48.4	20.9–49.1
White	%	3.4	24.3	40.6	35.2
(n=67)	CI	2.7–4.3	15.4–36.1	29.3–52.8	24.6–47.5
Black	%	5.4	51.5	31.9	16.5
(n=23)	CI	3.6–7.9	30.9–71.7	15.8–54.1	5.8–38.8
Hispanic	%	3.9	45.4	54.6	0.0
(n=2)	CI	0.9–14.8	0.0–100.0	0.0–100.0	–
8th	%	2.4	32.1	35.7	32.1
(n=27)	CI	1.6–3.4	16.9–52.5	19.6–55.9	16.9–52.5
10 th	%	4.1	26.2	45.2	28.6
(n=42)	CI	3.0–5.4	14.8–42.1	30.5–60.9	16.6–44.6
12th	%	6.8	36.8	29.8	33.3
(n=53)	CI	5.3–8.7	25.1–50.4	19.1–43.3	22.1–46.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.39. Ever Ridden with a Prescription Drugs Impaired Driver

		Have you ever ridden in a car driven by someone who was taking or was under the influence of prescription drugs?		
		No	Yes	Not sure
Total	%	77.5	11.1	11.5
(n=3,347)	CI	76.0–78.9	10.0–12.2	10.4–12.6
Male	%	79.9	8.9	11.2
(n=1,627)	CI	77.8–81.7	7.6–10.4	9.8–12.8
Female	%	76.4	12.1	11.5
(n=1,611)	CI	74.2–78.4	10.6–13.8	10.1–13.2
White	%	78.2	10.4	11.4
(n=2,296)	CI	76.5–79.9	9.2–11.8	10.1–12.7
Black	%	80.4	9.6	9.9
(n=551)	CI	76.9–83.6	7.4–12.4	7.7–12.7
Hispanic	%	86.2	4.7	9.1
(n=63)	CI	75.1–92.9	1.5–14.1	4.0–19.2
8th	%	79.0	7.4	13.7
(n=1,273)	CI	76.6–81.1	6.1–9.0	11.9–15.7
10th	%	73.7	12.4	13.9
(n=1,163)	CI	71.1–76.1	10.7–14.5	12.0–16.0
12th	%	80.0	13.4	6.6
(n=911)	CI	77.2–82.4	11.3–15.8	5.2–8.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.40. Frequency Riding with a Prescription Drugs Impaired Driver

		During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was taking or was under the influence of prescription drugs?					
		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	93.4	2.2	1.7	0.9	1.8	6.6
(n=3,402)	CI	92.5–94.2	1.7–2.7	1.3–2.2	0.6–1.3	1.4–2.3	5.8–7.5
Male	%	94.8	2.1	1.2	0.8	1.1	5.2
(n=1,658)	CI	93.6–95.8	1.5–2.9	0.8–1.9	0.5–1.4	0.7–1.7	4.2–6.4
Female	%	92.9	2.4	2.0	0.9	1.8	7.1
(n=1,633)	CI	91.6–94.1	1.7–3.3	1.4–2.8	0.6–1.5	1.2–2.5	5.9–8.4
White	%	94.5	2.1	1.4	0.6	1.5	5.5
(n=2,322)	CI	93.5–95.4	1.6–2.7	1.0–1.9	0.4–1.0	1.0–2.1	4.6–6.5
Black	%	93.3	2.3	1.9	0.7	1.8	6.7
(n=572)	CI	90.8–95.1	1.4–4.0	1.0–3.4	0.3–1.9	1.0–3.3	4.9–9.2
Hispanic	%	0.0	0.0	0.0	0.0	0.0	0.0
(n=64)	CI	–	–	–	–	–	–
8th	%	95.0	1.6	1.7	0.6	1.2	5.0
(n=1,288)	CI	93.6–96.0	1.0–2.4	1.1–2.6	0.3–1.2	0.7–1.9	4.0–6.4
10th	%	93.1	2.9	1.6	1.0	1.4	6.9
(n=1,182)	CI	91.5–94.4	2.1–4.0	1.0–2.5	0.6–1.8	0.9–2.3	5.6–8.5
12th	%	92.2	2.0	1.8	1.1	2.9	7.8
(n=932)	CI	90.3–93.8	1.3–3.2	1.1–2.9	0.6–2.0	2.0–4.2	6.2–9.7

Note: The confidence interval (CI) is 95 percent.

Table BH6.41. Ever Ridden with an Alcohol or Drug Impaired Driver

Have you ever ridden in a car driven by someone who was intoxicated by alcohol or drugs?		No	Yes	Not sure
Total	%	64.1	25.8	10.1
(n=3,419)	CI	62.5–65.7	24.3–27.3	9.1–11.1
Male	%	66.1	22.8	11.0
(n=1,666)	CI	63.8–68.4	20.9–24.9	9.6–12.6
Female	%	63.2	28.2	8.5
(n=1,643)	CI	60.8–65.5	26.1–30.5	7.3–10.0
White	%	65.0	24.7	10.2
(n=2,333)	CI	63.0–66.9	23.0–26.6	9.1–11.5
Black	%	68.6	23.2	8.2
(n=571)	CI	64.6–72.3	19.9–26.9	6.2–10.7
Hispanic	%	57.7	29.2	13.0
(n=66)	CI	45.2–69.3	19.2–41.8	6.8–23.6
8th	%	67.9	19.8	12.3
(n=1,298)	CI	65.3–70.4	17.7–22.0	10.6–14.2
10th	%	61.2	27.3	11.5
(n=1,188)	CI	58.4–63.9	24.8–29.9	9.8–13.4
12th	%	63.3	30.4	6.3
(n=933)	CI	60.2–66.3	27.6–33.4	4.9–8.0

Note: The confidence interval (CI) is 95 percent.

Table BH6.42. Frequency Riding with an Alcohol or Drug Impaired Driver

During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who was intoxicated by alcohol or drugs?		0 times	1 time	2–3 times	4–5 times	6 or more times	Combined: (One or more)
Total	%	84.8	5.4	4.9	1.5	3.4	15.2
(n=3,418)	CI	83.6–86.0	4.6–6.2	4.2–5.7	1.1–2.0	2.8–4.1	14.0–16.4
Male	%	87.1	4.3	4.3	1.6	2.8	12.9
(n=1,664)	CI	85.4–88.6	3.4–5.4	3.4–5.4	1.1–2.3	2.1–3.7	11.4–14.6
Female	%	83.3	6.6	5.4	1.5	3.2	16.7
(n=1,643)	CI	81.4–85.1	5.5–7.9	4.4–6.6	1.1–2.3	2.4–4.2	14.9–18.6
White	%	86.0	5.0	5.0	1.1	2.9	14.0
(n=2,332)	CI	84.5–87.3	4.2–6.0	4.2–6.0	0.8–1.7	2.3–3.7	12.7–15.5
Black	%	84.9	5.5	3.4	2.0	4.2	15.1
(n=575)	CI	81.6–87.6	3.9–7.7	2.2–5.3	1.1–3.6	2.8–6.3	12.4–18.4
Hispanic	%	83.3	8.8	3.7	2.9	1.4	16.7
(n=65)	CI	71.8–90.7	3.9–18.7	0.9–13.9	0.7–11.2	0.2–9.5	9.3–28.2
8th	%	88.6	4.5	4.0	1.2	1.8	11.4
(n=1,296)	CI	86.7–90.2	3.5–5.7	3.1–5.2	0.7–1.9	1.2–2.7	9.8–13.3
10th	%	84.1	5.5	5.2	1.7	3.5	15.9
(n=1,187)	CI	81.9–86.1	4.3–6.9	4.1–6.6	1.1–2.6	2.6–4.7	13.9–18.1
12th	%	81.7	6.2	5.4	1.7	5.0	18.3
(n=935)	CI	79.1–84.1	4.8–7.9	4.1–7.1	1.0–2.8	3.8–6.6	15.9–20.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.43. Parents Feel Misusing Prescription Drugs is Wrong

		How wrong do your parents feel it would be for you to <u>use prescription drugs not prescribed to you</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.8	3.3	6.1	87.8	93.9
(n=3,410)	CI	2.3–3.4	2.8–4.0	5.3–6.9	86.6–88.9	93.0–94.6
Male	%	2.7	2.4	5.9	89.0	94.9
(n=1,665)	CI	2.0–3.6	1.8–3.3	4.9–7.2	87.3–90.4	93.7–95.9
Female	%	2.2	4.0	6.3	87.5	93.8
(n=1,633)	CI	1.6–3.0	3.1–5.1	5.3–7.6	85.8–89.0	92.5–94.9
White	%	1.7	3.5	5.8	89.0	94.8
(n=2,328)	CI	1.2–2.3	2.8–4.3	4.9–6.8	87.7–90.2	93.8–95.7
Black	%	4.8	2.1	6.4	86.7	93.1
(n=568)	CI	3.3–6.9	1.2–3.6	4.6–8.8	83.6–89.3	90.7–95.0
Hispanic	%	0.0	1.5	5.7	92.8	98.5
(n=66)	CI	–	0.2–10.4	2.1–14.6	83.5–97.1	89.6–99.8
8th	%	2.1	2.5	5.0	90.3	95.4
(n=1,294)	CI	1.4–3.0	1.8–3.6	4.0–6.4	88.6–91.8	94.1–96.4
10th	%	2.8	4.1	6.3	86.8	93.1
(n=1,184)	CI	2.0–3.9	3.1–5.4	5.1–7.9	84.7–88.6	91.5–94.4
12th	%	3.5	3.3	6.9	86.3	93.2
(n=932)	CI	2.5–4.9	2.3–4.7	5.5–8.7	83.9–88.3	91.4–94.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.44. Parents Feel Drinking Alcohol is Wrong

		How wrong do your parents feel it would be for you to <u>drink alcohol</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.0	5.0	11.6	80.4	92.0
(n=3,414)	CI	2.5–3.7	4.3–5.8	10.5–12.7	79.0–81.7	91.0–92.9
Male	%	2.8	4.9	12.5	79.8	92.3
(n=1,668)	CI	2.1–3.8	3.9–6.0	11.0–14.2	77.7–81.7	90.9–93.5
Female	%	2.6	4.9	10.4	82.2	92.6
(n=1,634)	CI	1.9–3.5	3.9–6.0	9.0–12.0	80.3–84.0	91.2–93.8
White	%	2.4	5.3	12.4	79.9	92.3
(n=2,333)	CI	1.8–3.1	4.4–6.3	11.1–13.8	78.2–81.5	91.2–93.4
Black	%	3.7	3.3	7.4	85.6	93.0
(n=568)	CI	2.4–5.6	2.1–5.2	5.5–9.9	82.4–88.3	90.5–94.9
Hispanic	%	0.0	1.5	12.6	85.9	98.5
(n=66)	CI	–	0.2–10.4	6.3–23.7	74.6–92.6	89.6–99.8
8th	%	1.9	4.1	8.1	85.9	94.1
(n=1,294)	CI	1.2–2.8	3.1–5.3	6.7–9.7	83.9–87.7	92.6–95.2
10th	%	2.4	5.2	12.9	79.6	92.4
(n=1,187)	CI	1.6–3.4	4.1–6.6	11.1–14.9	77.2–81.8	90.8–93.8
12th	%	4.9	5.7	13.8	75.5	89.4
(n=933)	CI	3.7–6.5	4.4–7.4	11.8–16.2	72.7–78.2	87.2–91.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.45. Parents Feel Regularly Drinking Alcohol is Wrong

How wrong do your parents feel it would be for you to <u>drink beer, wine, or hard liquor (for example, vodka, whiskey, or gin) regularly?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	6.8	9.8	12.7	70.7	83.4
(n=3,414)	CI	6.0–7.7	8.8–10.9	11.6–13.9	69.1–72.2	82.1–84.6
Male	%	6.9	9.1	13.2	70.9	84.1
(n=1,668)	CI	5.7–8.2	7.7–10.6	11.6–14.9	68.7–73.1	82.2–85.8
Female	%	6.1	10.2	12.3	71.4	83.7
(n=1,634)	CI	5.0–7.4	8.8–11.8	10.8–14.0	69.1–73.6	81.8–85.4
White	%	6.6	10.3	13.4	69.8	83.2
(n=2,333)	CI	5.6–7.7	9.1–11.6	12.1–14.9	67.8–71.6	81.6–84.7
Black	%	5.4	6.1	9.3	79.2	88.5
(n=567)	CI	3.8–7.6	4.4–8.5	7.2–12.0	75.6–82.3	85.5–90.9
Hispanic	%	4.6	11.3	11.8	72.2	84.1
(n=66)	CI	1.5–13.9	5.3–22.2	5.9–22.4	59.8–82.0	72.5–91.4
8th	%	4.0	7.0	11.0	77.9	89.0
(n=1,295)	CI	3.1–5.2	5.8–8.5	9.4–12.9	75.6–80.1	87.1–90.6
10th	%	6.8	11.3	13.5	68.3	81.8
(n=1,187)	CI	5.5–8.4	9.7–13.3	11.7–15.6	65.6–70.9	79.6–83.9
12th	%	9.6	11.1	13.5	65.8	79.3
(n=932)	CI	7.9–11.6	9.2–13.3	11.5–15.9	62.7–68.8	76.6–81.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.46. Parents Feel Smoking Tobacco is Wrong

How wrong do your parents feel it would be for you to <u>smoke tobacco?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	3.2	3.8	10.4	82.6	93.0
(n=3,415)	CI	2.6–3.9	3.2–4.5	9.4–11.5	81.3–83.9	92.1–93.8
Male	%	3.6	4.7	11.0	80.7	91.7
(n=1,672)	CI	2.8–4.7	3.7–5.8	9.6–12.7	78.7–82.5	90.3–93.0
Female	%	2.3	2.7	9.4	85.6	95.0
(n=1,631)	CI	1.6–3.2	2.0–3.6	8.0–10.9	83.8–87.3	93.8–96.0
White	%	2.7	4.1	10.9	82.2	93.2
(n=2,330)	CI	2.1–3.5	3.3–5.0	9.7–12.3	80.6–83.8	92.0–94.2
Black	%	4.0	1.7	7.2	87.2	94.4
(n=570)	CI	2.6–6.0	0.9–3.2	5.3–9.6	84.2–89.7	92.1–96.0
Hispanic	%	0.0	1.5	14.7	83.8	98.5
(n=66)	CI	–	0.2–10.4	7.7–26.3	72.0–91.2	89.6–99.8
8th	%	1.5	2.5	7.9	88.0	96.0
(n=1,296)	CI	0.9–2.3	1.8–3.6	6.6–9.5	86.2–89.7	94.8–96.9
10th	%	2.3	3.1	9.8	84.9	94.6
(n=1,186)	CI	1.6–3.3	2.3–4.3	8.2–11.6	82.7–86.8	93.2–95.8
12th	%	6.0	5.9	13.5	74.7	88.2
(n=933)	CI	4.6–7.7	4.5–7.5	11.5–15.9	71.8–77.4	86.0–90.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.47. Parents Feel Smoking One or More Packs of Cigarettes is Wrong

How wrong do your parents feel it would be for you to <u>smoke one or more packs of cigarettes a day</u> ?						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	2.4	1.7	6.3	89.6	95.9
(n=3,412)	CI	2.0–3.0	1.3–2.2	5.5–7.2	88.5–90.6	95.1–96.5
Male	%	2.7	2.2	7.1	88.0	95.1
(n=1,664)	CI	2.0–3.7	1.6–3.0	6.0–8.5	86.3–89.5	93.9–96.0
Female	%	1.6	1.0	5.5	92.0	97.4
(n=1,637)	CI	1.0–2.3	0.6–1.7	4.4–6.7	90.5–93.2	96.5–98.1
White	%	1.8	2.0	6.4	89.8	96.2
(n=2,331)	CI	1.3–2.5	1.5–2.6	5.5–7.5	88.5–91.0	95.3–97.0
Black	%	3.4	0.2	5.5	90.9	96.4
(n=569)	CI	2.2–5.3	0.0–1.1	3.9–7.8	88.2–93.0	94.5–97.7
Hispanic	%	1.5	1.5	6.0	91.0	97.0
(n=66)	CI	0.2–10.4	0.2–10.4	2.2–15.4	81.0–96.0	88.4–99.3
8th	%	1.4	0.9	4.3	93.4	97.7
(n=1,293)	CI	0.9–2.2	0.5–1.6	3.3–5.5	91.9–94.7	96.7–98.4
10th	%	1.9	1.2	5.6	91.4	97.0
(n=1,186)	CI	1.2–2.8	0.7–2.0	4.4–7.0	89.7–92.9	95.8–97.8
12th	%	4.1	3.0	9.1	83.7	92.9
(n=933)	CI	3.0–5.6	2.1–4.3	7.5–11.2	81.2–86.0	91.0–94.4

Note: The confidence interval (CI) is 95 percent.

Table BH6.48. Parents Feel Using Electronic Cigarettes is Wrong

How wrong do your parents feel it would be for you to <u>use an e-cigarette</u> ?						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.2	8.5	13.2	73.1	86.3
(n=3,418)	CI	4.5–6.0	7.6–9.5	12.1–14.4	71.6–74.6	85.1–87.4
Male	%	5.7	8.4	14.5	71.5	85.9
(n=1,671)	CI	4.7–7.0	7.1–9.8	12.8–16.3	69.2–73.6	84.1–87.5
Female	%	3.9	8.5	11.7	76.0	87.6
(n=1,635)	CI	3.0–5.0	7.2–10.0	10.2–13.3	73.8–78.0	85.9–89.2
White	%	4.8	9.5	14.7	71.0	85.7
(n=2,334)	CI	4.0–5.8	8.4–10.8	13.3–16.2	69.1–72.8	84.2–87.1
Black	%	4.1	3.5	7.4	84.9	92.3
(n=570)	CI	2.7–6.2	2.3–5.5	5.5–9.9	81.7–87.7	89.8–94.3
Hispanic	%	0.0	8.7	15.1	76.1	91.3
(n=66)	CI	–	3.6–19.6	8.2–26.3	63.8–85.3	80.4–96.4
8th	%	3.2	6.2	10.9	79.6	90.5
(n=1,296)	CI	2.4–4.4	5.0–7.7	9.3–12.7	77.4–81.8	88.8–92.0
10th	%	3.7	8.9	14.9	72.5	87.4
(n=1,186)	CI	2.8–4.9	7.4–10.7	13.0–17.0	69.9–75.0	85.4–89.2
12th	%	8.8	10.4	13.8	67.0	80.8
(n=936)	CI	7.2–10.8	8.6–12.5	11.7–16.1	63.9–70.0	78.2–83.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.49. Parents Feel Smoking Marijuana is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.7	5.7	8.3	80.3	88.6
(n=3,411)	CI	4.9–6.5	5.0–6.6	7.4–9.3	78.9–81.7	87.5–89.6
Male	%	6.0	5.9	8.4	79.6	88.1
(n=1,666)	CI	4.9–7.3	4.9–7.2	7.2–9.9	77.6–81.5	86.4–89.6
Female	%	4.6	5.3	7.8	82.3	90.1
(n=1,632)	CI	3.7–5.8	4.3–6.5	6.6–9.2	80.4–84.1	88.5–91.5
White	%	4.4	5.5	8.3	81.8	90.1
(n=2,326)	CI	3.7–5.4	4.6–6.5	7.2–9.5	80.1–83.3	88.8–91.2
Black	%	6.9	4.4	7.1	81.5	88.6
(n=571)	CI	5.1–9.4	3.0–6.5	5.3–9.6	78.0–84.5	85.7–91.0
Hispanic	%	1.8	3.3	10.3	84.5	94.9
(n=66)	CI	0.2–12.3	0.8–12.7	4.9–20.6	73.2–91.6	84.9–98.4
8th	%	3.0	3.1	5.8	88.1	93.9
(n=1,295)	CI	2.2–4.1	2.3–4.2	4.6–7.2	86.2–89.8	92.5–95.1
10th	%	6.1	7.5	8.4	78.0	86.4
(n=1,183)	CI	4.8–7.6	6.1–9.2	7.0–10.2	75.5–80.3	84.4–88.3
12th	%	8.0	6.6	10.6	74.8	85.4
(n=933)	CI	6.4–9.9	5.2–8.4	8.8–12.8	71.9–77.5	83.0–87.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.50. Parents Feel Smoking Marijuana Once or Twice a Week is Wrong

		How wrong do your parents feel it would be for you to <u>smoke marijuana once or twice a week</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.4	4.4	7.3	82.9	90.2
(n=3,410)	CI	4.7–6.3	3.7–5.1	6.5–8.3	81.5–84.1	89.1–91.2
Male	%	5.5	4.8	7.9	81.8	89.7
(n=1,669)	CI	4.5–6.8	3.8–5.9	6.7–9.3	79.8–83.6	88.1–91.1
Female	%	4.7	3.7	6.7	84.9	91.6
(n=1,630)	CI	3.8–5.9	2.8–4.8	5.6–8.1	83.0–86.5	90.1–92.9
White	%	4.2	4.0	7.1	84.7	91.9
(n=2,329)	CI	3.4–5.1	3.2–4.9	6.2–8.3	83.2–86.2	90.7–93.0
Black	%	7.3	3.9	6.2	82.6	88.8
(n=571)	CI	5.4–9.8	2.6–5.9	4.5–8.6	79.2–85.5	85.9–91.2
Hispanic	%	3.4	1.5	9.2	85.9	95.1
(n=65)	CI	0.8–13.0	0.2–10.6	4.1–19.4	74.7–92.7	85.5–98.5
8th	%	2.7	1.8	5.8	89.7	95.5
(n=1,296)	CI	1.9–3.7	1.2–2.7	4.6–7.2	88.0–91.3	94.3–96.5
10th	%	5.8	5.8	7.7	80.7	88.4
(n=1,183)	CI	4.6–7.3	4.6–7.3	6.3–9.3	78.3–82.8	86.4–90.1
12th	%	7.9	5.5	8.5	78.0	86.6
(n=931)	CI	6.3–9.8	4.2–7.2	6.9–10.5	75.3–80.6	84.2–88.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.51. Friends Feel Misusing Prescription Drugs is Wrong

How wrong do your friends feel it would be for you to <u>use prescription drugs not prescribed to you?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	10.7	9.2	15.8	64.2	80.1
(n=3,405)	CI	9.7–11.8	8.3–10.3	14.6–17.1	62.6–65.9	78.7–81.4
Male	%	10.8	9.1	16.4	63.7	80.1
(n=1,664)	CI	9.4–12.4	7.8–10.6	14.7–18.3	61.3–66.0	78.1–82.0
Female	%	9.6	9.2	15.5	65.7	81.2
(n=1,629)	CI	8.2–11.1	7.9–10.8	13.8–17.4	63.3–68.0	79.2–83.1
White	%	9.8	9.7	16.3	64.3	80.6
(n=2,326)	CI	8.6–11.1	8.5–11.0	14.8–17.9	62.3–66.2	78.9–82.1
Black	%	10.4	9.1	13.6	66.9	80.6
(n=569)	CI	8.1–13.2	6.9–11.8	11.1–16.7	62.9–70.7	77.1–83.6
Hispanic	%	5.5	1.5	18.6	74.4	93.0
(n=65)	CI	1.7–16.1	0.2–10.5	10.7–30.4	61.9–83.9	82.4–97.4
8th	%	7.3	6.5	13.1	73.1	86.3
(n=1,294)	CI	6.0–8.8	5.3–8.0	11.4–15.1	70.6–75.5	84.3–88.0
10th	%	11.2	10.4	18.2	60.2	78.4
(n=1,178)	CI	9.5–13.1	8.8–12.3	16.1–20.5	57.4–63.0	76.0–80.7
12th	%	13.6	10.9	16.2	59.4	75.5
(n=933)	CI	11.6–16.0	9.0–13.0	13.9–18.7	56.2–62.5	72.7–78.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.52. Friends Feel Drinking Alcohol is Wrong

How wrong do your friends feel it would be for you to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	18.7	16.6	19.8	44.9	64.7
(n=3,410)	CI	17.4–20.1	15.4–17.9	18.5–21.2	43.2–46.5	63.0–66.3
Male	%	19.4	16.5	20.6	43.5	64.0
(n=1,666)	CI	17.6–21.4	14.8–18.4	18.7–22.6	41.1–45.9	61.7–66.3
Female	%	17.1	16.5	19.4	47.0	66.4
(n=1,632)	CI	15.3–19.0	14.8–18.5	17.5–21.4	44.6–49.4	64.0–68.6
White	%	18.5	17.4	21.3	42.8	64.0
(n=2,327)	CI	17.0–20.2	15.9–19.1	19.7–23.0	40.7–44.8	62.0–66.0
Black	%	16.5	12.1	16.7	54.7	71.4
(n=571)	CI	13.6–19.8	9.7–15.1	13.8–20.0	50.6–58.8	67.5–75.0
Hispanic	%	18.5	10.7	19.9	50.9	70.9
(n=65)	CI	10.4–30.7	5.0–21.1	11.7–31.8	38.6–63.2	58.1–81.0
8th	%	11.0	11.7	19.8	57.5	77.3
(n=1,294)	CI	9.4–12.8	10.1–13.6	17.7–22.0	54.8–60.2	74.9–79.5
10th	%	19.6	21.5	19.9	39.0	58.9
(n=1,182)	CI	17.4–21.9	19.3–24.0	17.7–22.3	36.2–41.8	56.1–61.7
12th	%	25.7	16.5	19.8	38.0	57.8
(n=934)	CI	23.0–28.6	14.2–19.0	17.3–22.4	35.0–41.2	54.6–60.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.53. Friends Feel Smoking Tobacco is Wrong

		How wrong do your friends feel it would be for you to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	16.1	13.8	19.1	50.9	70.1
(n=3,402)	CI	14.9–17.4	12.7–15.0	17.8–20.5	49.2–52.6	68.5–71.6
Male	%	17.0	15.1	18.8	49.0	67.9
(n=1,661)	CI	15.2–18.9	13.5–16.9	17.0–20.8	46.6–51.4	65.5–70.1
Female	%	14.1	12.5	19.9	53.5	73.5
(n=1,629)	CI	12.4–15.9	10.9–14.2	18.0–22.0	51.1–56.0	71.2–75.6
White	%	16.8	15.8	20.3	47.1	67.4
(n=2,324)	CI	15.3–18.4	14.3–17.3	18.7–22.0	45.1–49.1	65.4–69.3
Black	%	10.9	7.5	15.5	66.1	81.6
(n=568)	CI	8.6–13.7	5.6–10.0	12.8–18.8	62.0–69.9	78.2–84.6
Hispanic	%	8.5	8.1	18.4	65.0	83.4
(n=65)	CI	3.5–19.3	3.3–18.4	10.6–30.1	52.1–75.9	71.4–91.0
8th	%	10.3	9.9	17.5	62.3	79.8
(n=1,292)	CI	8.7–12.1	8.4–11.7	15.5–19.6	59.7–64.9	77.5–81.9
10th	%	15.3	15.5	20.7	48.4	69.2
(n=1,178)	CI	13.4–17.5	13.5–17.7	18.5–23.2	45.6–51.3	66.5–71.8
12th	%	22.9	16.1	19.2	41.9	61.0
(n=932)	CI	20.3–25.7	13.9–18.6	16.8–21.8	38.7–45.0	57.9–64.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.54. Friends Feel Using Electronic Cigarettes is Wrong

		How wrong do your friends feel it would be for you to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	30.4	15.6	13.0	41.0	54.0
(n=3,390)	CI	28.9–32.0	14.4–16.9	11.9–14.2	39.3–42.6	52.3–55.7
Male	%	31.4	15.2	13.2	40.2	53.4
(n=1,652)	CI	29.2–33.7	13.5–17.0	11.6–14.9	37.9–42.6	51.0–55.8
Female	%	28.4	16.2	13.2	42.1	55.4
(n=1,626)	CI	26.2–30.7	14.5–18.1	11.7–15.0	39.8–44.6	52.9–57.8
White	%	33.4	17.4	13.1	36.1	49.2
(n=2,319)	CI	31.4–35.3	15.9–19.0	11.8–14.6	34.2–38.1	47.2–51.3
Black	%	16.5	8.6	13.8	61.1	74.9
(n=564)	CI	13.6–19.8	6.5–11.2	11.2–16.9	57.0–65.1	71.1–78.3
Hispanic	%	20.7	13.4	9.0	56.9	65.9
(n=65)	CI	12.2–32.9	6.7–25.0	4.0–19.0	44.2–68.7	53.0–76.8
8th	%	19.6	13.1	15.1	52.2	67.3
(n=1,291)	CI	17.5–21.8	11.3–15.0	13.2–17.2	49.5–55.0	64.7–69.8
10th	%	31.4	19.1	12.7	36.8	49.5
(n=1,170)	CI	28.8–34.1	16.9–21.5	10.9–14.7	34.1–39.6	46.7–52.4
12th	%	40.6	14.5	11.2	33.7	44.9
(n=929)	CI	37.5–43.8	12.4–16.9	9.3–13.4	30.7–36.8	41.7–48.1

Note: The confidence interval (CI) is 95 percent.

Table BH6.55. Friends Feel Smoking Marijuana is Wrong

		How wrong do your friends feel it would be for you to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	25.7	13.9	13.1	47.4	60.4
(n=3,405)	CI	24.2–27.2	12.7–15.1	12.0–14.3	45.7–49.1	58.8–62.1
Male	%	26.1	14.0	13.4	46.5	59.8
(n=1,666)	CI	24.1–28.3	12.4–15.8	11.8–15.1	44.1–48.9	57.4–62.2
Female	%	23.8	14.2	12.9	49.1	62.0
(n=1,628)	CI	21.7–26.0	12.5–16.0	11.4–14.7	46.7–51.5	59.6–64.4
White	%	24.1	14.4	14.1	47.4	61.5
(n=2,324)	CI	22.3–25.9	13.0–15.9	12.7–15.6	45.4–49.5	59.5–63.5
Black	%	24.8	13.0	11.6	50.7	62.3
(n=572)	CI	21.3–28.5	10.4–16.0	9.2–14.5	46.5–54.8	58.2–66.2
Hispanic	%	18.7	13.3	10.7	57.3	68.0
(n=64)	CI	10.5–31.1	6.7–24.8	5.0–21.2	44.5–69.2	54.9–78.7
8th	%	13.8	8.1	13.0	65.1	78.1
(n=1,293)	CI	12.1–15.8	6.7–9.7	11.3–14.9	62.4–67.6	75.7–80.2
10th	%	29.9	16.2	13.8	40.1	53.8
(n=1,180)	CI	27.4–32.6	14.2–18.4	11.9–15.9	37.3–42.9	51.0–56.7
12th	%	33.3	17.4	12.5	36.8	49.3
(n=932)	CI	30.4–36.4	15.1–19.9	10.5–14.7	33.8–40.0	46.1–52.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.56. Friends Misused Prescription Drugs

		Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have <u>used prescription drugs not prescribed to you</u> ?					
		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	84.7	6.5	3.1	1.3	4.4	15.3
(n=3,394)	CI	83.4–85.9	5.7–7.4	2.6–3.8	0.9–1.7	3.7–5.2	14.1–16.6
Male	%	87.3	4.8	3.0	1.1	3.8	12.7
(n=1,653)	CI	85.6–88.9	3.9–6.0	2.2–3.9	0.7–1.8	2.9–4.8	11.1–14.4
Female	%	82.8	8.1	3.2	1.4	4.4	17.2
(n=1,629)	CI	80.9–84.6	6.9–9.6	2.4–4.2	1.0–2.2	3.5–5.5	15.4–19.1
White	%	84.9	7.0	3.2	1.2	3.6	15.1
(n=2,314)	CI	83.4–86.4	6.0–8.2	2.5–4.0	0.9–1.8	2.9–4.5	13.6–16.6
Black	%	88.5	3.9	1.5	1.1	4.9	11.5
(n=571)	CI	85.6–90.9	2.6–5.9	0.8–2.9	0.5–2.5	3.4–7.0	9.1–14.4
Hispanic	%	85.5	6.6	4.8	1.5	1.5	14.5
(n=64)	CI	74.0–92.4	2.4–16.9	1.5–14.3	0.2–10.7	0.2–10.7	7.6–26.0
8th	%	90.5	4.2	1.7	1.1	2.5	9.5
(n=1,287)	CI	88.8–92.0	3.2–5.4	1.1–2.6	0.6–1.8	1.8–3.5	8.0–11.2
10th	%	81.7	8.3	3.9	1.4	4.6	18.3
(n=1,172)	CI	79.4–83.8	6.9–10.1	2.9–5.2	0.9–2.3	3.5–6.0	16.2–20.6
12th	%	81.8	7.0	3.7	1.3	6.2	18.2
(n=935)	CI	79.2–84.2	5.5–8.8	2.7–5.1	0.7–2.2	4.8–7.9	15.8–20.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.57. Friends Tried Alcohol

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had one or more drinks of an alcoholic beverage?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	48.5	13.0	12.4	5.3	20.8	51.5
(n=3,405)	CI	46.8–50.2	11.9–14.2	11.4–13.6	4.6–6.1	19.5–22.3	49.8–53.2
Male	%	54.8	11.5	10.6	3.9	19.2	45.2
(n=1,656)	CI	52.3–57.2	10.0–13.1	9.2–12.2	3.1–5.0	17.3–21.2	42.8–47.7
Female	%	42.4	14.8	14.3	6.8	21.7	57.6
(n=1,638)	CI	40.0–44.8	13.2–16.6	12.7–16.1	5.6–8.1	19.7–23.8	55.2–60.0
White	%	44.0	13.8	14.1	5.8	22.2	56.0
(n=2,316)	CI	41.9–46.0	12.5–15.3	12.8–15.6	4.9–6.9	20.6–24.0	54.0–58.1
Black	%	66.3	9.8	7.4	3.1	13.4	33.7
(n=576)	CI	62.3–70.1	7.6–12.6	5.5–9.8	2.0–4.9	10.8–16.5	29.9–37.7
Hispanic	%	57.5	16.4	8.1	3.0	15.0	42.5
(n=65)	CI	44.8–69.2	9.2–27.7	3.3–18.4	0.7–11.8	7.8–26.7	30.8–55.2
8th	%	63.7	13.7	10.0	3.3	9.3	36.3
(n=1,285)	CI	61.0–66.3	11.9–15.7	8.5–11.8	2.5–4.5	7.8–11.0	33.7–39.0
10th	%	41.0	15.8	14.1	6.9	22.2	59.0
(n=1,185)	CI	38.2–43.8	13.9–18.0	12.2–16.2	5.6–8.5	19.9–24.7	56.2–61.8
12th	%	40.9	9.3	13.2	5.5	31.1	59.1
(n=935)	CI	37.8–44.0	7.6–11.4	11.1–15.5	4.2–7.2	28.2–34.1	56.0–62.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.58. Friends Binge Drinking

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have had five or more drinks on the same occasion?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	64.7	10.0	7.7	3.7	13.8	35.3
(n=3,399)	CI	63.1–66.4	9.0–11.0	6.8–8.7	3.1–4.5	12.7–15.1	33.6–36.9
Male	%	68.4	8.9	6.5	3.3	12.8	31.6
(n=1,653)	CI	66.1–70.7	7.6–10.4	5.4–7.9	2.5–4.3	11.2–14.6	29.3–33.9
Female	%	61.8	11.1	8.9	4.3	13.9	38.2
(n=1,634)	CI	59.4–64.1	9.7–12.8	7.6–10.4	3.4–5.4	12.3–15.7	35.9–40.6
White	%	60.7	11.3	9.2	4.0	14.9	39.3
(n=2,313)	CI	58.6–62.7	10.1–12.7	8.0–10.5	3.2–4.9	13.4–16.4	37.3–41.4
Black	%	79.2	6.0	3.8	2.7	8.3	20.8
(n=576)	CI	75.6–82.3	4.3–8.2	2.5–5.7	1.6–4.4	6.3–11.0	17.7–24.4
Hispanic	%	72.3	6.1	3.7	3.0	15.0	27.7
(n=65)	CI	59.5–82.2	2.2–15.6	0.9–13.9	0.7–11.8	7.8–26.7	17.8–40.5
8th	%	83.3	7.7	3.8	1.2	4.0	16.7
(n=1,286)	CI	81.2–85.2	6.4–9.3	2.9–5.0	0.8–2.0	3.0–5.2	14.8–18.8
10th	%	59.4	12.5	9.1	5.1	13.9	40.6
(n=1,178)	CI	56.5–62.1	10.8–14.5	7.6–10.9	4.0–6.5	12.0–16.0	37.9–43.5
12th	%	51.5	9.7	10.2	4.9	23.8	48.5
(n=935)	CI	48.3–54.7	7.9–11.7	8.4–12.3	3.7–6.5	21.2–26.6	45.3–51.7

Note: The confidence interval (CI) is 95 percent.

Table BH6.59. Friends Smoked Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have smoked part, or all of a cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	69.4	11.2	6.7	2.9	9.8	30.6
(n=3,402)	CI	67.8–70.9	10.2–12.3	5.9–7.6	2.4–3.5	8.9–10.9	29.1–32.2
Male	%	72.1	9.8	6.2	2.7	9.1	27.9
(n=1,663)	CI	69.9–74.3	8.5–11.4	5.2–7.6	2.0–3.6	7.8–10.6	25.7–30.1
Female	%	67.3	12.8	7.2	3.0	9.7	32.7
(n=1,629)	CI	65.0–69.6	11.3–14.5	6.0–8.6	2.3–4.0	8.3–11.3	30.4–35.0
White	%	65.4	12.8	7.7	3.4	10.7	34.6
(n=2,313)	CI	63.4–67.3	11.5–14.3	6.7–8.9	2.7–4.2	9.4–12.0	32.7–36.6
Black	%	84.8	6.5	3.0	0.8	4.9	15.2
(n=576)	CI	81.6–87.5	4.7–8.7	1.9–4.7	0.3–2.0	3.4–7.1	12.5–18.4
Hispanic	%	82.3	4.6	3.4	1.8	7.9	17.7
(n=65)	CI	70.4–90.1	1.4–13.6	0.8–12.9	0.2–12.4	3.3–18.1	9.9–29.6
8th	%	77.4	10.3	5.0	1.3	6.0	22.6
(n=1,287)	CI	75.0–79.6	8.8–12.1	3.9–6.3	0.8–2.1	4.8–7.4	20.4–25.0
10th	%	67.8	12.6	6.4	3.2	9.9	32.2
(n=1,177)	CI	65.1–70.4	10.8–14.7	5.2–8.0	2.4–4.4	8.3–11.8	29.6–34.9
12th	%	62.9	10.6	8.8	4.1	13.7	37.1
(n=938)	CI	59.7–65.9	8.8–12.7	7.1–10.8	3.0–5.6	11.6–16.0	34.1–40.3

Note: The confidence interval (CI) is 95 percent.

Table BH6.60. Friends Used Electronic Cigarettes

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used an e-cigarette?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	55.4	10.1	8.2	5.5	20.8	44.6
(n=3,401)	CI	53.7–57.1	9.2–11.2	7.3–9.1	4.8–6.3	19.4–22.2	42.9–46.3
Male	%	57.7	8.5	7.9	5.0	20.8	42.3
(n=1,657)	CI	55.3–60.1	7.2–9.9	6.7–9.4	4.0–6.2	18.9–22.9	39.9–44.7
Female	%	53.9	12.0	8.4	6.0	19.7	46.1
(n=1,634)	CI	51.5–56.3	10.5–13.7	7.1–9.9	4.9–7.3	17.8–21.7	43.7–48.5
White	%	49.1	11.4	9.5	6.8	23.3	50.9
(n=2,317)	CI	47.0–51.1	10.2–12.7	8.3–10.7	5.8–7.9	21.6–25.1	48.9–53.0
Black	%	82.3	4.7	2.7	1.9	8.4	17.7
(n=574)	CI	78.9–85.3	3.2–6.7	1.7–4.4	1.0–3.4	6.4–11.0	14.7–21.1
Hispanic	%	63.4	7.6	12.3	1.5	15.1	36.6
(n=65)	CI	50.6–74.6	3.1–17.5	6.2–23.2	0.2–10.5	7.9–26.9	25.4–49.4
8th	%	66.5	11.6	5.7	3.5	12.7	33.5
(n=1,287)	CI	63.9–69.1	10.0–13.5	4.5–7.1	2.6–4.6	10.9–14.6	30.9–36.1
10th	%	51.8	10.6	9.6	6.8	21.2	48.2
(n=1,179)	CI	48.9–54.6	8.9–12.5	8.1–11.5	5.5–8.4	19.0–23.7	45.4–51.1
12th	%	47.9	8.2	9.1	6.3	28.6	52.1
(n=935)	CI	44.7–51.1	6.6–10.1	7.4–11.1	4.9–8.0	25.8–31.5	48.9–55.3

Note: The confidence interval (CI) is 95 percent.

Table BH6.61. Friends Used Marijuana

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many, if any, of your friends have used marijuana or hashish?

		None	One friend	Two friends	Three friends	Four friends	Combined: (One or more)
Total	%	61.9	10.0	8.1	4.3	15.7	38.1
(n=3,409)	CI	60.2–63.5	9.1–11.1	7.2–9.1	3.6–5.0	14.5–17.0	36.5–39.8
Male	%	65.9	8.3	7.3	4.1	14.4	34.1
(n=1,661)	CI	63.5–68.2	7.1–9.8	6.1–8.7	3.2–5.2	12.8–16.2	31.8–36.5
Female	%	58.6	11.9	9.3	4.3	15.9	41.4
(n=1,637)	CI	56.2–61.0	10.4–13.6	7.9–10.8	3.4–5.4	14.2–17.8	39.0–43.8
White	%	62.2	10.8	8.4	4.3	14.4	37.8
(n=2,323)	CI	60.1–64.1	9.5–12.1	7.3–9.6	3.6–5.3	13.0–15.9	35.9–39.9
Black	%	66.9	6.9	5.8	3.2	17.2	33.1
(n=576)	CI	62.9–70.7	5.1–9.2	4.2–8.1	2.0–5.0	14.2–20.6	29.3–37.1
Hispanic	%	74.2	3.7	2.9	7.6	11.6	25.8
(n=65)	CI	61.7–83.8	0.9–13.9	0.7–11.2	3.1–17.5	5.5–22.8	16.2–38.3
8th	%	78.6	9.0	4.0	2.3	6.0	21.4
(n=1,295)	CI	76.3–80.8	7.6–10.7	3.1–5.2	1.6–3.3	4.8–7.5	19.2–23.7
10th	%	54.0	11.2	10.2	5.5	19.2	46.0
(n=1,180)	CI	51.1–56.8	9.5–13.1	8.6–12.0	4.3–7.0	17.1–21.6	43.2–48.9
12th	%	53.0	9.9	10.2	5.0	21.9	47.0
(n=934)	CI	49.8–56.2	8.1–12.0	8.4–12.3	3.8–6.6	19.4–24.7	43.8–50.2

Note: The confidence interval (CI) is 95 percent.

Table BH6.62. Perceived Risk of Misusing Prescription Drugs

How much do you think people risk harming themselves physically and in other ways when they use prescription drugs not prescribed to them?

		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.8	6.7	15.2	67.3	82.5
(n=3,390)	CI	9.8–11.9	5.9–7.6	14.0–16.5	65.7–68.9	81.2–83.8
Male	%	12.5	6.1	15.3	66.1	81.4
(n=1,657)	CI	11.0–14.2	5.1–7.4	13.6–17.1	63.8–68.3	79.4–83.2
Female	%	8.3	7.2	15.6	68.8	84.4
(n=1,622)	CI	7.1–9.8	6.1–8.6	13.9–17.5	66.5–71.1	82.6–86.1
White	%	6.7	6.4	16.0	70.8	86.9
(n=2,313)	CI	5.7–7.8	5.5–7.5	14.6–17.6	68.9–72.7	85.4–88.2
Black	%	25.7	6.9	12.8	54.7	67.4
(n=571)	CI	22.2–29.5	5.1–9.3	10.3–15.8	50.5–58.7	63.4–71.2
Hispanic	%	11.2	4.6	11.0	73.2	84.3
(n=63)	CI	5.3–22.1	1.4–13.6	5.2–21.9	60.6–83.0	72.7–91.5
8th	%	9.0	7.8	14.0	69.2	83.2
(n=1,285)	CI	7.6–10.7	6.4–9.4	12.2–16.0	66.6–71.7	81.1–85.2
10th	%	10.1	8.1	16.9	64.9	81.8
(n=1,173)	CI	8.5–12.0	6.6–9.8	14.9–19.2	62.1–67.6	79.5–83.9
12th	%	13.2	4.2	14.7	67.9	82.6
(n=932)	CI	11.2–15.5	3.1–5.6	12.6–17.1	64.8–70.8	80.1–84.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.63. Perceived Risk of Regularly Drinking Alcohol

		How much do you think people risk harming themselves physically and in other ways when they <u>take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	14.8	17.9	25.7	41.6	67.3
(n=3,408)	CI	13.7–16.1	16.6–19.2	24.2–27.2	39.9–43.2	65.7–68.8
Male	%	16.5	19.3	25.1	39.1	64.2
(n=1,668)	CI	14.8–18.4	17.5–21.3	23.1–27.3	36.8–41.5	61.9–66.5
Female	%	12.7	15.9	26.7	44.7	71.4
(n=1,626)	CI	11.1–14.4	14.2–17.8	24.6–28.9	42.2–47.1	69.1–73.6
White	%	11.6	19.3	27.5	41.6	69.1
(n=2,328)	CI	10.3–12.9	17.8–21.0	25.7–29.3	39.6–43.7	67.2–71.0
Black	%	26.4	12.8	18.1	42.6	60.7
(n=569)	CI	22.9–30.3	10.3–15.9	15.1–21.5	38.6–46.8	56.6–64.7
Hispanic	%	16.1	16.2	26.9	40.8	67.7
(n=65)	CI	9.0–27.3	8.8–27.9	17.4–39.2	29.2–53.5	55.1–78.2
8th	%	13.1	16.6	24.4	45.9	70.3
(n=1,292)	CI	11.3–15.0	14.7–18.8	22.2–26.9	43.2–48.6	67.7–72.7
10th	%	13.3	20.0	26.0	40.7	66.7
(n=1,180)	CI	11.5–15.3	17.8–22.4	23.5–28.5	38.0–43.6	64.0–69.3
12th	%	18.2	17.0	26.7	38.1	64.8
(n=936)	CI	15.9–20.8	14.7–19.5	24.0–29.6	35.0–41.2	61.7–67.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.64. Perceived Risk of Binge Drinking

		How much do you think people risk harming themselves physically and in other ways when they <u>take five or more drinks of an alcoholic beverage (beer, wine, liquor) once or twice a week?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	12.2	11.2	23.0	53.5	76.6
(n=3,404)	CI	11.1–13.4	10.2–12.4	21.6–24.5	51.8–55.2	75.1–78.0
Male	%	14.5	12.1	23.2	50.2	73.4
(n=1,661)	CI	12.9–16.3	10.6–13.7	21.2–25.3	47.8–52.7	71.2–75.5
Female	%	9.2	10.4	22.6	57.8	80.4
(n=1,629)	CI	7.9–10.7	9.0–12.0	20.6–24.7	55.3–60.2	78.3–82.3
White	%	8.5	11.7	25.6	54.2	79.8
(n=2,326)	CI	7.4–9.8	10.4–13.0	23.9–27.4	52.2–56.2	78.1–81.4
Black	%	26.2	8.4	15.0	50.4	65.4
(n=569)	CI	22.7–30.0	6.4–11.0	12.3–18.2	46.3–54.5	61.3–69.2
Hispanic	%	17.1	13.5	16.6	52.8	69.4
(n=65)	CI	9.5–28.7	7.0–24.5	9.2–28.0	40.4–65.0	56.8–79.7
8th	%	10.1	9.5	21.9	58.5	80.4
(n=1,287)	CI	8.6–11.9	8.1–11.3	19.7–24.2	55.7–61.1	78.1–82.4
10th	%	10.8	12.1	25.4	51.7	77.1
(n=1,178)	CI	9.1–12.7	10.4–14.1	23.0–28.0	48.9–54.6	74.7–79.4
12th	%	15.9	12.1	21.7	50.4	72.1
(n=939)	CI	13.7–18.3	10.1–14.3	19.2–24.4	47.2–53.6	69.1–74.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.65. Perceived Risk of Smoking Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke one or more packs of cigarettes a day?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	10.6	6.1	16.5	66.9	83.4
(n=3,413)	CI	9.6–11.7	5.3–6.9	15.3–17.8	65.3–68.5	82.1–84.6
Male	%	12.4	6.6	17.9	63.2	81.1
(n=1,669)	CI	10.8–14.0	5.5–7.9	16.1–19.8	60.8–65.5	79.1–82.9
Female	%	8.1	5.5	15.2	71.2	86.4
(n=1,630)	CI	6.8–9.5	4.5–6.7	13.6–17.1	68.9–73.4	84.7–88.0
White	%	6.7	6.2	17.8	69.2	87.0
(n=2,331)	CI	5.8–7.9	5.3–7.3	16.3–19.4	67.3–71.1	85.6–88.4
Black	%	24.5	6.4	11.1	58.0	69.1
(n=571)	CI	21.1–28.3	4.6–8.7	8.8–13.9	53.9–62.0	65.1–72.8
Hispanic	%	11.9	1.5	14.1	72.5	86.6
(n=65)	CI	5.9–22.4	0.2–10.5	7.6–24.8	60.3–82.1	75.8–93.0
8th	%	8.7	6.5	19.2	65.6	84.8
(n=1,296)	CI	7.3–10.4	5.3–8.0	17.1–21.4	63.0–68.2	82.8–86.7
10th	%	9.8	6.1	15.3	68.8	84.1
(n=1,180)	CI	8.2–11.6	4.9–7.6	13.4–17.5	66.1–71.4	81.9–86.1
12th	%	13.2	5.6	14.9	66.2	81.1
(n=937)	CI	11.2–15.6	4.3–7.3	12.8–17.4	63.1–69.2	78.5–83.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.66. Perceived Risk of Using Electronic Cigarettes

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke e-cigarettes, e-cigars, or e-hookahs?</u>				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	14.3	23.0	26.5	36.3	62.7
(n=3,403)	CI	13.1–15.5	21.6–24.4	25.0–28.0	34.7–37.9	61.1–64.4
Male	%	16.0	24.9	24.9	34.2	59.1
(n=1,659)	CI	14.3–17.8	22.9–27.1	22.9–27.1	31.9–36.5	56.7–61.4
Female	%	11.7	20.7	28.8	38.8	67.6
(n=1,630)	CI	10.2–13.4	18.8–22.7	26.6–31.1	36.5–41.2	65.3–69.9
White	%	11.0	24.7	28.1	36.2	64.3
(n=2,327)	CI	9.8–12.4	22.9–26.5	26.3–29.9	34.3–38.2	62.3–66.2
Black	%	27.3	16.4	18.1	38.2	56.3
(n=570)	CI	23.7–31.1	13.6–19.7	15.2–21.5	34.2–42.3	52.2–60.4
Hispanic	%	8.9	16.8	26.7	47.6	74.3
(n=64)	CI	3.9–19.0	9.4–28.3	17.0–39.3	35.3–60.1	61.9–83.7
8th	%	12.3	21.8	25.1	40.8	65.9
(n=1,291)	CI	10.6–14.2	19.7–24.2	22.8–27.5	38.1–43.5	63.2–68.4
10th	%	12.7	24.6	28.0	34.6	62.6
(n=1,174)	CI	11.0–14.8	22.3–27.2	25.5–30.7	31.9–37.3	59.8–65.3
12th	%	17.9	22.4	26.2	33.4	59.7
(n=938)	CI	15.6–20.5	19.9–25.2	23.5–29.1	30.5–36.5	56.5–62.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.67. Perceived Risk of Trying Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>try marijuana once or twice</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	33.2	22.3	14.9	29.7	44.5
(n=3,402)	CI	31.6–34.8	20.9–23.7	13.7–16.1	28.2–31.2	42.9–46.2
Male	%	34.6	21.9	14.6	28.9	43.5
(n=1,660)	CI	32.3–36.9	20.0–24.0	13.0–16.4	26.8–31.1	41.2–45.9
Female	%	30.4	23.2	15.6	30.8	46.4
(n=1,628)	CI	28.2–32.8	21.2–25.3	13.9–17.4	28.6–33.1	44.0–48.8
White	%	30.0	22.8	15.9	31.3	47.2
(n=2,323)	CI	28.2–32.0	21.1–24.5	14.4–17.4	29.5–33.2	45.1–49.2
Black	%	40.6	18.7	13.5	27.2	40.7
(n=568)	CI	36.6–44.8	15.7–22.1	10.9–16.6	23.7–31.0	36.7–44.8
Hispanic	%	28.0	29.7	10.5	31.8	42.3
(n=65)	CI	18.2–40.6	19.5–42.3	5.0–20.9	21.4–44.4	30.6–54.9
8th	%	20.4	22.2	16.2	41.2	57.4
(n=1,286)	CI	18.2–22.6	20.0–24.6	14.3–18.4	38.5–43.9	54.7–60.1
10th	%	34.4	24.8	16.4	24.3	40.8
(n=1,177)	CI	31.7–37.2	22.4–27.4	14.4–18.7	22.0–26.9	38.0–43.6
12th	%	44.8	19.8	11.8	23.6	35.4
(n=939)	CI	41.7–48.0	17.3–22.4	9.9–14.1	21.0–26.4	32.4–38.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.68. Perceived Risk of Regularly Smoking Marijuana

		How much do you think people risk harming themselves physically and in other ways when they <u>smoke marijuana once or twice a week</u> ?				
		No risk	Slight risk	Moderate risk	Great risk	Combined: (Great risk, Moderate risk)
Total	%	25.5	15.6	20.2	38.7	58.9
(n=3,398)	CI	24.0–27.0	14.4–16.9	18.9–21.6	37.1–40.4	57.3–60.6
Male	%	27.4	15.6	19.6	37.5	57.1
(n=1,662)	CI	25.3–29.6	13.9–17.4	17.7–21.6	35.2–39.8	54.6–59.4
Female	%	22.6	15.3	21.6	40.5	62.0
(n=1,623)	CI	20.6–24.8	13.6–17.2	19.6–23.6	38.1–42.9	59.6–64.4
White	%	21.3	14.7	21.7	42.3	64.0
(n=2,325)	CI	19.7–23.1	13.3–16.2	20.1–23.4	40.3–44.3	62.0–65.9
Black	%	37.3	16.4	16.3	30.0	46.3
(n=566)	CI	33.4–41.5	13.5–19.7	13.5–19.6	26.3–33.9	42.2–50.4
Hispanic	%	26.0	15.8	21.6	36.5	58.1
(n=65)	CI	16.6–38.4	8.6–27.4	13.0–33.7	25.5–49.2	45.5–69.8
8th	%	16.2	11.2	20.2	52.4	72.6
(n=1,288)	CI	14.3–18.3	9.6–13.0	18.1–22.5	49.7–55.2	70.1–75.0
10th	%	26.0	17.4	22.1	34.6	56.7
(n=1,176)	CI	23.5–28.5	15.3–19.7	19.8–24.5	31.9–37.4	53.8–59.5
12th	%	34.4	18.2	18.4	29.0	47.4
(n=934)	CI	31.5–37.5	15.8–20.8	16.0–21.0	26.2–32.0	44.2–50.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.69. Wrong to Misuse Prescription Drugs

How wrong do you think it is for someone your age to <u>use prescription drugs not prescribed to them?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	5.0	5.9	12.2	76.9	89.0
(n=3,408)	CI	4.3–5.8	5.2–6.8	11.1–13.3	75.4–78.3	87.9–90.1
Male	%	4.8	5.9	12.4	76.9	89.3
(n=1,665)	CI	3.9–6.0	4.9–7.2	10.9–14.1	74.7–78.8	87.7–90.7
Female	%	4.3	5.7	12.3	77.7	90.0
(n=1,628)	CI	3.4–5.4	4.7–7.0	10.8–14.0	75.5–79.6	88.4–91.3
White	%	3.9	5.9	12.0	78.2	90.2
(n=2,328)	CI	3.2–4.8	5.0–7.0	10.7–13.4	76.5–79.9	88.9–91.3
Black	%	6.4	4.4	11.0	78.2	89.2
(n=567)	CI	4.6–8.8	3.0–6.4	8.7–13.9	74.6–81.5	86.3–91.6
Hispanic	%	3.7	6.1	10.5	79.8	90.3
(n=65)	CI	0.9–13.9	2.2–15.6	5.0–20.9	67.7–88.1	79.5–95.7
8th	%	3.2	3.9	10.3	82.6	92.9
(n=1,294)	CI	2.4–4.4	2.9–5.1	8.7–12.0	80.5–84.6	91.4–94.2
10th	%	5.3	7.1	14.0	73.6	87.6
(n=1,178)	CI	4.2–6.8	5.8–8.7	12.1–16.1	71.0–76.0	85.5–89.3
12th	%	6.6	6.8	12.2	74.4	86.6
(n=936)	CI	5.2–8.3	5.3–8.6	10.3–14.4	71.6–77.1	84.3–88.7

Note: The confidence interval (CI) is 95 percent.

Table BH6.70. Wrong to Regularly Drink Alcohol

How wrong do you think it is for someone your age to <u>have one or two drinks of an alcoholic beverage nearly every day?</u>						
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	7.2	12.4	19.7	60.7	80.4
(n=3,418)	CI	6.4–8.1	11.3–13.6	18.4–21.1	59.1–62.4	79.0–81.7
Male	%	7.2	12.6	19.1	61.1	80.1
(n=1,667)	CI	6.1–8.6	11.1–14.4	17.2–21.0	58.7–63.4	78.1–82.0
Female	%	6.1	12.1	20.2	61.6	81.7
(n=1,636)	CI	5.0–7.4	10.6–13.9	18.3–22.2	59.2–63.9	79.8–83.6
White	%	6.5	13.1	20.2	60.2	80.4
(n=2,329)	CI	5.6–7.7	11.7–14.5	18.6–21.9	58.1–62.2	78.7–82.0
Black	%	7.1	8.7	14.4	69.9	84.3
(n=576)	CI	5.2–9.5	6.6–11.3	11.8–17.5	66.0–73.5	81.0–87.1
Hispanic	%	9.9	10.8	21.6	57.7	79.3
(n=65)	CI	4.4–20.8	5.1–21.4	13.0–33.7	45.0–69.4	67.1–87.8
8th	%	3.8	6.6	17.0	72.5	89.5
(n=1,298)	CI	2.9–5.0	5.4–8.1	15.1–19.2	70.0–74.9	87.7–91.1
10th	%	7.3	14.1	21.9	56.6	78.6
(n=1,182)	CI	6.0–9.0	12.2–16.2	19.7–24.4	53.8–59.4	76.1–80.8
12th	%	10.5	16.5	20.1	52.9	73.0
(n=938)	CI	8.7–12.6	14.3–19.0	17.7–22.8	49.7–56.1	70.1–75.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.71. Wrong to Smoke Tobacco

		How wrong do you think it is for someone your age to <u>smoke tobacco</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	8.0	10.2	17.6	64.2	81.8
(n=3,417)	CI	7.1–8.9	9.2–11.3	16.4–18.9	62.6–65.8	80.5–83.1
Male	%	8.7	11.0	16.9	63.5	80.4
(n=1,668)	CI	7.4–10.2	9.5–12.6	15.2–18.8	61.1–65.8	78.3–82.3
Female	%	6.0	9.5	18.3	66.2	84.5
(n=1,634)	CI	4.9–7.3	8.1–11.0	16.5–20.3	63.8–68.5	82.6–86.2
White	%	8.0	11.2	18.3	62.5	80.8
(n=2,330)	CI	6.9–9.2	9.9–12.6	16.8–20.0	60.5–64.5	79.1–82.4
Black	%	5.4	5.5	13.3	75.8	89.1
(n=575)	CI	3.8–7.6	3.9–7.8	10.7–16.4	72.0–79.1	86.2–91.4
Hispanic	%	10.2	7.8	11.0	71.1	82.0
(n=65)	CI	4.6–21.3	3.2–17.8	5.2–21.7	58.3–81.2	70.0–89.9
8th	%	3.9	5.4	15.3	75.4	90.7
(n=1,299)	CI	3.0–5.1	4.3–6.8	13.4–17.4	73.0–77.7	89.0–92.2
10th	%	6.3	10.9	19.0	63.7	82.7
(n=1,179)	CI	5.1–7.9	9.3–12.8	16.9–21.4	60.9–66.4	80.5–84.8
12th	%	13.7	14.5	18.5	53.3	71.8
(n=939)	CI	11.7–16.1	12.4–16.9	16.1–21.1	50.1–56.4	68.8–74.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.72. Wrong to Use Electronic Cigarettes

		How wrong do you think it is for someone your age to <u>use an e-cigarette</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	15.8	17.7	16.2	50.4	66.5
(n=3,414)	CI	14.5–17.0	16.5–19.1	14.9–17.4	48.7–52.1	64.9–68.1
Male	%	15.7	17.3	16.2	50.8	66.9
(n=1,665)	CI	14.0–17.6	15.6–19.3	14.5–18.0	48.3–53.2	64.6–69.2
Female	%	14.3	18.2	16.6	50.9	67.5
(n=1,634)	CI	12.7–16.1	16.4–20.2	14.9–18.5	48.4–53.3	65.2–69.8
White	%	16.7	19.6	16.4	47.3	63.7
(n=2,329)	CI	15.2–18.3	18.0–21.3	14.9–17.9	45.3–49.4	61.7–65.6
Black	%	8.8	10.2	13.1	67.8	81.0
(n=573)	CI	6.7–11.5	8.0–13.0	10.6–16.2	63.9–71.6	77.5–84.0
Hispanic	%	9.7	20.9	12.3	57.0	69.4
(n=65)	CI	4.3–20.5	12.3–33.1	6.1–23.2	44.4–68.9	56.6–79.7
8th	%	9.4	12.3	15.8	62.5	78.3
(n=1,299)	CI	7.9–11.1	10.6–14.2	13.9–17.9	59.9–65.1	76.0–80.5
10th	%	16.0	20.3	17.2	46.5	63.7
(n=1,176)	CI	14.0–18.2	18.1–22.7	15.2–19.5	43.6–49.3	60.9–66.4
12th	%	22.0	20.6	15.4	42.0	57.4
(n=939)	CI	19.5–24.7	18.1–23.3	13.3–17.9	38.9–45.1	54.2–60.5

Note: The confidence interval (CI) is 95 percent.

Table BH6.73. Wrong to Smoke Marijuana

		How wrong do you think it is for someone your age to <u>smoke marijuana</u> ?				
		Not at all wrong	A little bit wrong	Wrong	Very Wrong	Combined: (Very wrong, Wrong)
Total	%	16.2	12.3	13.6	58.0	71.5
(n=3,415)	CI	15.0–17.5	11.2–13.4	12.5–14.8	56.3–59.6	70.0–73.1
Male	%	16.4	11.6	13.1	59.0	72.0
(n=1,669)	CI	14.7–18.3	10.1–13.2	11.5–14.8	56.6–61.3	69.8–74.2
Female	%	15.1	12.6	14.1	58.2	72.4
(n=1,631)	CI	13.4–17.0	11.0–14.3	12.5–15.9	55.8–60.6	70.1–74.5
White	%	14.8	11.5	14.0	59.7	73.7
(n=2,331)	CI	13.3–16.3	10.3–12.9	12.7–15.5	57.6–61.7	71.8–75.5
Black	%	16.2	10.8	11.9	61.1	73.0
(n=574)	CI	13.3–19.5	8.5–13.7	9.5–14.8	57.0–65.1	69.2–76.6
Hispanic	%	13.3	17.2	9.0	60.6	69.5
(n=65)	CI	6.6–24.7	9.6–28.9	4.0–19.0	47.8–72.0	56.8–79.9
8th	%	6.5	7.1	11.6	74.8	86.4
(n=1,295)	CI	5.3–8.0	5.8–8.6	9.9–13.4	72.4–77.1	84.4–88.2
10th	%	18.4	14.3	15.7	51.6	67.3
(n=1,182)	CI	16.3–20.7	12.5–16.5	13.7–17.9	48.7–54.4	64.5–69.9
12th	%	23.8	15.3	13.4	47.4	60.8
(n=938)	CI	21.2–26.6	13.2–17.8	11.4–15.8	44.2–50.6	57.7–63.9

Note: The confidence interval (CI) is 95 percent.

Table BH6.74. Parents Talked about Prescription Drug Misuse

		During the past 12 months, have you talked with at least one of your parents about the dangers of using prescription drugs not prescribed to you? By parents, we mean your adult guardians, whether they live with you or not.			
		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	66.9	23.3	9.8	33.1
(n=3,407)	CI	65.3–68.5	21.9–24.8	8.8–10.8	31.5–34.7
Male	%	66.2	23.6	10.2	33.8
(n=1,663)	CI	63.9–68.5	21.6–25.7	8.8–11.8	31.5–36.1
Female	%	67.2	23.6	9.2	32.8
(n=1,631)	CI	64.8–69.4	21.6–25.7	7.9–10.7	30.6–35.2
White	%	67.0	23.7	9.3	33.0
(n=2,327)	CI	65.0–68.8	22.0–25.5	8.2–10.6	31.2–35.0
Black	%	69.1	20.7	10.2	30.9
(n=573)	CI	65.2–72.7	17.5–24.2	8.0–13.0	27.3–34.8
Hispanic	%	57.2	30.5	12.3	42.8
(n=65)	CI	44.5–68.9	20.3–43.1	6.1–23.2	31.1–55.5
8th	%	63.0	26.4	10.7	37.0
(n=1,292)	CI	60.3–65.5	24.0–28.8	9.1–12.5	34.5–39.7
10th	%	68.3	22.1	9.6	31.7
(n=1,179)	CI	65.6–70.9	19.8–24.5	8.1–11.5	29.1–34.4
12th	%	69.5	21.5	9.0	30.5
(n=936)	CI	66.4–72.3	19.0–24.3	7.3–11.0	27.7–33.6

Note: The confidence interval (CI) is 95 percent.

Table BH6.75. Parents Talked about Alcohol and Other Drugs (ATOD)

During the past 12 months, have you talked with at least one of your parents about the dangers of alcohol, tobacco, or drug use? By parents, we mean your adult guardians, whether they live with you or not.

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	47.7	33.3	19.0	52.3
(n=3,410)	CI	46.1–49.4	31.7–34.9	17.7–20.3	50.6–53.9
Male	%	48.8	32.0	19.2	51.2
(n=1,666)	CI	46.4–51.2	29.8–34.3	17.3–21.1	48.8–53.6
Female	%	46.2	35.1	18.7	53.8
(n=1,632)	CI	43.7–48.6	32.8–37.5	16.9–20.7	51.4–56.3
White	%	45.7	34.9	19.4	54.3
(n=2,331)	CI	43.7–47.7	33.0–36.9	17.8–21.0	52.3–56.3
Black	%	58.5	25.2	16.3	41.5
(n=571)	CI	54.4–62.5	21.8–29.0	13.5–19.5	37.5–45.6
Hispanic	%	42.0	44.0	14.0	58.0
(n=65)	CI	30.3–54.6	32.2–56.6	7.3–25.2	45.4–69.7
8th	%	44.7	34.6	20.7	55.3
(n=1,293)	CI	42.0–47.4	32.1–37.3	18.6–23.0	52.6–58.0
10th	%	45.3	33.3	21.4	54.7
(n=1,179)	CI	42.4–48.1	30.7–36.1	19.2–23.8	51.9–57.6
12th	%	53.4	32.0	14.6	46.6
(n=938)	CI	50.2–56.6	29.1–35.0	12.5–17.0	43.4–49.8

Note: The confidence interval (CI) is 95 percent.

Table BH6.76. Exposure to Prescription Drug Misuse Media Messaging

During the past 12 months, do you recall hearing, reading, or watching an advertisement about the dangers of using prescription drugs not prescribed to you?

		No	Yes, once	Yes, more than once	Combined: (Yes, once, Yes, more than once)
Total	%	40.5	30.6	28.9	59.5
(n=3,408)	CI	38.8–42.2	29.1–32.2	27.4–30.4	57.8–61.2
Male	%	41.8	28.4	29.7	58.2
(n=1,663)	CI	39.4–44.2	26.3–30.7	27.6–32.0	55.8–60.6
Female	%	38.8	33.1	28.1	61.2
(n=1,633)	CI	36.4–41.2	30.9–35.5	26.0–30.3	58.8–63.6
White	%	38.5	31.9	29.6	61.5
(n=2,329)	CI	36.5–40.5	30.0–33.8	27.8–31.5	59.5–63.5
Black	%	51.9	25.4	22.7	48.1
(n=573)	CI	47.8–56.0	21.9–29.1	19.5–26.3	44.0–52.2
Hispanic	%	38.5	31.9	29.6	61.5
(n=65)	CI	27.2–51.2	21.5–44.5	19.5–42.3	48.8–72.8
8th	%	31.4	31.4	37.2	68.6
(n=1,294)	CI	28.9–33.9	29.0–34.0	34.6–39.9	66.1–71.1
10th	%	43.9	28.9	27.3	56.1
(n=1,178)	CI	41.0–46.7	26.4–31.5	24.8–29.9	53.3–59.0
12th	%	46.3	31.6	22.1	53.7
(n=936)	CI	43.2–49.5	28.7–34.6	19.5–24.8	50.5–56.8

Note: The confidence interval (CI) is 95 percent.