Drug Overdoses in Tennessee

3,826 Tennesseans died due to a drug overdose in 2022.

Fentanyl-related deaths increased between 2017 and 2021 by 446%.

Emerging substances like **Xylazine**

a veterinary sedative, contributed to 324 overdose deaths in TN between 2019 and 2022.

How can drug overdoses be prevented?

Store & Dispose of Medications Correctly

Support Local Substance Use Prevention Coalitions



Participate in Drug Take-Back Events



High Impact Area (HIA)

programs are managed by Substance
Misuse Response Coordinators to
achieve a more comprehensive,
equitable, and locally managed
response to areas of TN most
burdened by the overdose epidemic.

See website for more information on the work of the West, Middle, Southeast, East, and Northeast HIAs managed by TDH's Overdose Response Coordination Office (ORCO).

To access ORCO's website and resources, scan code:



- 1.Fatal Drug Overdose Timeseries Trends. Tennessee State Government TN.gov. (n.d.). https://www.tn.gov/health/health-program-areas/pdo/pdo/data-dashboard.html
- Korona-Bailey, J., Onyango, E. M., Hall, K., Jayasundara, J. H., & Mukhopadhyay, S. (2023).
 Xylazine-Involved Fatal and Nonfatal Drug Overdoses in Tennessee From 2019 to 2022. JAMA Network Open, 6(7), e2324001-e2324001. https://doi.org/10.1001/jamanetworkopen.2023.24001

Paid for by the Tennessee Department of Health, Overdose Response Coordination Office and the Center for Disease Control and Prevention. Overdose Data to Action in States Grant NU17CE010208.



Tennessee Department of Health Authorization No. 355997, 6400 copies. May 2024. This public document was promulgated at a cost of \$0.21 per copy.



www.tn.gov/health/orco



You may be at increased risk of **OVERDOSE**

If you...

- Have had a previous overdose
- Have a substance use disorder
- Use more than one substance (aka polysubstance use)
- Are exposed to Fentanyl or Xylazine (aka "Trang") in your supply

Remember,

most overdoses are **ACCIDENTAL.**

OPIOID OVERDOSE

Warning Signs



Dizziness and disorientation



Blue lips

Choking or

gurgling sounds



Slow or No **Breathing**



Cold and/or clammy skin



Falling asleep or hard to wake up



Pinpoint Pupils

Reversal

If you suspect an Opioid Overdose...

IMMEDIATELY

- **USE NALOXONE** if you have it
- Call 911
- Keep the person awake
- Lay the person on their side to prevent choking
- Stay until help arrives

Where to get Naloxone:



- Local Health **Department** (limited supply)
- Local **Pharmacy**

For Overdose Prevention Training and Resources Contact:

Virtual Training:

https://apps.health.tn.gov/naloxone/savealife/

Agency or Individual:

TN.gov/behavioral-health/ROPS

Looking for Treatment?



Visit the website to find treatment for substance use disorders.



Call or text 24/7/365 for addiction treatment referrals.